

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/12, 5/10, C07K 14/705, 16/28, C12N 15/62, C07K 17/14, 19/00, G01N 33/545, A61K 38/17, 47/48	A2	(11) International Publication Number: WO 96/40915 (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09052 (22) International Filing Date: 6 June 1996 (06.06.96) (30) Priority Data: 08/479,744 7 June 1995 (07.06.95) US (71) Applicants: DANA-FARBER CANCER INSTITUTE [US/US]; 44 Binney Street, Boston, MA 02115 (US). REPLIGEN CORPORATION [US/US]; One Kendall Square, Cambridge, MA 02139 (US). (72) Inventors: FREEMAN, Gordon, J.; 305 Walnut Street, Brook- line, MA 02146 (US). NADLER, Lee, M.; 36 Cross Hill Road, Newton, MA 02159 (US). GRAY, Gary, S.; 32 Mil- ton Road, Brookline, MA 02146 (US). (74) Agents: MANDRAGOURAS, Amy, E. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: NOVEL CTLA4/CD28 LIGANDS AND USES THEREFOR		
(57) Abstract Nucleic acids encoding novel CTLA4/CD28 ligands which costimulate T cell activation are disclosed. In one embodiment, the nucleic acid has a sequence which encodes a fusion protein consisting of at least part of the B lymphocyte antigen, B7-2, fused to one or more constant domains of immunoglobulin heavy chain. Preferably, the nucleic acid is a DNA molecule comprising at least a portion of a nucleotide sequence shown in Figure 8, SEQ ID NO:1 or Figure 14, SEQ ID NO:23. The nucleic acid sequences of the invention can be integrated into various expression vectors, which in turn direct the synthesis of the corresponding proteins or peptides in a variety of hosts, particularly eukaryotic cells, such as mammalian and insect cell culture. Also disclosed are host cells transformed to produce proteins or peptides encoded by the nucleic acid sequences of the invention and isolated proteins and peptides which comprise at least a portion of a novel B lymphocyte antigen. Proteins and peptides described herein can be administered to subjects to enhance or suppress T cell-mediated immune responses.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

NOVEL CTLA4/CD28 LIGANDS AND USES THEREFOR

Background of the Invention

10

To induce antigen-specific T cell activation and clonal expansion, two signals provided by antigen-presenting cells (APCs) must be delivered to the surface of resting T lymphocytes (Jenkins, M. and Schwartz, R. (1987) *J. Exp. Med.* 165, 302-319; Mueller, D.L., et al. (1990) *J. Immunol.* 144, 3701-3709; Williams, I.R. and Unanue, E.R. (1990) *J. Immunol.* 145, 85-93). The first signal, which confers specificity to the immune response, is mediated via the T cell receptor (TCR) following recognition of foreign antigenic peptide presented in the context of the major histocompatibility complex (MHC). The second signal, termed costimulation, induces T cells to proliferate and become functional (Schwartz, R.H. (1990) *Science* 248, 1349-1356). Costimulation is neither antigen-specific, nor MHC restricted and is thought to be provided by one or more distinct cell surface molecules expressed by APCs (Jenkins, M.K., et al. (1988) *J. Immunol.* 140, 3324-3330; Linsley, P.S., et al. (1991) *J. Exp. Med.* 173, 721-730; Gimmi, C.D., et al., (1991) *Proc. Natl. Acad. Sci. USA* 88, 6575-6579; Young, J.W., et al. (1992) *J. Clin. Invest.* 90, 229-237; Koulova, L., et al. (1991) *J. Exp. Med.* 173, 759-762; Reiser, H., et al. (1992) *Proc. Natl. Acad. Sci. USA* 89, 271-275; van-Seventer, G.A., et al. (1990) *J. Immunol.* 144, 4579-4586; LaSalle, J.M., et al., (1991) *J. Immunol.* 147, 774-80; Dustin, M.I., et al., (1989) *J. Exp. Med.* 169, 503; Armitage, R.J., et al. (1992) *Nature* 357, 80-82; Liu, Y., et al. (1992) *J. Exp. Med.* 175, 437-445).

Considerable evidence suggests that the B7 protein, expressed on APCs, is one such critical costimulatory molecule (Linsley, P.S., et al., (1991) *J. Exp. Med.* 173, 721-730; Gimmi, C.D., et al., (1991) *Proc. Natl. Acad. Sci. USA* 88, 6575-6579; Koulova, L., et al., (1991) *J. Exp. Med.* 173, 759-762; Reiser, H., et al. (1992) *Proc. Natl. Acad. Sci. USA* 89, 271-275; Linsley, P.S. et al. (1990) *Proc. Natl. Acad. Sci. USA* 87, 5031-5035; Freeman, G.J. et al. (1991) *J. Exp. Med.* 174, 625-631.). B7 is the counter-receptor for two ligands

30

expressed on T lymphocytes. The first ligand, termed CD28, is constitutively expressed on resting T cells and increases after activation. After signaling through the T cell receptor, ligation of CD28 induces T cells to proliferate and secrete IL-2 (Linsley, P.S., et al. (1991) *J. Exp. Med.* 173, 721-730; Gimmi, C.D., et al. (1991) *Proc. Natl. Acad. Sci. USA.* 88, 6575-6579; Thompson, C.B., et al. (1989) *Proc. Natl. Acad. Sci. USA.* 86, 1333-1337; June, C.H., et al. (1990) *Immunol. Today.* 11, 211-6; Harding, F.A., et al. (1992) *Nature.* 356, 607-609.). The second ligand, termed CTLA4 is homologous to CD28 but is not expressed on resting T cells and appears following T cell activation (Brunet, J.F., et al., (1987) *Nature* 328, 267-270). DNA sequences encoding the human and murine CTLA4 protein are described in Dariavich, et al. (1988) *Eur. J. Immunol.* 18(12), 1901-1905; Brunet, J.F., et al. (1987) *supra*; Brunet, J.F. et al. (1988) *Immunol. Rev.* 103:21-36; and Freeman, G.J., et al. (1992) *J. Immunol.* 149, 3795-3801. Although B7 has a higher affinity for CTLA4 than for CD28 (Linsley, P.S., et al., (1991) *J. Exp. Med.* 174, 561-569), the function of CTLA4 is still unknown.

The importance of the B7:CD28/CTLA4 costimulatory pathway has been demonstrated *in vitro* and in several *in vivo* model systems. Blockade of this costimulatory pathway results in the development of antigen specific tolerance in murine and humans systems (Harding, F.A., et al. (1992) *Nature.* 356, 607-609; Lenschow, D.J., et al. (1992) *Science.* 257, 789-792; Turka, L.A., et al. (1992) *Proc. Natl. Acad. Sci. USA.* 89, 11102-11105; Gimmi, C.D., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90, 6586-6590; Boussiotis, V., et al. (1993) *J. Exp. Med.* 178, 1753-1763). Conversely, expression of B7 by B7 negative murine tumor cells induces T-cell mediated specific immunity accompanied by tumor rejection and long lasting protection to tumor challenge (Chen, L., et al. (1992) *Cell* 71, 1093-1102; Townsend, S.E. and Allison, J.P. (1993) *Science* 259, 368-370; Baskar, S., et al. (1993) *Proc. Natl. Acad. Sci.* 90, 5687-5690.). Therefore, manipulation of the B7:CD28/CTLA4 pathway offers great potential to stimulate or suppress immune responses in humans.

Summary of the Invention


This invention pertains to isolated nucleic acids encoding novel molecules which costimulate T cell activation. Preferred costimulatory molecules include antigens on the surface of B lymphocytes, professional antigen presenting cells (e.g., monocytes, dendritic cells, Langerhan cells) and other cells (e.g., keratinocytes, endothelial cells, astrocytes, fibroblasts, oligodendrocytes) which present antigen to immune cells, and which bind either CTLA4, CD28, both CTLA4 and CD28 or other known or as yet undefined receptors on immune cells. Such costimulatory molecules are referred to herein as CTLA4/CD28 binding counter-receptors or B lymphocyte antigens, and are capable of

providing costimulation to activated T cells to thereby induce T cell proliferation and/or cytokine secretion. Preferred B lymphocyte antigens include B7-2 and B7-3 and soluble fragments or derivatives thereof which bind CTLA4 and/or CD28 and have the ability to inhibit or induce costimulation of immune cells. In one embodiment, an isolated nucleic acid which encodes a peptide having the activity of the human B7-2 B lymphocyte antigen is provided. Preferably, the nucleic acid is a cDNA molecule having a nucleotide sequence encoding human B7-2, as shown in Figure 8 (SEQ ID NO:1). In another embodiment, the nucleic acid is a cDNA molecule having a nucleotide sequence encoding murine B7-2, as shown in Figure 14 (SEQ ID NO:22).

The invention also features nucleic acids which encode a peptide having B7-2 activity and at least about 50%, more preferably at least about 60% and most preferably at least about 70% homologous with an amino acid sequence shown in Figure 8 (SEQ ID NO:2) or an amino acid sequence shown in Figure 14 (SEQ ID NO:23). Nucleic acids which encode peptides having B7-2 activity and at least about 80%, more preferably at least about 90%, more preferably at least about 95% and most preferably at least about 98% or at least about 99% homologous with an amino acid sequence shown in Figure 8 (SEQ ID NO:2) or an amino acid sequence shown in Figure 14 (SEQ ID NO:23) are also within the scope of the invention. In another embodiment, the peptide having B7-2 activity is encoded by a nucleic acid which hybridizes under high or low stringency conditions to a nucleic acid which encodes a peptide having an amino acid sequence of Figure 8 (SEQ ID NO:2) or a peptide having an amino acid sequence shown in Figure 14 (SEQ ID NO:23).

The invention further pertains to an isolated nucleic acid comprising a nucleotide sequence encoding a peptide having B7-2 activity and having a length of at least 20 amino acid residues. Peptides having B7-2 activity and consisting of at least 40 amino acid residues in length, at least 60 amino acid residues in length, at least 80 amino acid residues in length, at least 100 amino acid residues in length or at least 200 or more amino acid residues in length are also within the scope of this invention. Particularly preferred nucleic acids encode a peptide having B7-2 activity, a length of at least 20 amino acid residues or more and at least 50% or greater homology (preferably at least 70%) with a sequence shown in Figure 8 (SEQ ID NO:2).

In one preferred embodiment, the invention features an isolated DNA encoding a peptide having B7-2 activity and an amino acid sequence represented by a formula:



In the formula, Y consists essentially of amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2). X_n and Z_m are additional amino acid residue(s) linked to Y by an amide bond. X_n and Z_m are amino acid residues selected from amino acid residues contiguous to Y in the amino acid sequence shown in Figure 8 (SEQ ID NO:2). X_n is amino acid residue(s) selected from amino acids contiguous to the amino terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2), i.e., selected from amino acid residue 23 to 1. Z_m is amino acid residue(s) selected from amino acids contiguous to the carboxy terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2), i.e., selected from amino acid residue 246 to 329. According to the formula, n is a number from 0 to 23 ($n=0-23$) and m is a number from 0 to 84 ($m=0-84$). A particularly preferred DNA encodes a peptide having an amino acid sequence represented by the formula X_n -Y- Z_m , where Y is amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2) and $n=0$ and $m=0$.

The invention also features an isolated DNA encoding a B7-2 fusion protein which includes a nucleotide sequence encoding a first peptide having B7-2 activity and a nucleotide sequence encoding a second peptide corresponding to a moiety that alters the solubility, binding affinity, stability or valency of the first peptide. Preferably, the first peptide having B7-2 activity includes an extracellular domain portion of the B7-2 protein (e.g., about amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2)) and the second peptide is an immunoglobulin constant region, for example, a human C γ 1 or C γ 4 domain, including the hinge, CH2 and CH3 region, to produce a B7-2 immunoglobulin fusion protein (B7-2Ig)(see Capon et al. (1989) *Nature* 337, 525-531 and Capon U.S. 5,116,964).

The nucleic acids obtained in accordance with the present invention can be inserted into various expression vectors, which in turn direct the synthesis of the corresponding protein or peptides in a variety of hosts, particularly eucaryotic cells, such as mammalian and insect cell culture, and procaryotic cells such as *E. coli*. Expression vectors within the scope of the invention comprise a nucleic acid encoding at least one peptide having the activity of a novel B lymphocyte antigen as described herein, and a promoter operably linked to the nucleic acid sequence. In one embodiment, the expression vector contains a DNA encoding a peptide having the activity of the B7-2 antigen and a DNA encoding a peptide having the activity of another B lymphocyte antigen, such as the previously characterized B7 activation antigen, referred to herein as B7-1. Such expression vectors can be used to transfect host cells to thereby produce proteins and peptides, including fusion proteins, encoded by nucleic acids as described herein.

Nucleic acid probes useful for assaying a biological sample for the presence of B cells expressing the B lymphocyte antigens B7-2 and B7-3 are also within the scope of the invention.

The invention further pertains to isolated peptides having the activity of a novel B lymphocyte antigen, including the B7-2 and B7-3 protein antigens. A preferred peptide having B7-2 activity is produced by recombinant expression and comprises an amino acid sequence shown in Figure 8 (SEQ ID NO: 2). Another preferred peptide having B7-2 activity comprises an amino acid sequence shown in Figure 14 (SEQ ID NO:23). A particularly preferred peptide having the activity of the B7-2 antigen includes at least a portion of the mature form of the protein, such as an extracellular domain portion (e.g., about amino acid residues 24-245 of SEQ ID NO:2) which can be used to enhance or suppress T-cell mediated immune responses in a subject. Other preferred peptides having B7-2 activity include peptides having an amino acid sequence represented by a formula:

$$X_n-Y-Z_m$$

In the formula, Y is amino acid residues selected from the group consisting of: amino acid residues 55-68 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 81-89 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 128-142 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 160-169 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 188-200 of the sequence shown in Figure 8 (SEQ ID NO:2); and amino acid residues 269-282 of the sequence shown in Figure 8 (SEQ ID NO:2). In the formula X_n and Z_m are additional amino acid residue(s) linked to Y by an amide bond and are selected from amino acid residues contiguous to Y in the amino acid sequence shown in Figure 8 (SEQ ID NO:2). X_n is amino acid residue(s) selected from amino acids contiguous to the amino terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2). Z_m is amino acid residue(s) selected from amino acids contiguous to the carboxy terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2). According to the formula, n is a number from 0 to 30 ($n=0-30$) and m is a number from 0 to 30 ($m=0-30$).

Fusion proteins or hybrid fusion proteins including a peptide having the activity of a novel B lymphocyte antigen (e.g., B7-2, B7-3) are also featured. For example, a fusion protein comprising a first peptide which includes an extracellular domain portion of a novel B lymphocyte antigen fused to second peptide, such as an immunoglobulin constant region, that alters the solubility, binding affinity, stability and/or valency of the first peptide are provided. In one embodiment, a fusion protein is produced comprising a first peptide which includes amino acid residues of an extracellular domain portion of the B7-2 protein joined to a second peptide which includes amino acid residues of a sequence corresponding to the hinge, CH2 and CH3 regions of C γ 1 or C γ 4 to form a B7-2Ig fusion protein. In another embodiment, a hybrid fusion protein is produced comprising a first peptide which includes an extracellular domain portion of the B7-1 antigen and an extracellular domain portion of the

B7-2 antigen and a second peptide which includes amino acid residues corresponding to the hinge, CH2 and CH3 of C γ 1 (see e.g., Linsley et al. (1991) *J. Exp. Med.* 1783:721-730; Capon et al. (1989) *Nature* 337, 525-531; and Capon U.S. 5,116,964). In a yet another embodiment, a hybrid fusion protein comprises the immunoglobulin-like variable domain of B7-2, but not the immunoglobulin-like constant domain of B7-2, linked to the constant region of an immunoglobulin molecule. In a preferred embodiment, the B7-2Ig fusion protein includes the variable domain of human B7-2, preferably from about amino acid residue 24 to about amino acid residue 133 of the human B7-2 protein (as shown SEQ ID NO: 2), fused to the constant region of an IgG molecule.

Isolated peptides and fusion proteins of the invention can be administered to a subject to either upregulate or inhibit the expression of one or more B lymphocyte antigens or block the ligation of one or more B lymphocyte antigens to their natural ligand on immune cells, such as T cells, to thereby provide enhancement or suppression of cell-mediated immune responses *in vivo*.

Another embodiment of the invention provides antibodies, preferably monoclonal antibodies, specifically reactive with a peptide of a novel B lymphocyte antigen or fusion protein as described herein. Preferred antibodies are anti-human B7-2 monoclonal antibodies produced by hybridoma cells HF2.3D1, HA5.2B7 and HA3.1F9. These hybridoma cells have been deposited with the American Type Culture Collection at ATCC Accession No. HB 11686 (HF2.3D1), ATCC Accession No. HB 11687 (HA5.2B7), and ATCC Accession No. HB 11688 (HA3.1F9).

A still further aspect of the invention involves the use of the nucleic acids of the invention, especially the cDNAs, to enhance the immunogenicity of a mammalian cell. In preferred embodiments, the mammalian cell is a tumor cell, such as a sarcoma, a lymphoma, a melanoma, a neuroblastoma, a leukemia or a carcinoma, or an antigen presenting cell, such as a macrophage, which is transfected to allow expression of a peptide having the activity of a novel B lymphocyte antigen of the invention on the surface of the cell. Macrophages that express a peptide having the activity of a B lymphocyte antigen, such as the B7-2 antigen, can be used as antigen presenting cells, which, when pulsed with an appropriate pathogen-related antigen or tumor antigen, enhance T cell activation and immune stimulation.

Mammalian cells can be transfected with a suitable expression vector containing a nucleic acid encoding a peptide having the activity of a novel B lymphocyte antigen, such as the B7-2 antigen, *ex vivo* and then introduced into the host mammal, or alternatively, cells can be transfected with the gene *in vivo* via gene therapy techniques. For example, the nucleic acid encoding a peptide having B7-2 activity can be transfected alone, or in combination with nucleic acids encoding other costimulatory molecules. In enhancing the immunogenicity of tumors which do not express Class I or Class II MHC molecules, it

may be beneficial to additionally transfect appropriate class I or II genes into the mammalian cells to be transfected with a nucleic acid encoding a peptide having the activity of a B lymphocyte antigen, as described herein.

The invention also provides methods for inducing both general
5 immunosuppression and antigen-specific tolerance in a subject by, for example, blocking the functional interaction of the novel B lymphocyte antigens of the invention, e.g., B7-2 and B7-3, to their natural ligand(s) on T cells or other immune system cells, to thereby block co-stimulation through the receptor-ligand pair. In one embodiment, inhibitory molecules that can be used to block the interaction of the natural human B7-2 antigen to its natural ligands
10 (e.g., CTLA4 and CD28) include a soluble peptide having B7-2 binding activity but lacking the ability to costimulate immune cells, antibodies that block the binding of B7-2 to its ligands and fail to deliver a co-stimulatory signal (so called "blocking antibodies", such as blocking anti-B7-2 antibodies), B7-2-Ig fusion proteins, which can be produced in accordance with the teachings of the present invention, as well as soluble forms of B7-2
15 receptors, such as CTLA4Ig or CD28Ig. Such blocking agents can be used alone or in combination with agents which block interaction of other costimulatory molecules with their natural ligands (e.g., anti-B7 antibody). Inhibition of T cell responses and induction of T cell tolerance according to the methods described herein may be useful prophylactically, in preventing transplantation rejection (solid organ, skin and bone marrow) and graft versus host
20 disease, especially in allogeneic bone marrow transplantation. The methods of the invention may also be useful therapeutically, in the treatment of autoimmune diseases, allergy and allergic reactions, transplantation rejection, and established graft versus host disease in a subject.

Another aspect of the invention features methods for upregulating immune
25 responses by delivery of a costimulatory signal to T cells through use of a stimulatory form of B7-2 antigen, which include soluble, multivalent forms of B7-2 protein, such as a peptide having B7-2 activity and B7-2 fusion proteins. Delivery of a stimulatory form of B7-2 in conjunction with antigen may be useful prophylactically to enhance the efficacy of vaccination against a variety of pathogens and may also be useful therapeutically to
30 upregulate an immune response against a particular pathogen during an infection or against a tumor in a tumor-bearing host.

The invention also features methods of identifying molecules which can inhibit either the interaction of B lymphocyte antigens, e.g., B7-2, B7-3, with their receptors or interfere with intracellular signalling through their receptors. Methods for identifying
35 molecules which can modulate the expression of B lymphocyte antigens on cells are also provided. In addition, methods for identifying cytokines produced in response to

costimulation of T cells by novel B lymphocyte antigens are within the scope of the invention.

Brief Description of the Drawings

5 *Figure 1A-B* are graphic representations of the responses of CD28⁺ T cells, as assessed by ³H-thymidine incorporation or IL-2 secretion, to costimulation provided by either B7 (B7-1) transfected CHO cells (panel a) or syngeneic activated B lymphocytes (panel b) cultured in media, anti-CD3 alone, or anti-CD3 in the presence of the following monoclonal antibodies or recombinant proteins: αB7 (133, anti-B7-1); CTLA4Ig; Fab α
10 CD28; control Ig fusion protein (isotype control for CTLA4Ig); or αB5 (anti-B5, the isotype control for anti-B7-1).

Figure 2A-C are graphs of log fluorescence intensity of cell surface expression of B7-1 on splenic B cells activated with surface immunoglobulin (sIg) crosslinking. The total (panel a), B7-1 positive (B7-1⁺, panel b) and B7-1 negative (B7-1⁻, panel c) activated B
15 cells were stained with anti-B7-1 monoclonal antibody (133) and fluorescein isothiocyanate (FITC) labeled goat anti-mouse immunoglobulin and analyzed by flow cytometry.

Figure 3A-B are graphic representations of the responses of CD28⁺ T cells, as assessed by ³H-thymidine incorporation and IL-2 secretion, to costimulation provided by B7-1⁺ (panel a) or B7-1⁻ (panel b) activated syngeneic B lymphocytes cultured in media, anti-CD3 alone, or anti-CD3 in the presence of the following monoclonal antibodies or
20 recombinant proteins: αBB-1 (133, anti-B7-1 and anti-B7-3); αB7 (anti-B7-1); CTLA4Ig; Fab αCD28; control Ig fusion protein or αB5 (anti-B5).

Figure 4 is a graphic representation of the cell surface expression of B7-1, B7-3 and total CTLA4 counter-receptors on fractionated B7-1⁺ and B7-1⁻ activated B
25 lymphocytes.

Figure 5 is a graphic representation of temporal surface expression of B7-1 (CTLA4Ig and mAbs BB-1 and 133), B7-3 (CTLA4Ig and mAb BB1) and B7-2 (CTLA4Ig) counter-receptors on splenic B cells activated by sIg crosslinking.

Figure 6 is a graphic representation of temporal surface expression of B7-1 (CTLA4Ig and mAbs BB-1 and 133), B7-3 (CTLA4Ig and mAb BB1) and B7-2 (CTLA4Ig) counter-receptors on splenic B cells activated by MHC class II crosslinking.
30

Figure 7A-B are graphic representations of the response of CD28⁺ T cells, as assessed by ³H-thymidine incorporation and IL-2 secretion, to costimulation provided by syngeneic B lymphocytes activated by sIg crosslinking for 24 hours (panel a) or 48 hours (panel b) and cultured in media, anti-CD3 alone, or anti-CD3 in the presence of the following monoclonal antibodies or recombinant protein: αB7(133, anti-B7-1); αBB1 (anti-B7-1, anti-B7-3) CTLA4Ig; Fab αCD28; and αB5(anti-B5).
35

Figure 8 is the nucleotide and deduced amino acid sequence of the human B lymphocyte antigen B7-2 (hB7-2-clone29).

Figure 9 is a graphic representation of COS cells transfected with control plasmid (pCDNAI), plasmid expressing B7-1 (B7-1), or plasmid expressing B7-2 (B7-2) stained with either control mAb (IgM), anti-B7-1 (mAbs 133 and BB-1), recombinant protein CTLA4Ig, or isotype matched control Ig protein followed by the appropriate second FITC labelled immunoglobulin and analyzed by flow cytometry.

Figure 10A-B show RNA blot analyses of B7-2 expression in unstimulated and anti-Ig activated human splenic B cells and cell lines (panel a) and human myelomas (panel b).

Figure 11 is a graphic representation of the proliferation of CD28⁺ T cells, as assessed by ³H-thymidine incorporation or IL-2 secretion, to submitogenic stimulation with phorbol myristic acid (PMA) and COS cells transfected with vector alone or vectors directing the expression of either B7-1 or B7-2.

Figure 12 is a graphic representation of the inhibition by mAbs and recombinant proteins of the proliferation of CD28⁺ T cells, as assessed by ³H-thymidine incorporation and IL-2 secretion, to stimulation by PMA and COS cells transfected with vector alone (vector), or with a vector expressing B7-1 (B7-1) or B7-2 (B7-2). Inhibition studies were performed with the addition of either no antibody (no mAb), anti-B7 mAb 133 (133), anti-B7 mAb BB-1 (BB1), anti-B5 mAb (B5), Fab fragment of anti-CD28 (CD28 Fab), CTLA4Ig (CTLA4Ig), or Ig control protein (control Ig) to the PMA stimulated COS cell admixed CD28⁺ T cells.

Figure 13 shows the sequence homology between the human B7-2 protein (h B7-2) deduced amino acid sequence (SEQ ID NO: 2) and the amino acid sequence of both the human B7-1 protein (h B7-1) (SEQ ID NO: 28 and 29) and the murine B7-1 protein (m B7) (SEQ ID NO: 30 and 31).

Figure 14 is the nucleotide and deduced amino acid sequence of the murine B7-2 antigen (mB7-2) (SEQ ID NO: 22 and 23).

Figure 15 is a graphic representation of the competitive inhibition of binding of biotinylated-CTLA4Ig to immobilized B7-2 Ig by B7 family-Ig fusion proteins. The Ig fusion proteins examined as competitors were: full-length B7-2 (hB7.2), full-length B7-1 (hB7.1), the variable region-like domain of B7-2 (hB7.2V) or the constant region-like domain of B7-2 (hB7.2C).

Figure 16A-B are graphic representations of the competitive inhibition of binding of biotinylated-B7-1-Ig (panel A) or B7-2-Ig (panel B) to immobilized CTLA4-Ig by increasing concentrations of unlabelled B7-1-Ig (panel A) or B7-2-Ig (panel B). The experimentally determined IC₅₀ values are indicated in the upper right corner of the panels.

Figure 17 depicts flow cytometric profiles of cells stained with an anti-hB7-2 monoclonal antibody, HA3.1F9. Cells stained with the antibody were CHO cells transfected to express human B7-2 (CHO-hB7.2), NIH 3T3 cells transfected to express human B7-2 (3T3-hB7.2) and control transfected NIH 3T3 cells (3T3-neo). The anti-hB7.2 antibody B70
5 was used as a positive control.

Figure 18 depicts flow cytometric profiles of cells stained with an anti-hB7-2 monoclonal antibody, HA5.2B7. Cells stained with the antibody were CHO cells transfected to express human B7-2 (CHO-hB7.2), NIH 3T3 cells transfected to express human B7-2 (3T3-hB7.2) and control transfected NIH 3T3 cells (3T3-neo). The anti-hB7.2 antibody B70
10 was used as a positive control.

Figure 19 depicts flow cytometric profiles of cells stained with an anti-hB7-2 monoclonal antibody, HF2.3D1. Cells stained with the antibody were CHO cells transfected to express human B7-2 (CHO-hB7.2), NIH 3T3 cells transfected to express human B7-2 (3T3-hB7.2) and control transfected NIH 3T3 cells (3T3-neo). The anti-hB7.2 antibody B70
15 was used as a positive control.

Figure 20 is a graphic representation of the direct binding of soluble biotinylated CTLA4Ig to B7-1Ig, B7-1VIg, B7-1CIg, B7-2Ig, B7-2VIg, B7-2CIg, or human IgG (hIgG) bound to plates.

Figures 21A-E depict flow cytometric profiles of binding of B7-2Ig (Panel C), B7-2VIg (Panel D), B7-1Ig (Panel E), or secondary antibody alone (Panel B) to CTLA4+
20 CHO cells. Panel A is a negative control representing untransfected CHO cells.

Figure 22 depicts flow cytometric profiles of binding of control Ig, B7-1Ig, B7-2Ig, B7-2VIg, and anti-CD28 to CHO cells expressing CD28.

Figure 23 represents a histogram showing proliferation of CD28+ T cells
25 stimulated with 1 ng/ml PMA alone or with either of the following costimulatory signals: CHO/B7-1 cells, CHO/B7-2 cells, control Ig (30µg/ml), or B7-1Ig, B7-2Ig, or B7-2VIg (30µg or 100µg/ml each).

Figure 24 represents a histogram showing proliferation of, and IL-2 production by CD28+ T cells incubated with anti-CD3 attached to plates and B7-1Ig (10, 3 or
30 1µg/ml) or B7-2Ig (19, 3 or 1µg/ml) or B7-2VIg (3.0 - 0.01µg/ml).

Figure 25 represents the amount of IL-2 produced by CD28+ T cells after 1, 2, or 3 days of incubation of the cells with anti-CD3 alone or together with either CHO/B7-2 cells or B7-2VIg fusion protein.

Figure 26 represents the amount of IL-2 secreted by CD28+ T cells after 1, 2,
35 or 5 days of incubation of the cells with anti-CD3 alone or with either anti-CD28, B7-1Ig, B7-2Ig, or B7-2VIg.

Figure 27 is a graphical representation of the growth of CD28+ T cells incubated with anti-CD3 alone, or with B7-1Ig, B7-2Ig, B7-2VIg, or together with either anti-CD28.

5 Detailed Description of the Invention

In addition to the previously characterized B lymphocyte activation antigen B7 (referred to herein as B7-1), human B lymphocytes express other novel molecules which costimulate T cell activation. These costimulatory molecules include antigens on the surface of B lymphocytes, professional antigen presenting cells (e.g., monocytes, dendritic cells,
10 Langerhan cells) and other cells (e.g., keratinocytes, endothelial cells, astrocytes, fibroblasts, oligodendrocytes) which present antigen to immune cells, and which bind either CTLA4, CD28, both CTLA4 and CD28 or other known or as yet undefined receptors on immune cells. Costimulatory molecules within the scope of the invention are referred to herein as CTLA4/CD28 ligands (counter-receptors) or B lymphocyte antigens. Novel B lymphocyte
15 antigens which provide cotimulation to activated T cells to thereby induce T cell proliferation and/or cytokine secretion include the B7-2 (human and murine) and the B7-3 antigens described and characterized herein.

The B lymphocyte antigen B7-2 is expressed by human B cells at about 24 hours following stimulation with either anti-immunoglobulin or anti-MHC class II
20 monoclonal antibody. The B7-2 antigen induces detectable IL-2 secretion and T cell proliferation. At about 48 to 72 hours post activation, human B cells express both B7-1 and a third CTLA4 counter-receptor, B7-3, identified by a monoclonal antibody BB-1, which also binds B7-1 (Yokochi, T., et al. (1982) *J. Immunol.* 128, 823-827). The B7-3 antigen is also expressed on B7-1 negative activated B cells and can costimulate T cell proliferation without
25 detectable IL-2 production, indicating that the B7-1 and B7-3 molecules are distinct. B7-3 is expressed on a wide variety of cells including activated B cells, activated monocytes, dendritic cells, Langerhan cells and keratinocytes. At 72 hours post B cell activation, the expression of B7-1 and B7-3 begins to decline. The presence of these costimulatory molecules on the surface of activated B lymphocytes indicates that T cell costimulation is
30 regulated, in part, by the temporal expression of these molecules following B cell activation.

Accordingly, one aspect of this invention pertains to isolated nucleic acids comprising a nucleotide sequence encoding a novel costimulatory molecule, such as the B lymphocyte antigen, B7-2, fragments of such nucleic acids, or equivalents thereof. The term "nucleic acid" as used herein is intended to include such fragments or equivalents. The term
35 "equivalent" is intended to include nucleotide sequences encoding functionally equivalent B lymphocyte antigens or functionally equivalent peptides having an activity of a novel B lymphocyte antigen, i.e., the ability to bind to the natural ligand(s) of the B lymphocyte

antigen on immune cells, such as CTLA4 and/or CD28 on T cells, and inhibit (e.g., block) or stimulate (e.g., enhance) immune cell costimulation. Such nucleic acids are considered equivalents of the human B7-2 nucleotide sequence provided in Figure 8 (SEQ ID NO:1) and the murine B7-2 nucleotide sequence provided in Figure 14 (SEQ ID NO:22) and are within
5 the scope of this invention.

In one embodiment, the nucleic acid is a cDNA encoding a peptide having an activity of the B7-2 B lymphocyte antigen. Preferably, the nucleic acid is a cDNA molecule consisting of at least a portion of a nucleotide sequence encoding human B7-2, as shown in Figure 8 (SEQ ID NO:1) or at least a portion of a nucleotide sequence encoding murine B7-2,
10 as shown in Figure 14 (SEQ ID NO:22). A preferred portion of the cDNA molecule of Figure 8 (SEQ ID NO:1) or Figure 14 (SEQ ID NO:22) includes the coding region of the molecule.

In another embodiment, the nucleic acid of the invention encodes a peptide having an activity of B7-2 and comprising an amino acid sequence shown in Figure 8 (SEQ ID NO:2) or Figure 14 (SEQ ID NO:23). Preferred nucleic acids encode a peptide having B7-2 activity and at least about 50% homology, more preferably at least about 60% homology and most preferably at least about 70% homology with an amino acid sequence shown in Figure 8 (SEQ ID NO:2). Nucleic acids which encode peptides having B7-2 activity and at least about 90%, more preferably at least about 95%, and most preferably at least about 98-99% homologous with a sequence set forth in Figure 8 (SEQ ID NO:2) are
20 also within the scope of the invention. Homology refers to sequence similarity between two peptides having the activity of a novel B lymphocyte antigen, such as B7-2, or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequences is occupied by the same nucleotide base or amino acid, then the
25 molecules are homologous at that position. A degree (or percentage) of homology between sequences is a function of the number of matching or homologous positions shared by the sequences.

Another aspect of the invention provides a nucleic acid which hybridizes
30 under high or low stringency conditions to a nucleic acid which encodes a peptide having all or a portion of an amino acid sequence shown in Figure 8 (SEQ ID NO:2) or a peptide having all or a portion of an amino acid sequence shown in Figure 14 (SEQ ID NO:23). Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C are
35 known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50°C to a high


stringency of about 0.2 x SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C to high stringency conditions, at about 65°C.

Isolated nucleic acids encoding a peptide having an activity of a novel B lymphocyte antigen, as described herein, and having a sequence which differs from nucleotide sequence shown in Figure 8 (SEQ ID NO:1) or Figure 14 (SEQ ID NO:22) due to degeneracy in the genetic code are also within the scope of the invention. Such nucleic acids encode functionally equivalent peptides (e.g., a peptide having B7-2 activity) but differ in sequence from the sequence of Figure 8 or Figure 14 due to degeneracy in the genetic code. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid, or synonyms (for example, CAU and CAC are synonyms for histidine) may occur due to degeneracy in the genetic code. As one example, DNA sequence polymorphisms within the nucleotide sequence of a B7-2 (especially those within the third base of a codon) may result in "silent" mutations in the DNA which do not affect the amino acid encoded. However, it is expected that DNA sequence polymorphisms that do lead to changes in the amino acid sequences of the B7-2 antigen will exist within a population. It will be appreciated by one skilled in the art that these variations in one or more nucleotides (up to about 3-4% of the nucleotides) of the nucleic acids encoding peptides having the activity of a novel B lymphocyte antigen may exist among individuals within a population due to natural allelic variation. Any and all such nucleotide variations and resulting amino acid polymorphisms are within the scope of the invention. Furthermore, there may be one or more isoforms or related, cross-reacting family members of the novel B lymphocyte antigens described herein. Such isoforms or family members are defined as proteins related in function and amino acid sequence to a B lymphocyte antigen (e.g., the B7-2 antigen), but encoded by genes at different loci.

A "fragment" of a nucleic acid encoding a novel B lymphocyte antigen is defined as a nucleotide sequence having fewer nucleotides than the nucleotide sequence encoding the entire amino acid sequence of the B lymphocyte antigen and which encodes a peptide having an activity of the B lymphocyte antigen (i.e., the ability to bind to the natural ligand(s) of the B lymphocyte antigen on immune cells, such as CTLA4 and/or CD28 on T cells and either stimulate or inhibit immune cell costimulation). Thus, a peptide having B7-2 activity binds CTLA4 and/or CD28 and stimulates or inhibits a T cell mediated immune response, as evidenced by, for example, cytokine production and/or T cell proliferation by T cells that have received a primary activation signal. In one embodiment, the nucleic acid fragment encodes a peptide of the B7-2 antigen which retains the ability of the antigen to bind CTLA4 and/or CD28 and deliver a costimulatory signal to T lymphocytes. In another embodiment, the nucleic acid fragment encodes a peptide including an extracellular portion

of the human B7-2 antigen (e.g., approximately amino acid residues 24-245 of the sequence provided in Figure 8 (SEQ ID NO:2)) which can be used to bind CTLA4 and/or CD28 and, in monovalent form, inhibit costimulation, or in multivalent form, induce or enhance costimulation.

- 5 Preferred nucleic acid fragments encode peptides of at least 20 amino acid residues in length, preferably at least 40 amino acid residues and length, and more preferably at least 60 amino acid residues in length. Nucleic acid fragments which encode peptides of at least 80 amino acid residues in length, at least 100 amino acid residues in length, and at least 200 or more amino acids in length are also within the scope of the invention. Particularly
10 preferred nucleic acid fragments encode a peptide having the activity of human B7-2 and an amino acid sequence represented by a formula:



- 15 In the formula, Y comprises amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2). X_n and Z_m are additional amino acid residue(s) linked to Y by an amide bond. X_n and Z_m are selected from amino acid residues contiguous to Y in the amino acid sequence shown in Figure 8 (SEQ ID NO:2). In the formula, X_n is amino acid residue(s) selected from amino acids contiguous to the amino terminus of Y in the sequence shown in
20 Figure 8 (SEQ ID NO:2), i.e., from amino acid residue 23 to 1. Z_m is amino acid residue(s) selected from amino acids contiguous to the carboxy terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2), i.e., from amino acid residue 246 to 329. In addition, in the formula, n is a number from 0 to 23 ($n=0-23$) and m is a number from 0 to 84 ($m=0-84$). A particularly preferred peptide has an amino acid sequence represented by the formula X_n-Y-Z_m
25 as above, where $n=0$ and $m=0$.

- Nucleic acid fragments within the scope of the invention include those capable of hybridizing with nucleic acid from other animal species for use in screening protocols to detect novel proteins that are cross-reactive with the B lymphocyte antigens described herein. These and other fragments are described in detail herein. Generally, the nucleic acid
30 encoding a fragment of a B lymphocyte antigen will be selected from the bases coding for the mature protein, however, in some instances it may be desirable to select all or part of a fragment or fragments from the leader sequence or non-coding portion of a nucleotide sequence. Nucleic acids within the scope of the invention may also contain linker sequences, modified restriction endonuclease sites and other sequences useful for molecular cloning,
35 expression or purification of recombinant protein or fragments thereof. These and other modifications of nucleic acid sequences are described in further detail herein.

A nucleic acid encoding a peptide having an activity of a novel B lymphocyte antigen, such as the B7-2 antigen, may be obtained from mRNA present in activated B lymphocytes. It should also be possible to obtain nucleic acid sequences encoding B lymphocyte antigens from B cell genomic DNA. For example, the gene encoding the B7-2 antigen can be cloned from either a cDNA or a genomic library in accordance with protocols herein described. A cDNA encoding the B7-2 antigen can be obtained by isolating total mRNA from an appropriate cell line. Double stranded cDNAs can then prepared from the total mRNA. Subsequently, the cDNAs can be inserted into a suitable plasmid or viral (e.g., bacteriophage) vector using any one of a number of known techniques. Genes encoding novel B lymphocyte antigens can also be cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention. The nucleic acids of the invention can be DNA or RNA. A preferred nucleic acid is a cDNA encoding the human B7-2 antigen having the sequence depicted in Figure 8 (SEQ ID NO:1). Another preferred nucleic acid is a cDNA encoding the murine B7-2 antigen having the sequence shown on Figure 14 (SEQ ID NO:22).

This invention further pertains to expression vectors containing a nucleic acid encoding at least one peptide having the activity of a novel B lymphocyte antigen, as described herein, operably linked to at least one regulatory sequence. "Operably linked" is intended to mean that the nucleotide acid sequence is linked to a regulatory sequence in a manner which allows expression of the nucleotide sequence (e.g., in cis or trans). Regulatory sequences are art-recognized and are selected to direct expression of the desired protein in an appropriate host cell. Accordingly, the term regulatory sequence includes promoters, enhancers and other expression control elements. Such regulatory sequences are known to those skilled in the art or one described in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transfected and/or the type of protein desired to be expressed. In one embodiment, the expression vector includes a nucleic acid encoding at least a portion of the B7-2 protein, such as an extracellular domain portion. In another embodiment, the expression vector includes a DNA encoding a peptide having an activity of the B7-2 antigen and a DNA encoding a peptide having an activity of another B lymphocyte antigen, such as B7-1. cDNAs encoding the human B7-1 and mouse B7-1 antigens are shown in SEQ ID NO:28 and SEQ ID NO:30, respectively. The deduced amino acid sequences of these antigens are also shown in SEQ ID NO:29 and SEQ ID NO:31, respectively. Such expression vectors can be used to transfect cells to thereby produce proteins or peptides, including fusion proteins or peptides encoded by nucleic acid sequences as described herein. These and other embodiments are described in further detail herein.

The invention also features methods of producing peptides having an activity of a novel B lymphocyte antigen. For example, a host cell transfected with a nucleic acid vector directing expression of a nucleotide sequence encoding a peptide having an activity of the B7-2 protein can be cultured in a medium under appropriate conditions to allow
5 expression of the peptide to occur. In addition, one or more expression vectors containing DNA encoding a peptide having an activity of B7-2 and DNA encoding another peptide, such as a peptide having an activity of a second B lymphocyte antigen (e.g., B7-1, B7-3) can be used to transfect a host cell to coexpress these peptides or produce fusion proteins or peptides. In one embodiment, a recombinant expression vector containing DNA encoding a
10 B7-2 fusion protein is produced. A B7-2 fusion protein can be produced by recombinant expression of a nucleotide sequence encoding a first peptide having B7-2 activity and a nucleotide sequence encoding second peptide corresponding to a moiety that alters the solubility, affinity, stability or valency of the first peptide, for example, an immunoglobulin constant region. Preferably, the first peptide consists of a portion of the extracellular domain
15 of the human B7-2 antigen (e.g., approximately amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2)). The second peptide can include an immunoglobulin constant region, for example, a human C γ 1 domain or C γ 4 domain (e.g., the hinge, CH2 and CH3 regions of human IgC γ 1, or human IgC γ 4, see e.g., Capon et al. US 5,116,964, incorporated herein by reference). A resulting B7-2Ig fusion protein may have altered B7-2
20 solubility, binding affinity, stability and/or valency (i.e., the number of binding sites available per molecule) and may increase the efficiency of protein purification. Fusion proteins and peptides produced by recombinant technique may be secreted and isolated from a mixture of cells and medium containing the protein or peptide. Alternatively, the protein or peptide may be retained cytoplasmically and the cells harvested, lysed and the protein isolated. A cell
25 culture typically includes host cells, media and other byproducts. Suitable mediums for cell culture are well known in the art. Protein and peptides can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins and peptides. Techniques for transfecting host cells and purifying proteins and peptides are described in further detail herein.

30 Particularly preferred human B7-2Ig fusion proteins include the extracellular domain portion or variable region-like domain of human B7-2 coupled to an immunoglobulin constant region. The immunoglobulin constant region may contain genetic modifications which reduce or eliminate effector activity inherent in the immunoglobulin structure. For example, DNA encoding the extracellular portion of human B7-2 (hB7-2), as well as DNA
35 encoding the variable region-like domain of human B7-2 (hB7.2V) or the constant region-like domain of human B7-2 (hB7.2C) can be joined to DNA encoding the hinge, CH2 and

CH3 regions of human IgC γ 1 and/or IgC γ 4 modified by site directed mutagenesis. The preparation and characterization of these fusion proteins is described in detail in Example 7.

In a specific embodiment, the protein of the invention is a variable region form of the B cell activation antigen B7-2. The language "a variable region form of the B cell activation antigen B7-2" is intended to include forms of B7-2 which contain the immunoglobulin-like variable domain of B7-2, but do not comprise the immunoglobulin-like constant domain of B7-2. In a preferred embodiment, the variable region form of B7-2 comprises an amino acid sequence starting at about amino acid position 18 to 30 and ending about amino acid position 128 to 140 of human B7-2 protein (SEQ ID NO: 2). In a most preferred embodiment, the variable form of B7-2 comprises about amino acids 24 to 133 of human B7-2 protein (SEQ ID NO: 2). The variable region form of B7-2 can further be operatively linked directly to a transmembrane domain, such as the transmembrane domain of B7-2, to form a variable region form of B7-2 that can be expressed on a cell surface. "Operatively" is intended to mean in such a way that the molecule formed by operatively linking two or more domains or peptides is functional. The transmembrane domain of human B7-2 comprises about amino residues 246 to 268 of human B7-2 protein. Thus, in one embodiment, the variable region form of B7-2 is operatively linked to a peptide having a first amino acid located between about amino acid residue 238 and about amino acid residue 252, and a last amino acid residue located between about amino acid residue 260 and about amino acid residue 274 of human B7-2 protein of sequence SEQ ID NO: 2. The incorporation of a transmembrane domain in a protein of the invention, allows the protein to be expressed on a cell surface when a nucleic acid encoding the protein is expressed in the cell.

In another embodiment, the variable region form of B7-2 is operatively linked to a cytoplasmic domain, such as a cytoplasmic domain of B7-2. The cytoplasmic domain of human B7-2 comprises about amino acid residues 269 to 329 of human B7-2 protein of SEQ ID NO: 2. Accordingly, in one embodiment, the variable region form of B7-2 is operatively linked to a second B7-2 peptide, having a first amino acid residue located between about amino acids 260 and 275 of human B7-2 and a terminal amino acid residue located between about amino acid 323 to about amino acid 335 of human B7-2 of SEQ ID NO: 2. In another embodiment, a variable region form of B7-2 is operatively linked to a second B7-2 peptide of about amino acid residues 269 to 329 of human B7-2.

In a further embodiment, a variable region form of B7-2 operatively linked to a peptide corresponding to about the transmembrane domain of B7-2 is further operatively linked to a peptide corresponding significantly to the cytoplasmic domain of B7-2. Thus, proteins within the scope of the invention include those comprising an amino acid sequence from about position 24 to about position 133 of SEQ ID NO: 2, operatively linked to an

amino acid sequence from about position 246 to about position 268 of SEQ ID NO: 2 (V-region and transmembrane domains). Other proteins within the scope of the invention include those comprising an amino acid sequence from about position 24 to about position 133 of SEQ ID NO: 2, operatively linked to an amino acid sequence from about position 246 to about position 329 of SEQ ID NO: 2 (V-region, transmembrane and cytoplasmic domains). Yet other proteins within the scope of the invention include the leader sequence of B7-2 (e.g. positions 1-23) at the N-terminus. Proteins including other portions of B7-2 protein operatively linked to each other, but not including the immunoglobulin-like constant domain of B7-2, are also within the scope of the invention. In other embodiments, B7-2 proteins that contain an immunoglobulin-like constant domain of B7-2 in the absence of the variable region are contemplated.

A variable region form of B7-2, can also be linked to at least one heterologous polypeptide. The term "heterologous polypeptide" is intended to include any polypeptide, such as a polypeptide that directs the protein of the invention to a specific cellular compartment. In one embodiment, the heterologous polypeptide is a signal peptide that allows for the protein to be secreted from the cell. Another heterologous polypeptide within the scope of the invention is a signal peptide that allows for the protein to be expressed on the surface of the cell. In yet another embodiment, the heterologous polypeptide is a constant region of an immunoglobulin molecule. In an even more preferred embodiment, the heterologous polypeptide comprises the hinge, CH2, and CH3 domains of IgG1, as described herein.

The variable region form of B7-2 can further be attached to a linker polypeptide. A "linker polypeptide" as defined herein includes any polypeptide that bridges two peptides in the protein of the invention. Alternatively, the linker peptide is attached to either or both ends of the protein. Thus, a linker peptide attached to one or both ends of the protein can for example facilitate binding of the protein of the invention to a solid support. The linker peptide can also be a fragment of a bacterial or viral protein.

The fusion proteins described above can be, for example, human or murine. The nucleic acid molecules encoding the above described fusion proteins, as well as expression vectors and host cells expressing the fusion proteins are also within the scope of the invention.

Transfected cells which express peptides having an activity of one or more B lymphocyte antigens (e.g., B7-2, B7-3) on the surface of the cell are also within the scope of this invention. In one embodiment, a host cell such as a COS cell is transfected with an expression vector directing the expression of a peptide having B7-2 activity on the surface of the cell. Such a transfected host cell can be used in methods of identifying molecules which inhibit binding of B7-2 to its counter-receptor on T cells or which interfere with intracellular

signaling of costimulation to T cells in response to B7-2 interaction. In another embodiment, a tumor cell such as a sarcoma, a melanoma, a leukemia, a lymphoma, a carcinoma or a neuroblastoma is transfected with an expression vector directing the expression of at least one peptide having the activity of a novel B lymphocyte antigen on the surface of the tumor cell.


5 In some instances, it may be beneficial to transfect a tumor cell to coexpress major histocompatibility complex (MHC) proteins, for example MHC class II α and β chain proteins or an MHC class I α chain protein, and, if necessary, a β 2 microglobulin protein. Such transfected tumor cells can be used to induce tumor immunity in a subject. These and other embodiments are described in further detail herein.

10 The nucleic acid sequences of the invention can also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura et al. U.S. Patent No. 4,598,049; Caruthers et al. U.S. Patent No. 4,458,066; and
15 Itakura U.S. Patent Nos. 4,401,796 and 4,373,071, incorporated by reference herein).

Another aspect of the invention pertains to isolated peptides having an activity of a novel B lymphocyte antigen (e.g., B7-2, B7-3). A peptide having an activity of a B lymphocyte antigen may differ in amino acid sequence from the B lymphocyte antigen, such as the human B7-2 sequence depicted in Figure 8 (SEQ ID NO:2), or murine B7-2 sequence
20 depicted in Figure 14 (SEQ ID NO:22), but such differences result in a peptide which functions in the same or similar manner as the B lymphocyte antigen or which has the same or similar characteristics of the B lymphocyte antigen. For example, a peptide having an activity of the B7-2 protein is defined herein as a peptide having the ability to bind to the natural ligand(s) of the B7-2 protein on immune cells, such as CTLA4 and/or CD28 on T
25 cells and either stimulate or inhibit immune cell costimulation. Thus, a peptide having B7-2 activity binds CTLA4 and/or CD28 and stimulates or inhibits a T cell mediated immune response (as evidenced by, for example, cytokine production and/or proliferation by T cells that have received a primary activation signal). One embodiment provides a peptide having B7-2 binding activity, but lacking the ability to deliver a costimulatory signal to T cells.
30 Such a peptide can be used to inhibit or block T cell proliferation and/or cytokine secretion in a subject. Alternatively, a peptide having both B7-2 binding activity and the ability to deliver a costimulatory signal to T cells is used to stimulate or enhance T cell proliferation and/or cytokine secretion in a subject. Various modifications of the B7-2 protein to produce these and other functionally equivalent peptides are described in detail herein. The term "peptide"
35 as used herein, refers to peptides, proteins and polypeptides.

A peptide can be produced by modification of the amino acid sequence of the human B7-2 protein shown in Figure 8 (SEQ ID NO:2) or the murine B7-2 protein shown in

Figure 14 (SEQ ID NO:23), such as a substitution, addition or deletion of an amino acid residue which is not directly involved in the function of B7-2 (i.e., the ability of B7-2 to bind CTLA4 and/or CD28 and/or stimulate or inhibit T cell costimulation). Peptides of the invention are typically at least 20 amino acid residues in length, preferably at least 40 amino acid residues in length, and most preferably 60 amino acid residues in length. Peptides having B7-2 activity and including at least 80 amino acid residues in length, at least 100 amino acid residues in length, or at least 200 or more amino acid residues in length are also within the scope of the invention. A preferred peptide includes an extracellular domain portion of the human B7-2 antigen (e.g., about amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2). Other preferred peptides have an amino acid sequence represented by a formula:



where Y is amino acid residues selected from the group consisting of: amino acid residues 55-68 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 81-89 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 128-142 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 160-169 of the sequence shown in Figure 8 (SEQ ID NO:2); amino acid residues 188-200 of the sequence shown in Figure 8 (SEQ ID NO:2); and amino acid residues 269-282 of the sequence shown in Figure 8 (SEQ ID NO:2). In the formula, X_n and Z_m are additional amino acid residues linked to Y by an amide bond. X_n and Z_m are amino acid residues selected from amino acids contiguous to Y in the amino acid sequence shown in Figure 8 (SEQ ID NO:2). X_n is amino acid residues selected from amino acids contiguous to the amino terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2). Z_m is amino acid residues selected from amino acids contiguous to the carboxy terminus of Y in the sequence shown in Figure 8 (SEQ ID NO:2). According to the formula, n is a number from 0 to 30 ($n=0-30$) and m is a number from 0 to 30 ($m=0-30$). A particularly preferred peptide has an amino acid sequence represented by the formula X_n-Y-Z_m , where $n=0$ and $m=0$.

Another embodiment of the invention provides a substantially pure preparation of a peptide having an activity of a novel B lymphocyte antigen such as B7-2 or B7-3. Such a preparation is substantially free of proteins and peptides with which the peptide naturally occurs in a cell or with which it naturally occurs when secreted by a cell.

The term "isolated" as used throughout this application refers to a nucleic acid, protein or peptide having an activity of a novel B lymphocyte antigen, such as B7-2, substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized.

An isolated nucleic acid is also free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the organism from which the nucleic acid is derived.

The various peptides, polypeptides, proteins, and fusion proteins of the invention can be prepared as soluble forms. Alternatively, the proteins of the invention can be expressed on the surface of a cell, such as a CHO cell and can be prepared according to methods well known in the art. The proteins of the invention can also be coupled to a solid phase support, such as a bead or a plate. In a specific embodiment, a B7-2Ig fusion protein is attached to a solid phase support, such as a bead, for example a biodegradable bead. In a preferred embodiment, a variable region form of B7-2 is attached to a solid support. In a most preferred embodiment, a B7-2VIg fusion protein comprising an amino acid sequence of about position 24 to 133 of human B7-2Ig (SEQ ID NO: 2) linked to the constant domain of an IgG molecule is attached to a solid phase support.

These molecules can then be attached to a solid phase surface via several possible methods. For example, the proteins of the invention, such as a variable region form of B7-2, can be crosslinked to the beads via covalent modification using tosyl linkage. In this method, the proteins of the invention are typically in 0.05M borate buffer, pH 9.5 and added to tosyl-activated magnetic immunobeads (Dynal Inc., Great Neck, NY) according to manufacturer's instructions. After a 24 hr incubation at 22°C, the beads are collected and washed extensively. It is not mandatory that immunomagnetic beads be used, as other methods are also satisfactory. For example, proteins of the invention may also be immobilized on polystyrene beads or culture vessel surfaces.

It is also possible to attach the proteins, such as B7-2VIg to a solid phase surface through an avidin- or streptavidin-biotin complex. In this particular embodiment, the soluble protein is first crosslinked to biotin and then reacted with the solid phase surface to which avidin or streptavidin molecules are bound. It is also possible to crosslink the protein with avidin or streptavidin and to react these with a solid phase surface that is covered with biotin molecules.

These and other aspects of this invention are described in detail in the following subsections.

I. Isolation of Nucleic Acid From Cell Lines

Suitable cells for use in isolating nucleic acids encoding peptides having an activity of a novel B lymphocyte antigen include cells capable of producing mRNA coding for B lymphocyte antigens (e.g., B7-1, B7-2, B7-3) and appropriately translating the mRNA into the corresponding protein. One source of mRNA is normal human splenic B cells, either resting or activated by treatment with an anti-immunoglobulin antibody or an anti-MHC class

II antibody, or from subsets of neoplastic B cells. Expression of the human B7-2 antigen is detectable in resting B cells and in activated B cells, with mRNA levels increasing 4-fold from resting levels following stimulation. Total cellular RNA can be obtained using standard techniques from resting or activated B cells during these intervals and utilized in the construction of a cDNA library.

In addition, various subsets of neoplastic B cells may express B7-2 and B7-3 and can alternatively serve as a source of the mRNA for construction of a cDNA library. For example, tumor cells isolated from patients with non-Hodgkins lymphoma express B7-1 mRNA. B cells from nodular, poorly differentiated lymphoma (NPDL), diffuse large cell lymphoma (LCL) and Burkitt's lymphoma cell lines are also suitable sources of human B7-1 mRNA and, potentially B7-2 and B7-3 mRNA. Myelomas generally express B7-2, but not B7-1 mRNA, and, thus can provide a source of B7-2 mRNA. The Burkitt's lymphoma cell line Raji is one source of B lymphocyte antigen mRNA. Preferably, B7-2 mRNA is obtained from a population of both resting and activated normal human B cells. Activated B cells can be obtained by stimulation over a broad spectrum of time (e.g., from minutes to days) with, for example, an anti-immunoglobulin antibody or an anti-MCH class II antibody.

II. Isolation of mRNA and Construction of cDNA Library

Total cellular mRNA can be isolated by a variety of techniques, e.g., by using the guanidinium-thiocyanate extraction procedure of Chirgwin et al., *Biochemistry* **18**, 5294-5299 (1979). According to this method, Poly (A+) mRNA is prepared and purified for use in a cDNA library construction using oligo (dT) cellulose selection. cDNA is then synthesized from the poly(A+) RNA using oligo(dT) priming and reverse transcriptase. Moloney MLV reverse transcriptase (available from Gibco/BRL, Bethesda, MD) or AMV reverse transcriptase (available from Seikagaku America, Inc., St. Petersburg, FL) are preferably employed.

Following reverse transcription, the mRNA/DNA hybrid molecule is converted to double stranded DNA using conventional techniques and incorporated into a suitable vector. The experiments herein employed *E. coli* DNA polymerase I and ribonuclease H in the conversion to double stranded cDNA.

Cloning of the cDNAs can be accomplished using any of the conventional techniques for joining double stranded DNA with an appropriate vector. The use of synthetic adaptors is particularly preferred, since it alleviates the possibility of cleavage of the cDNA with restriction enzyme prior to cloning. Using this method, non-self complementary, kinased adaptors are added to the DNA prior to ligation with the vector. Virtually any adaptor can be employed. As set forth in more detail in the examples below, non-self

complementary BstXI adaptors are preferably added to the cDNA for cloning, for ligation into a pCDM8 vector prepared for cloning by digestion with BstXI.

Eucaryotic cDNA can be expressed when placed in the sense orientation in a vector that supplies an appropriate eucaryotic promoter and origin of replication and other elements including enhancers, splice acceptors and/or donor sequences and polyadenylation signals. The cDNAs of the present invention are placed in suitable vectors containing a eucaryotic promoter, an origin of replication functional in *E. coli*, an SV40 origin of replication which allows growth in COS cells, and a cDNA insertion site. Suitable vectors include π H3 (Seed and Aruffo, *Proc. Natl. Acad. Sci.*, 84:3365-3369 (1987)), π H3m (Aruffo and Seed, *Proc. Natl. Acad. Sci.*, 84:8573-8577 (1987)), pCDM7 and pCDM8 (Seed, *Nature*, 329:840-841 (1987), with the pCDM8 vector being particularly preferred (available commercially from Invitrogen, San Diego, CA).

III. Transfection of Host Cells and Screening for Novel B Lymphocyte Activation Antigens

The thus prepared cDNA library is then used to clone the gene of interest by expression cloning techniques. A basic expression cloning technique has been described by Seed and Aruffo, *Proc. Natl. Acad. Sci. USA*, 84:3365-3369 (1987) and Aruffo and Seed, *Proc. Natl. Acad. Sci. USA*, 84:8573-8577 (1987), although modifications to this technique may be necessary.

According to one embodiment, plasmid DNA is introduced into a simian COS cell line (Gluzman, *Cell* 23:175 (1981)) by known methods of transfection (e.g., DEAE-Dextran) and allowed to replicate and express the cDNA inserts. The transfectants expressing B7-1 antigen are depleted with an anti-B7-1 monoclonal antibody (e.g., 133 and B1.1) and anti-murine IgG and IgM coated immunomagnetic beads. Transfectants expressing human B7-2 antigen can be positively selected by reacting the transfectants with the fusion proteins CTLA4Ig and CD28Ig, followed by panning with anti-human Ig antibody coated plates. Although human CTLA4Ig and CD28Ig fusion proteins were used in the examples described herein, given the cross-species reactivity between B7-1 and, for example murine B7-1, it can be expected that other fusion proteins reactive with another cross-reactive species could be used. After panning, episomal DNA is recovered from the panned cells and transformed into a competent bacterial host, preferably *E. coli*. Plasmid DNA is subsequently reintroduced into COS cells and the cycle of expression and panning repeated at least two times. After the final cycle, plasmid DNA is prepared from individual colonies, transfected into COS cells and analyzed for expression of novel B lymphocyte antigens by indirect immunofluorescence with, for example, CTLA4Ig and CD28Ig.

IV. Sequencing of Novel B Lymphocyte Antigens

Plasmids are prepared from those clones which are strongly reactive with the CTLA4Ig and/or CD28Ig. These plasmids are then sequenced. Any of the conventional sequencing techniques suitable for sequencing tracts of DNA about 1.0 kb or larger can be employed.

As described in Example 4, a human B7-2 clone (clone29) was obtained containing an insert of 1,120 base pairs with a single long open reading frame of 987 nucleotides and approximately 27 nucleotides of 3' noncoding sequences (Figure 8, SEQ ID NO:1). The predicted amino acid sequence encoded by the open reading frame of the protein is shown below the nucleotide sequence in Figure 8. The encoded human B7-2 protein, is predicted to be 329 amino acid residues in length (SEQ ID NO:2). This protein sequence exhibits many features common to other type I Ig superfamily membrane proteins. Protein translation is predicted to begin at the methionine codon (ATG, nucleotides 107 to 109) based on the DNA homology in this region with the consensus eucaryotic translation initiation site (see Kozak, M. (1987) *Nucl. Acids Res.* 15:8125-8148). The amino terminus of the B7-2 protein (amino acids 1 to 23) has the characteristics of a secretory signal peptide with a predicted cleavage between the alanines at positions 23 and 24 (von Heijne (1987) *Nucl. Acids Res.* 14:4683). Processing at this site would result in a B7-2 membrane bound protein of 306 amino acids having an unmodified molecular weight of approximately 34 kDa. This protein would consist of an approximate extracellular Ig superfamily V and C like domains of from about amino acid residue 24 to 245, a hydrophobic transmembrane domain of from about amino acid residue 246 to 268, and a long cytoplasmic domain of from about amino acid residue 269 to 329. The homologies to the Ig superfamily are due to the two contiguous Ig-like domains in the extracellular region bound by the cysteines at positions 40 to 110 and 157 to 218. The extracellular domain also contains eight potential N-linked glycosylation sites and, like B7-1, is probably glycosylated. Glycosylation of the human B7-2 protein may increase the molecular weight to about 50-70 kDa. The cytoplasmic domain of human B7-2, while somewhat longer than B7-1, contains a common region of multiple cysteines followed by positively charged amino acids which presumably function as signaling or regulatory domains within an antigen-presenting cell (APC). Comparison of both the nucleotide and amino acid sequences of the human B7-2 with the GenBank and EMBL databases yielded significant homology (about 26% amino acid sequence identity) with human B7-1. Since human B7-1, human B7-2 and murine B7-1 all bind to human CTLA4 and CD28, the homologous amino acids probably represent those necessary to comprise a CTLA4 or CD28 binding sequence. *E. coli* transfected with a vector containing a cDNA insert encoding human B7-2 (clone 29) was deposited with the American Type Culture Collection (ATCC) on July 26, 1993 as Accession No. 69357.

V. Cloning Novel B Lymphocyte Antigens from Other Mammalian Species

The present invention is not limited to human nucleic acid molecules and contemplates that novel B lymphocyte antigen homologues from other mammalian species that express B lymphocyte antigens can be cloned and sequenced using the techniques described herein. B lymphocyte antigens isolated for one species (e.g., humans) which exhibit cross-species reactivity may be used to modify T cell mediated immune responses in a different species (e.g., mice). Isolation of cDNA clones from other species can also be accomplished using human cDNA inserts, such as human B7-2 cDNA, as hybridization probes.

As described in Example 6, a murine B7-2 clone (mB7-2, clone 4) was obtained containing an insert of 1,163 base pairs with a single long open reading frame of 927 nucleotides and approximately 126 nucleotides of 3' noncoding sequences (Figure 14, SEQ ID NO:22). The predicted amino acid sequence encoded by the open reading frame of the protein is shown below the nucleotide sequence in Figure 14. The encoded murine B7-2 protein, is predicted to be 309 amino acid residues in length (SEQ ID NO:23). This protein sequence exhibits many features common to other type I Ig superfamily membrane proteins. Protein translation is predicted to begin at the methionine codon (ATG, nucleotides 111 to 113) based on the DNA homology in this region with the consensus eucaryotic translation initiation site (see Kozak, M. (1987) *Nucl. Acids Res.* 15:8125-8148). The amino terminus of the murine B7-2 protein (amino acids 1 to 23) has the characteristics of a secretory signal peptide with a predicted cleavage between the alanine at position 23 and the valine at position 24 (von Heijne (1987) *Nucl. Acids Res.* 14:4683). Processing at this site would result in a murine B7-2 membrane bound protein of 286 amino acids having an unmodified molecular weight of approximately 32 kDa. This protein would consist of an approximate extracellular Ig superfamily V and C like domains of from about amino acid residue 24 to 246, a hydrophobic transmembrane domain of from about amino acid residue 247 to 265, and a long cytoplasmic domain of from about amino acid residue 266 to 309. The homologies to the Ig superfamily are due to the two contiguous Ig-like domains in the extracellular region bound by the cysteines at positions 40 to 110 and 157 to 216. The extracellular domain also contains nine potential N-linked glycosylation sites and, like murine B7-1, is probably glycosylated. Glycosylation of the murine B7-2 protein may increase the molecular weight to about 50-70 kDa. The cytoplasmic domain of murine B7-2 contains a common region which has a cysteine followed by positively charged amino acids which presumably functions as signaling or regulatory domain within an APC. Comparison of both the nucleotide and amino acid sequences of murine B7-2 with the GenBank and EMBL databases yielded significant homology (about 26% amino acid sequence identity) with human and murine B7-

1. Murine B7-2 exhibits about 50% identity and 67% similarity with its human homologue, hB7-2. *E. coli* (DH106/p3) transfected with a vector (plasmid pmBx4) containing a cDNA insert encoding murine B7-2 (clone 4) was deposited with the American Type Culture Collection (ATCC) on August 18, 1993 as Accession No. 69388.

5 Nucleic acids which encode novel B lymphocyte antigens from other species, such as the murine B7-2, can be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at
10 prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, murine B7-2 cDNA or an appropriate sequence thereof can be used to clone genomic B7-2 in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express B7-2 protein. Methods for generating transgenic animals,
15 particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for B7-2 transgene incorporation with tissue specific enhancers, which could result in T cell costimulation and enhanced T cell proliferation and autoimmunity. Transgenic animals that include a copy of a B7-2 transgene introduced into the germ line of the animal at
20 an embryonic stage can be used to examine the effect of increased B7 expression. Such animals can be used as tester animals for reagents thought to confer protection from, for example, autoimmune disease. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the disease, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the
25 disease.

 Alternatively, the non-human homologues of B7-2 can be used to construct a B7-2 "knock out" animal which has a defective or altered B7-2 gene as a result of homologous recombination between the endogenous B7-2 gene and altered B7-2 genomic DNA introduced into an embryonic cell of the animal. For example, murine B7-2 cDNA can
30 be used to clone genomic B7-2 in accordance with established techniques. A portion of the genomic B7-2 DNA (e.g., such as an exon which encodes an extracellular domain) can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas, K.R. and Capecchi, M. R.
35 (1987) *Cell* 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected (see

e.g., Li, E. et al. (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras (see e.g., Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harbouring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for their ability to accept grafts, reject tumors and defend against infectious diseases and can be used in the study of basic immunobiology.

VI. Expression of B Lymphocyte Antigens

Host cells transfected to express peptides having the activity of a novel B lymphocyte antigen are also within the scope of the invention. The host cell may be any procaryotic or eucaryotic cell. For example, a peptide having B7-2 activity may be expressed in bacterial cells such as *E. coli*, insect cells (baculovirus), yeast, or mammalian cells such as Chinese hamster ovary cells (CHO) and NS0 cells. Other suitable host cells may be found in Goeddel, (1990) *supra* or are known to those skilled in the art.

For example, expression in eucaryotic cells such as mammalian, yeast, or insect cells can lead to partial or complete glycosylation and/or formation of relevant inter- or intra-chain disulfide bonds of recombinant protein. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). Baculovirus vectors available for expression of proteins in cultured insect cells (SF 9 cells) include the pAc series (Smith et al., (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow, V.A., and Summers, M.D., (1989) *Virology* 170:31-39). Generally, COS cells (Gluzman, Y., (1981) *Cell* 23:175-182) are used in conjunction with such vectors as pCDM8 (Seed, B., (1987) *Nature* 329:840) for transient amplification/expression in mammalian cells, while CHO (dhfr⁻ Chinese Hamster Ovary) cells are used with vectors such as pMT2PC (Kaufman et al. (1987), *EMBO J.* 6:187-195) for stable amplification/expression in mammalian cells. A preferred cell line for production of recombinant protein is the NS0 myeloma cell line available from the ECACC (catalog #85110503) and described in Galfre, G. and Milstein, C. ((1981) *Methods in Enzymology* 73(13):3-46; and *Preparation of Monoclonal Antibodies: Strategies and Procedures*, Academic Press, N.Y., N.Y). Vector DNA can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofectin, or electroporation.

Suitable methods for transforming host cells can be found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory press (1989)), and other laboratory textbooks. When used in mammalian cells, the expression vector's control functions are often provided by viral material. For example, commonly used
5 promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and most frequently, Simian Virus 40.

It is known that a small fraction of cells (about 1 out of 10^5) typically integrate DNA into their genomes. In order to identify these integrants, a gene that contains a selectable marker (i.e., resistance to antibiotics) is generally introduced into the host cells
10 along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Selectable markers may be introduced on the same plasmid as the gene of interest or may be introduced on a separate plasmid. Cells containing the gene of interest can be identified by drug selection; cells that have incorporated the selectable marker gene will survive, while the other cells die. The
15 surviving cells can then be screened for production of novel B lymphocyte antigens by cell surface staining with ligands to the B cell antigens (e.g., CTLA4Ig and CD28Ig). Alternatively, the protein can be metabolically radiolabeled with a labeled amino acid and immunoprecipitated from cell supernatant with an anti-B lymphocyte antigen monoclonal antibody or a fusion protein such as CTLA4Ig or CD28Ig.

Expression in procaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids usually to the amino terminus of the expressed target gene. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the target
25 recombinant protein; and 3) to aid in the purification of the target recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the target recombinant protein to enable separation of the target recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition
30 sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase, maltose E binding protein, or protein A, respectively, to the target recombinant protein.

E. coli expression systems include the inducible expression vectors pTrc
35 (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11 (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89; commercially available from Novagen). In the pTrc vector system, the inserted gene is

expressed with a pelB signal sequence by host RNA polymerase transcription from a hybrid trp-lac fusion promoter. After induction, the recombinant protein can be purified from the periplasmic fraction. In the pET 11 vector system, the target gene is expressed as non-fusion protein by transcription from the T7 gn10-lac 0 fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host *E. coli* strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 under the transcriptional control of the lacUV 5 promoter. In this system, the recombinant protein can be purified from inclusion bodies in a denatured form and, if desired, renatured by step gradient dialysis to remove denaturants.

One strategy to maximize recombinant B7-2 expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy would be to alter the nucleic acid sequence of the B7-2 gene or other DNA to be inserted into an expression vector so that the individual codons for each amino acid would be those preferentially utilized in highly expressed *E. coli* proteins (Wada *et al.*, (1992) *Nuc. Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention could be carried out by standard DNA synthesis techniques.

Novel B lymphocyte antigens and portions thereof, expressed in mammalian cells or otherwise, can be purified according to standard procedures of the art, including ammonium sulfate precipitation, fractionation column chromatography (e.g. ion exchange, gel filtration, electrophoresis, affinity chromatography, etc.) and ultimately, crystallization (see generally, "Enzyme Purification and Related Techniques", *Methods in Enzymology*, 22:233-577 (1971)). Once purified, partially or to homogeneity, the recombinantly produced B lymphocyte antigens or portions thereof can be utilized in compositions suitable for pharmaceutical administration as described in detail herein.

VII. Modifications of Nucleic Acid and Amino Acid Sequences of the Invention and Assays for B7 Lymphocyte Antigen Activity

It will be appreciated by those skilled in the art that other nucleic acids encoding peptides having the activity of a novel B lymphocyte antigen can be isolated by the above process. Different cell lines can be expected to yield DNA molecules having different sequences of bases. Additionally, variations may exist due to genetic polymorphisms or cell-mediated modifications of the genetic material. Furthermore, the DNA sequence of a B lymphocyte antigen can be modified by genetic techniques to produce proteins or peptides with altered amino acid sequences. Such sequences are considered within the scope of the

present invention, where the expressed peptide is capable of either inducing or inhibiting activated T cell mediated immune responses and immune function.

A number of processes can be used to generate equivalents or fragments of an isolated DNA sequence. Small subregions or fragments of the nucleic acid encoding the B7-2 protein, for example 1-30 bases in length, can be prepared by standard, synthetic organic chemical means. The technique is also useful for preparation of antisense oligonucleotides and primers for use in the generation of larger synthetic fragments of B7-2 DNA.

Larger subregions or fragments of the genes encoding B lymphocyte antigens can be expressed as peptides by synthesizing the relevant piece of DNA using the polymerase chain reaction (PCR) (Sambrook, Fritsch and Maniatis, *2 Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y., (1989)), and ligating the thus obtained DNA into an appropriate expression vector. Using PCR, specific sequences of the cloned double stranded DNA are generated, cloned into an expression vector, and then assayed for CTLA4/CD28 binding activity. For example, to express a secreted (soluble) form of the human B7-2 protein, using PCR, a DNA can be synthesized which does not encode the transmembrane and cytoplasmic regions of the protein. This DNA molecule can be ligated into an appropriate expression vector and introduced into a host cell such as CHO, where the B7-2 protein fragment is synthesized and secreted. The B7-2 protein fragment can then readily be obtained from the culture media.

In another embodiment, mutations can be introduced into a DNA by any one of a number of methods, including those for producing simple deletions or insertions, systematic deletions, insertions or substitutions of clusters of bases or substitutions of single bases, to generate variants or modified equivalents of B lymphocyte antigen DNA. For example, changes in the human B7-2 cDNA sequence shown in Figure 8 (SEQ ID NO:1) or murine B7-2 cDNA sequence shown in Figure 14 (SEQ ID NO:22) such as amino acid substitutions or deletions are preferably obtained by site-directed mutagenesis. Site directed mutagenesis systems are well known in the art. Protocols and reagents can be obtained commercially from Amersham International PLC, Amersham, U.K.

Peptides having an activity of a novel B lymphocyte antigen, i.e., the ability to bind to the natural ligand(s) of a B lymphocyte antigen on T cells and either stimulate (amplify) or inhibit (block) activated T cell mediated immune responses, as evidenced by, for example, cytokine production and/or T cell proliferation by T cells that have received a primary activation signal are considered within the scope of the invention. More specifically, peptides that bind to T lymphocytes, for example CD28⁺ cells, may be capable of delivering a costimulatory signal to the T lymphocytes, which, when transmitted in the presence of antigen and class II MHC, or other material capable of transmitting a primary signal to the T cell, results in activation of cytokine genes within the T cell. Alternatively, such a peptide

can be used in conjunction with class I MHC to thereby activate CD8⁺ cytolytic T cells. In addition, soluble, monomeric forms of the B7-2 protein, may retain the ability to bind to their natural ligand(s) on CD28⁺ T cells but, perhaps because of insufficient cross-linking with the ligand, fail to deliver the secondary signal essential for enhanced cytokine production and cell division. Such peptides, which provide a means to induce a state of anergy or tolerance in the cells, are also considered within the scope of the invention.

Screening the peptides for those which retain a characteristic B lymphocyte antigen activity as described herein can be accomplished using one or more of several different assays. For example, the peptides can be screened for specific reactivity with an anti-B7-2 monoclonal antibody reactive with cell surface B7-2 or with a fusion protein, such as CTLA4Ig or CD28Ig. Specifically, appropriate cells, such as COS cells, can be transfected with a B7-2 DNA encoding a peptide and then analyzed for cell surface phenotype by indirect immunofluorescence and flow cytometry to determine whether the peptide has B7-2 activity. Cell surface expression of the transfected cells is evaluated using a monoclonal antibody specifically reactive with cell surface B7-2 or with a CTLA4Ig or CD28Ig fusion protein. Production of secreted forms of B7-2 is evaluated using anti-B7-2 monoclonal antibody or CTLA4Ig or CD28 fusion protein for immunoprecipitation.

Other, more preferred, assays take advantage of the functional characteristics of the B7-2 antigen. As previously set forth, the ability of T cells to synthesize cytokines depends not only on occupancy or cross-linking of the T cell receptor for antigen (the "primary activation signal" provided by, for example anti-CD3, or phorbol ester to produce an "activated T cell"), but also on the induction of a costimulatory signal, in this case, by interaction with a B lymphocyte antigen, such as B7-2, B7-1 or B7-3. The binding of B7-2 to its natural ligand(s) on, for example, CD28⁺ T cells, has the effect of transmitting a signal to the T cell that induces the production of increased levels of cytokines, particularly of interleukin-2, which in turn stimulates the proliferation of the T lymphocytes. Other assays for B7-2 function thus involve assaying for the synthesis of cytokines, such as interleukin-2, interleukin-4 or other known or unknown novel cytokines, and/or assaying for T cell proliferation by CD28⁺ T cells which have received a primary activation signal.

In vitro, T cells can be provided with a first or primary activation signal by anti-T3 monoclonal antibody (e.g. anti-CD3) or phorbol ester or, more preferably, by antigen in association with class II MHC. T cells which have received a primary activation signal are referred to herein as activated T cells. B7-2 function is assayed by adding a source of B7-2 (e.g., cells expressing a peptide having B7-2 activity or a secreted form of B7-2) and a primary activation signal such as antigen in association with Class II MHC to a T cell culture and assaying the culture supernatant for interleukin-2, gamma interferon, or other known or unknown cytokine. For example, any one of several conventional assays for interleukin-2

can be employed, such as the assay described in *Proc. Natl. Acad. Sci. USA*, 86:1333 (1989) the pertinent portions of which are incorporated herein by reference. A kit for an assay for the production of interferon is also available from Genzyme Corporation (Cambridge, MA.). T cell proliferation can also be measured as described in the Examples below. Peptides that
5 retain the characteristics of the B7-2 antigen as described herein may result in increased per cell production of cytokines, such as IL-2, by T cells and may also result in enhanced T cell proliferation when compared to a negative control in which a costimulatory signal is lacking.

The same basic functional assays can also be used to screen for peptides having B7-2 activity, but which lack the ability to deliver a costimulatory signal, but in the
10 case of such peptides, addition of the B7-2 protein will not result in a marked increase in proliferation or cytokine secretion by the T cells. The ability of such proteins to inhibit or completely block the normal B7-2 costimulatory signal and induce a state of anergy can be determined using subsequent attempts at stimulation of the T cells with antigen presenting cells that express cell surface B7-2 and present antigen. If the T cells are unresponsive to the
15 subsequent activation attempts, as determined by IL-2 synthesis and T cell proliferation, a state of anergy has been induced. See, e.g., Gimmi, C.D. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90, 6586-6590; and Schwartz (1990) *Science*, 248, 1349-1356, for assay systems that can be used as the basis for an assay in accordance with the present invention.

It is possible to modify the structure of a peptide having the activity of a novel
20 B lymphocyte antigen for such purposes as increasing solubility, enhancing therapeutic or prophylactic efficacy, or stability (e.g., shelf life *ex vivo* and resistance to proteolytic degradation *in vivo*). Such modified peptides are considered functional equivalents of the B lymphocyte antigens as defined herein. For example, a peptide having B7-2 activity can be modified so that it maintains the ability to co-stimulate T cell proliferation and/or produce
25 cytokines. Those residues shown to be essential to interact with the CTLA4/CD28 receptors on T cells can be modified by replacing the essential amino acid with another, preferably similar amino acid residue (a conservative substitution) whose presence is shown to enhance, diminish, but not eliminate, or not effect receptor interaction. In addition, those amino acid residues which are not essential for receptor interaction can be modified by being replaced by
30 another amino acid whose incorporation may enhance, diminish, or not effect reactivity.

Another example of modification of a peptide having the activity of a novel B lymphocyte antigen is substitution of cysteine residues preferably with alanine, serine, threonine, leucine or glutamic acid residues to minimize dimerization via disulfide linkages. In addition, amino acid side chains of a peptide having B7-2 activity can be chemically
35 modified. Another modification is cyclization of the peptide.

In order to enhance stability and/or reactivity, peptides having B7-2 activity can be modified to incorporate one or more polymorphisms in the amino acid sequence of the

antigen resulting from any natural allelic variation. Additionally, D-amino acids, non-natural amino acids, or non-amino acid analogs can be substituted or added to produce a modified protein within the scope of this invention. Furthermore, the peptides can be modified using polyethylene glycol (PEG) according to the method of A. Schon and co-workers (Wie et al.,
5 *supra*) to produce a peptide conjugated with PEG. In addition, PEG can be added during chemical synthesis of the peptide. Other modifications of the peptides include reduction/alkylation (Tarr in: *Methods of Protein Microcharacterization*, J. E. Silver ed., Humana Press, Clifton NJ 155-194 (1986)); acylation (Tarr, *supra*); chemical coupling to an appropriate carrier (Mishell and Shiigi, eds, *Selected Methods in Cellular Immunology*, WH
10 Freeman, San Francisco, CA (1980), U.S. Patent 4,939,239; or mild formalin treatment (Marsh (1971), *Int. Arch. of Allergy and Appl. Immunol.* 41:199-215).

To facilitate purification and potentially increase solubility of a peptide, it is possible to add an amino acid fusion moiety to the protein backbone. For example, hexahistidine can be added to the peptide for purification by immobilized metal ion affinity
15 chromatography (Hochuli, E. et al., (1988) *Bio/Technology* 6:1321-1325). In addition, to facilitate isolation of a B lymphocyte antigen free of irrelevant sequences, specific endoprotease cleavage sites can be introduced between the sequences of a fusion moiety and the peptide. It may be necessary to increase the solubility of a peptide by adding functional groups to the peptide, or by omitting hydrophobic regions of the peptide.

20

VII. Uses of Nucleic Acid Sequences Encoding B Lymphocyte Antigens and Peptides Having B7-2 Activity

A. Molecular Probes

25 The nucleic acids of this invention are useful diagnostically, for tracking the progress of disease, by measuring the activation status of B lymphocytes in biological samples or for assaying the effect of a molecule on the expression of a B lymphocyte antigen (e.g., detecting cellular mRNA levels). In accordance with these diagnostic assays, the nucleic acid sequences are labeled with a detectable marker, e.g., a radioactive, fluorescent,
30 or biotinylated marker and used in a conventional dot blot or Northern hybridization procedure to probe mRNA molecules of total or poly(A⁺) RNAs from a biological sample.

B. Antibody Production

35 The peptides and fusion proteins produced from the nucleic acid molecules of the present invention can also be used to produce antibodies specifically reactive with B lymphocyte antigens. For example, by using a full-length B7-2 protein, or a peptide fragment thereof, having an amino acid sequence based on the predicted amino acid sequence of B7-2,

anti-protein/anti-peptide polyclonal antisera or monoclonal antibodies can be made using standard methods. A mammal, (e.g., a mouse, hamster, or rabbit) can be immunized with an immunogenic form of the protein or peptide which elicits an antibody response in the mammal. The immunogen can be, for example, a recombinant B7-2 protein, or fragment thereof, a synthetic peptide fragment or a cell that expresses a B lymphocyte antigen on its surface. The cell can be for example, a splenic B cell or a cell transfected with a nucleic acid encoding a B lymphocyte antigen of the invention (e.g., a B7-2 cDNA) such that the B lymphocyte antigen is expressed on the cell surface. The immunogen can be modified to increase its immunogenicity. For example, techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay can be used with the immunogen as antigen to assess the levels of antibodies.

Following immunization, antisera can be obtained and, if desired, polyclonal antibodies isolated from the sera. To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused with myeloma cells by standard somatic cell fusion procedures thus immortalizing these cells and yielding hybridoma cells. Such techniques are well known in the art. For example, the hybridoma technique originally developed by Kohler and Milstein (*Nature* (1975) 256:495-497) as well as other techniques such as the human B-cell hybridoma technique (Kozbar et al., *Immunol. Today* (1983) 4:72), the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al. *Monoclonal Antibodies in Cancer Therapy* (1985) (Allen R. Bliss, Inc., pages 77-96), and screening of combinatorial antibody libraries (Huse et al., *Science* (1989) 246:1275). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the peptide and monoclonal antibodies isolated.

The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with a peptide having the activity of a novel B lymphocyte antigen or fusion protein as described herein. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')₂ fragments can be generated by treating antibody with pepsin. The resulting F(ab')₂ fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The antibody of the present invention is further intended to include bispecific and chimeric molecules having an anti-B lymphocyte antigen (i.e., B7-2, B7-3) portion.

Particularly preferred antibodies are anti-human B7-2 monoclonal antibodies produced by hybridomas HA3.1F9, HA5.2B7 and HF2.3D1. The preparation and

characterization of these antibodies is described in detail in Example 8. Monoclonal antibody HA3.1F9 was determined to be of the IgG1 isotype; monoclonal antibody HA5.2B7 was determined to be of the IgG2b isotype; and monoclonal antibody HF2.3D1 was determined to be of the IgG2a isotype. Hybridoma cells were deposited with the American Type Culture
5 Collection, which meets the requirements of the Budapest Treaty, on July 19, 1994 as ATCC Accession No. HB11688 (hybridoma HA3.1F9), ATCC Accession No. HB11687 (HA5.2B7) and ATCC Accession No. HB11686 (HF2.3D1).

When antibodies produced in non-human subjects are used therapeutically in humans, they are recognized to varying degrees as foreign and an immune response may be
10 generated in the patient. One approach for minimizing or eliminating this problem, which is preferable to general immunosuppression, is to produce chimeric antibody derivatives, i.e., antibody molecules that combine a non-human animal variable region and a human constant region. Chimeric antibody molecules can include, for example, the antigen binding domain from an antibody of a mouse, rat, or other species, with human constant regions. A variety of
15 approaches for making chimeric antibodies have been described and can be used to make chimeric antibodies containing the immunoglobulin variable region which recognizes the gene product of the novel B lymphocyte antigens of the invention. See, for example, Morrison et al., *Proc. Natl. Acad. Sci. U.S.A.* 81:6851 (1985); Takeda et al., *Nature* 314:452 (1985), Cabilly et al., U.S. Patent No. 4,816,567; Boss et al., U.S. Patent No. 4,816,397;
20 Tanaguchi et al., European Patent Publication EP171496; European Patent Publication 0173494, United Kingdom Patent GB 2177096B. It is expected that such chimeric antibodies would be less immunogenic in a human subject than the corresponding non-chimeric antibody.

For human therapeutic purposes, the monoclonal or chimeric antibodies
25 specifically reactive with a peptide having the activity of a B lymphocyte antigen as described herein can be further humanized by producing human variable region chimeras, in which parts of the variable regions, especially the conserved framework regions of the antigen-binding domain, are of human origin and only the hypervariable regions are of non-human origin. General reviews of "humanized" chimeric antibodies are provided by
30 Morrison, S. L. (1985) *Science* 229:1202-1207 and by Oi et al. (1986) *BioTechniques* 4:214. Such altered immunoglobulin molecules may be made by any of several techniques known in the art, (e.g., Teng et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80:7308-7312 (1983); Kozbor et al., *Immunology Today*, 4:7279 (1983); Olsson et al., *Meth. Enzymol.*, 92:3-16 (1982)), and are preferably made according to the teachings of PCT Publication WO92/06193 or EP 0239400.
35 Humanized antibodies can be commercially produced by, for example, Scotgen Limited, 2 Holly Road, Twickenham, Middlesex, Great Britain. Suitable "humanized" antibodies can be alternatively produced by CDR or CEA substitution (see U.S. Patent 5,225,539 to Winter;

Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060). Humanized antibodies which have reduced immunogenicity are preferred for immunotherapy in human subjects.

Immunotherapy with a humanized antibody will likely reduce the necessity for any
5 concomitant immunosuppression and may result in increased long term effectiveness for the treatment of chronic disease situations or situations requiring repeated antibody treatments.

As an alternative to humanizing a monoclonal antibody from a mouse or other species, a human monoclonal antibody directed against a human protein can be generated. Transgenic mice carrying human antibody repertoires have been created which can be
10 immunized with a human B lymphocyte antigen, such as B7-2. Splenocytes from these immunized transgenic mice can then be used to create hybridomas that secrete human monoclonal antibodies specifically reactive with a human B lymphocyte antigen (see, e.g., Wood et al. PCT publication WO 91/00906, Kucherlapati et al. PCT publication WO 91/10741; Lonberg et al. PCT publication WO 92/03918; Kay et al. PCT publication
15 92/03917; Lonberg, N. et al. (1994) *Nature* 368:856-859; Green, L.L. et al. (1994) *Nature Genet.* 7:13-21; Morrison, S.L. et al. (1994) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Bruggeman et al. (1993) *Year Immunol* 7:33-40; Tuailon et al. (1993) *PNAS* 90:3720-3724; and Bruggeman et al. (1991) *Eur J Immunol* 21:1323-1326).

Monoclonal antibody compositions of the invention can also be produced by
20 other methods well known to those skilled in the art of recombinant DNA technology. An alternative method, referred to as the "combinatorial antibody display" method, has been developed to identify and isolate antibody fragments having a particular antigen specificity, and can be utilized to produce monoclonal antibodies that bind a B lymphocyte antigen of the invention (for descriptions of combinatorial antibody display see e.g., Sastry et al. (1989)
25 *PNAS* 86:5728; Huse et al. (1989) *Science* 246:1275; and Orlandi et al. (1989) *PNAS* 86:3833). After immunizing an animal with a B lymphocyte antigen, the antibody repertoire of the resulting B-cell pool is cloned. Methods are generally known for directly obtaining the DNA sequence of the variable regions of a diverse population of immunoglobulin molecules by using a mixture of oligomer primers and PCR. For instance, mixed oligonucleotide
30 primers corresponding to the 5' leader (signal peptide) sequences and/or framework 1 (FR1) sequences, as well as primer to a conserved 3' constant region primer can be used for PCR amplification of the heavy and light chain variable regions from a number of murine antibodies (Larrick et al. (1991) *Biotechniques* 11:152-156). A similar strategy can also been used to amplify human heavy and light chain variable regions from human antibodies
35 (Larrick et al. (1991) *Methods: Companion to Methods in Enzymology* 2:106-110).

In an illustrative embodiment, RNA is isolated from activated B cells of, for example, peripheral blood cells, bone marrow, or spleen preparations, using standard

protocols (e.g., U.S. Patent No. 4,683,202; Orlandi, et al. *PNAS* (1989) 86:3833-3837; Sastry et al., *PNAS* (1989) 86:5728-5732; and Huse et al. (1989) *Science* 246:1275-1281.) First-strand cDNA is synthesized using primers specific for the constant region of the heavy chain(s) and each of the κ and λ light chains, as well as primers for the signal sequence.

- 5 Using variable region PCR primers, the variable regions of both heavy and light chains are amplified, each alone or in combination, and ligated into appropriate vectors for further manipulation in generating the display packages. Oligonucleotide primers useful in amplification protocols may be unique or degenerate or incorporate inosine at degenerate positions. Restriction endonuclease recognition sequences may also be incorporated into the
- 10 primers to allow for the cloning of the amplified fragment into a vector in a predetermined reading frame for expression.

- The V-gene library cloned from the immunization-derived antibody repertoire can be expressed by a population of display packages, preferably derived from filamentous phage, to form an antibody display library. Ideally, the display package comprises a system
- 15 that allows the sampling of very large diverse antibody display libraries, rapid sorting after each affinity separation round, and easy isolation of the antibody gene from purified display packages. In addition to commercially available kits for generating phage display libraries (e.g., the Pharmacia *Recombinant Phage Antibody System*, catalog no. 27-9400-01; and the Stratagene *SurfZAP*TM phage display kit, catalog no. 240612), examples of methods and
 - 20 reagents particularly amenable for use in generating a diverse antibody display library can be found in, for example, Ladner et al. U.S. Patent No. 5,223,409; Kang et al. International Publication No. WO 92/18619; Dower et al. International Publication No. WO 91/17271; Winter et al. International Publication WO 92/20791; Markland et al. International Publication No. WO 92/15679; Breitling et al. International Publication WO 93/01288;
 - 25 McCafferty et al. International Publication No. WO 92/01047; Garrard et al. International Publication No. WO 92/09690; Ladner et al. International Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum Antibod Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J* 12:725-734; Hawkins et al. (1992) *J Mol Biol* 226:889-896; Clackson et al. (1991) *Nature* 352:624-
 - 30 628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrad et al. (1991) *Bio/Technology* 9:1373-1377; Hoogenboom et al. (1991) *Nuc Acid Res* 19:4133-4137; and Barbas et al. (1991) *PNAS* 88:7978-7982.

- In certain embodiments, the V region domains of heavy and light chains can be expressed on the same polypeptide, joined by a flexible linker to form a single-chain Fv
- 35 fragment, and the scFV gene subsequently cloned into the desired expression vector or phage genome. As generally described in McCafferty et al., *Nature* (1990) 348:552-554, complete V_H and V_L domains of an antibody, joined by a flexible (Gly₄-Ser)₃ linker can be used to

produce a single chain antibody which can render the display package separable based on antigen affinity. Isolated scFV antibodies immunoreactive with a peptide having activity of a B lymphocyte antigen can subsequently be formulated into a pharmaceutical preparation for use in the subject method.

5 Once displayed on the surface of a display package (e.g., filamentous phage), the antibody library is screened with a B lymphocyte antigen protein, or peptide fragment thereof, to identify and isolate packages that express an antibody having specificity for the B lymphocyte antigen. Nucleic acid encoding the selected antibody can be recovered from the display package (e.g., from the phage genome) and subcloned into other expression vectors
10 by standard recombinant DNA techniques.

 The antibodies of the current invention can be used therapeutically to inhibit T cell activation through blocking receptor:ligand interactions necessary for costimulation of the T cell. These so-called "blocking antibodies" can be identified by their ability to inhibit T cell proliferation and/or cytokine production when added to an *in vitro* costimulation assay as
15 described herein. The ability of blocking antibodies to inhibit T cell functions may result in immunosuppression and/or tolerance when these antibodies are administered *in vivo*.

C. Protein Purification

 The polyclonal or monoclonal antibodies of the current invention, such as an
20 antibody specifically reactive with a recombinant or synthetic peptide having B7-2 activity or B7-3 activity can also be used to isolate the native B lymphocyte antigen from cells. For example, antibodies reactive with the peptide can be used to isolate the naturally-occurring or native form of B7-2 from activated B lymphocytes by immunoaffinity chromatography. In addition, the native form of B7-3 can be isolated from B cells by immunoaffinity
25 chromatography with monoclonal antibody BB-1.

D. Other Therapeutic Reagents

 The nucleic acid sequences and novel B lymphocyte antigens described herein can be used in the development of therapeutic reagents having the ability to either upregulate
30 (e.g., amplify) or downregulate (e.g., suppress or tolerize) T cell mediated immune responses. For example, peptides having B7-2 activity, including soluble, monomeric forms of the B7-2 antigen or a B7-2 fusion protein, e.g., B7-2Ig, and anti-B7-2 antibodies that fail to deliver a costimulatory signal to T cells that have received a primary activation signal, can be used to block the B7-2 ligand(s) on T cells and thereby provide a specific means by which to cause
35 immunosuppression and/or induce tolerance in a subject. Such blocking or inhibitory forms of B lymphocyte antigens and fusion proteins and blocking antibodies can be identified by their ability to inhibit T cell proliferation and/or cytokine production when added to an *in*

vitro costimulation assay as previously described herein. In contrast to the monomeric form, stimulatory forms of B7-2, such as an intact cell surface B7-2, retain the ability to transmit the costimulatory signal to the T cells, resulting in an increased secretion of cytokines when compared to activated T cells that have not received the secondary signal.

5 In addition, fusion proteins comprising a first peptide having an activity of B7-2 fused to a second peptide having an activity of another B lymphocyte antigen (e.g., B7-1) can be used to modify T cell mediated immune responses. Alternatively, two separate peptides having an activity of B lymphocyte antigens, for example, B7-2 and B7-1, or a combination of blocking antibodies (e.g., anti-B7-2 and anti-B7-1 monoclonal antibodies)
10 can be combined as a single composition or administered separately (simultaneously or sequentially), to upregulate or downregulate T cell mediated immune responses in a subject. Furthermore, a therapeutically active amount of one or more peptides having B7-2 activity and or B7-1 activity can be used in conjunction with other immunomodulating reagents to influence immune responses. Examples of other immunomodulating reagents include
15 blocking antibodies, e.g., against CD28 or CTLA4, against other T cell markers or against cytokines, fusion proteins, e.g., CTLA4Ig, or immunosuppressive drugs, e.g., cyclosporine A or FK506.

The peptides produced from the nucleic acid molecules of the present invention may also be useful in the construction of therapeutic agents which block T cell
20 function by destruction of the T cell. For example, as described, secreted forms of a B lymphocyte antigen can be constructed by standard genetic engineering techniques. By linking a soluble form of B7-1, B7-2 or B7-3 to a toxin such as ricin, an agent capable of preventing T cell activation can be made. Infusion of one or a combination of immunotoxins, e.g., B7-2-ricin, B7-1-ricin, into a patient may result in the death of T cells, particularly of
25 activated T cells that express higher amounts of CD28 and CTLA4. Soluble forms of B7-2 in a monovalent form alone may be useful in blocking B7-2 function, as described above, in which case a carrier molecule may also be employed.

Another method of preventing the function of a B lymphocyte antigen is through the use of an antisense or triplex oligonucleotide. For example, an oligonucleotide
30 complementary to the area around the B7-1, B7-2 or B7-3 translation initiation site, (e.g., for B7-1, TGGCCCATGGCTTCAGA, (SEQ ID NO:20) nucleotides 326-309 and for B7-2, GCCAAAATGGATCCCCA (SEQ ID NO:21)), can be synthesized. One or more antisense oligonucleotides can be added to cell media, typically at 200 µg/ml, or administered to a patient to prevent the synthesis of B7-1, B7-2 and/or B7-3. The antisense oligonucleotide is
35 taken up by cells and hybridizes to the appropriate B lymphocyte antigen mRNA to prevent translation. Alternatively, an oligonucleotide which binds double-stranded DNA to form a

triplex construct to prevent DNA unwinding and transcription can be used. As a result of either, synthesis of one or more B lymphocyte antigens is blocked.

In a specific embodiment, T cells are obtained from a subject and cultured *ex vivo* to expand the population of T cells. In a further embodiment the T cells are then administered to a subject. T cells can be stimulated to proliferate *in vitro* by, for example, providing to the T cells a primary activation signal and a costimulatory signal, as described in detail in the Examples section. A preferred costimulatory molecule for stimulating proliferation of activated T cells, such as T cells stimulated through their T cell receptor, is aB7-2VIg fusion protein. However, other forms of B7-2Ig fusion proteins can also be used to costimulate proliferation of T cells. In a specific embodiment of the invention, activated T cells are costimulated with a B7-2VIg protein, such that a response by the activated T cells is stimulated. In one embodiment T cells are cultured *ex vivo* according to the method described in PCT Application No. WO 94/29436, using B7-2Ig, or more preferably B7-2VIg as the costimulatory molecule. The costimulatory molecule can be soluble, attached to a cell membrane or attached to a solid surface, such as a bead. In a preferred embodiment, a T helper-type 2 (Th2) response is preferentially stimulated.

E. Therapeutic Uses by Downregulation of Immune Responses

Given the structure and function of the novel B lymphocyte antigens disclosed herein, it is possible to downregulate the function of a B lymphocyte antigen, and thereby downregulate immune responses, in a number of ways. Downregulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Downregulating or preventing one or more B lymphocyte antigen functions, e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7

lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural
5 ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this manner prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may
10 avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens. For example, it may be desirable to block the function of B7-2 and B7-1, B7-2 and B7-3, B7-1 and B7-3 or B7-2, B7-1 and B7-3 by administering a soluble form of a combination of peptides having an activity of each of
15 these antigens or a blocking antibody (separately or together in a single composition) prior to transplantation. Alternatively, inhibitory forms of B lymphocyte antigens can be used with other suppressive agents such as blocking antibodies against other T cell markers or against cytokines, other fusion proteins, e.g., CTLA4Ig, or immunosuppressive drugs.

The efficacy of particular blocking reagents in preventing organ transplant
20 rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. The functionally important aspects of B7-1 are conserved structurally between species and it is therefore likely that other B lymphocyte antigens can function across species, thereby allowing use of reagents composed of human proteins in animal systems. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and
25 xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow et al., *Science*, 257: 789-792 (1992) and Turka et al., *Proc. Natl. Acad. Sci. USA*, 89: 11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect
30 of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking B lymphocyte antigen function, e.g., by use of a peptide having B7-2 activity alone or in combination with a peptide having B7-1 activity and/or a peptide having B7-3 activity, may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive
35 against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block

costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 840-856).

The IgE antibody response in atopic allergy is highly T cell dependent and, thus, inhibition of B lymphocyte antigen induced T cell activation may be useful therapeutically in the treatment of allergy and allergic reactions. An inhibitory form of B7-2 protein, such as a peptide having B7-2 activity alone or in combination with a peptide having the activity of another B lymphocyte antigen, such as B7-1, can be administered to an allergic subject to inhibit T cell mediated allergic responses in the subject. Inhibition of B lymphocyte antigen costimulation of T cells may be accompanied by exposure to allergen in conjunction with appropriate MHC molecules. Allergic reactions may be systemic or local in nature, depending on the route of entry of the allergen and the pattern of deposition of IgE on mast cells or basophils. Thus, it may be necessary to inhibit T cell mediated allergic responses locally or systemically by proper administration of an inhibitory form of B7-2 protein.

Inhibition of T cell activation through blockage of B lymphocyte antigen function may also be important therapeutically in viral infections of T cells. For example, in the acquired immune deficiency syndrome (AIDS), viral replication is stimulated by T cell activation. Blocking B7-2 function could lead to a lower level of viral replication and thereby ameliorate the course of AIDS. In addition, it may also be necessary to block the function of a combination of B lymphocyte antigens i.e., B7-1, B7-2 and B7-3. Surprisingly, HTLV-I infected T cells express B7-1 and B7-2. This expression may be important in the growth of HTLV-I infected T cells and the blockage of B7-1 function together with the function of B7-2 and/or B7-3 may slow the growth of HTLV-I induced leukemias. Alternatively, stimulation of viral replication by T cell activation may be induced by contact with a stimulatory form of B7-2 protein, for such purposes as generating retroviruses (e.g., various HIV isolates) in sufficient quantities for isolation and use.

F. Therapeutic Uses by Upregulation of Immune Responses

Upregulation of a B lymphocyte antigen function, as a means of upregulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. Viral infections are cleared primarily by cytolytic T cells. In accordance with the present invention, it is believed that B7-2 and thus, B7-1 and B7-3 with their natural ligand(s) on T cells may result in an increase in the cytolytic activity of at least some T cells. It is also believed that B7-2, B7-1, and B7-3 are involved in the initial activation and generation of CD8+ cytotoxic T cells. The addition of a soluble peptide having B7-2 activity, alone, or in combination with a peptide having the activity of another B lymphocyte antigen, in a multi-valent form, to stimulate T cell activity through the costimulation pathway would thus be therapeutically useful in situations where more rapid or thorough clearance of virus would be beneficial. These would include viral skin diseases such as Herpes or shingles, in which cases the multi-valent soluble peptide having B7-2 activity or combination of such peptide and/or a peptide having B7-1 activity and/or a peptide having B7-3 activity is delivered topically to the skin. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide having B7-2 activity (alone or in combination with a peptide having B7-1 activity and/or a peptide having B7-3 activity) or together with a stimulatory form of a soluble peptide having B7-2 activity (alone or in combination with a peptide having B7-1 activity and/or a peptide having B7-3 activity) and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a peptide having the activity of a B lymphocyte antigen as described herein such that the cells express all or a portion of a B lymphocyte antigen on their surface, e.g., B7-2 or B7-3, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

Stimulatory forms of B lymphocyte antigens may also be used prophylactically in vaccines against various pathogens. Immunity against a pathogen, e.g., a virus, could be induced by vaccinating with a viral protein along with a stimulatory form of a peptide having B7-2 activity or another peptide having the activity of B lymphocyte antigen in an appropriate adjuvant. Alternately, an expression vector which encodes genes for both a

pathogenic antigen and a peptide having the activity of a B lymphocyte antigen, e.g., a vaccinia virus expression vector engineered to express a nucleic acid encoding a viral protein and a nucleic acid encoding a peptide having B7-2 activity as described herein, can be used for vaccination. Presentation of B7-2 with class I MHC proteins by, for example, a cell
5 transfected to coexpress a peptide having B7-2 activity and MHC class I α chain protein and β_2 microglobulin may also result in activation of cytolytic CD8⁺ T cells and provide immunity from viral infection. Pathogens for which vaccines may be useful include hepatitis B, hepatitis C, Epstein-Barr virus, cytomegalovirus, HIV-1, HIV-2, tuberculosis, malaria and schistosomiasis.

10 In another application, upregulation or enhancement of B lymphocyte antigen function may be useful in the induction of tumor immunity. Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide having the activity of a B lymphocyte antigen, such as B7-2, can be administered to a subject to overcome tumor-specific tolerance in the subject. If
15 desired, the tumor cell can be transfected to express a combination of peptides having the activity of a number of B lymphocyte antigens (e.g., B7-1, B7-2, B7-3). For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2 activity alone, or in conjunction with a peptide having B7-1 activity and/or B7-3 activity. The transfected tumor cells are returned to the patient to
20 result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition,
25 tumor cells which lack MHC class I or MHC class II molecules, or which fail to express sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I α chain protein and β_2 microglobulin protein or an MHC class II α chain protein and an MHC class II β chain protein to thereby express MHC class I or MHC class II
30 proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell.

Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA
35 encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Expression of B7-1 by B7 negative murine tumor cells has been shown to induce T cell mediated specific immunity

accompanied by tumor rejection and prolonged protection to tumor challenge in mice (Chen, L., et al. (1992) *Cell* 71, 1093-1102; Townsend, S.E. and Allison, J.P. (1993) *Science* 259, 368-370; Baskar, S., et al. (1993) *Proc. Natl. Acad. Sci.* 90, 5687-5690). Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome
5 tumor-specific tolerance in the subject.

In another aspect, a stimulatory form of one or more soluble peptides having an activity of a B lymphocyte antigen can be administered to a tumor-bearing patient to provide a costimulatory signal to T cells in order to induce anti-tumor immunity.

10 G. Administration of Therapeutic Forms of B Lymphocyte Antigens

The peptides of the invention are administered to subjects in a biologically compatible form suitable for pharmaceutical administration *in vivo* to either enhance or suppress T cell mediated immune response. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the protein to be administered in which any toxic
15 effects are outweighed by the therapeutic effects of the protein. The term subject is intended to include living organisms in which an immune response can be elicited, e.g., mammals. Examples of subjects include humans, dogs, cats, mice, rats, and transgenic species thereof. Administration of a peptide having the activity of a novel B lymphocyte antigen as described herein can be in any pharmacological form including a therapeutically active amount of
20 peptide alone or in combination with a peptide having the activity of another B lymphocyte antigen and a pharmaceutically acceptable carrier. Administration of a therapeutically active amount of the therapeutic compositions of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of a peptide having B7-2 activity may vary
25 according to factors such as the disease state, age, sex, and weight of the individual, and the ability of peptide to elicit a desired response in the individual. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

30 The active compound (e.g., peptide) may be administered in a convenient manner such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the active compound may be coated in a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound.

35 To administer a peptide having B7-2 activity by other than parenteral administration, it may be necessary to coat the peptide with, or co-administer the peptide with, a material to prevent its inactivation. For example, a peptide having B7-2 activity may

be administered to an individual in an appropriate carrier, diluent or adjuvant, co-administered with enzyme inhibitors or in an appropriate carrier such as liposomes. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Adjuvant is used in its broadest sense and includes any immune stimulating compound such as
5 interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether. Enzyme inhibitors include pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEP) and trasylol. Liposomes include water-in-oil-in-water emulsions as well as conventional liposomes (Strejan et al., (1984) *J. Neuroimmunol* 7:27).

10 The active compound may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

Pharmaceutical compositions suitable for injectable use include sterile
15 aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases, the composition must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a
20 solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by
25 various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, asorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum
30 monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating active compound (e.g., peptide having B7-2 activity) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a
35 sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-

drying which yields a powder of the active ingredient (e.g., peptide) plus any additional desired ingredient from a previously sterile-filtered solution thereof.

When the active compound is suitably protected, as described above, the protein may be orally administered, for example, with an inert diluent or an assimilable edible carrier. As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

H. Identification of Cytokines Induced by Costimulation

The nucleic acid sequences encoding peptides having the activity of novel B lymphocyte antigens as described herein can be used to identify cytokines which are produced by T cells in response to stimulation by a form of B lymphocyte antigen, e.g., B7-2. T cells can be suboptimally stimulated *in vitro* with a primary activation signal, such as phorbol ester, anti-CD3 antibody or preferably antigen in association with an MHC class II molecule, and given a costimulatory signal by a stimulatory form of B7-2 antigen, for instance by a cell transfected with nucleic acid encoding a peptide having B7-2 activity and expressing the peptide on its surface or by a soluble, stimulatory form of the peptide. Known cytokines released into the media can be identified by ELISA or by the ability of an antibody which blocks the cytokine to inhibit T cell proliferation or proliferation of other cell types that is induced by the cytokine. An IL-4 ELISA kit is available from Genzyme (Cambridge MA), as is an IL-7 blocking antibody. Blocking antibodies against IL-9 and IL-12 are available from Genetics Institute (Cambridge, MA).

An *in vitro* T cell costimulation assay as described above can also be used in a method for identifying novel cytokines which may be induced by costimulation. If a

particular activity induced upon costimulation, e.g., T cell proliferation, cannot be inhibited by addition of blocking antibodies to known cytokines, the activity may result from the action of an unknown cytokine. Following costimulation, this cytokine could be purified from the media by conventional methods and its activity measured by its ability to induce T cell proliferation.

To identify cytokines which prevent the induction of tolerance, an *in vitro* T cell costimulation assay as described above can be used. In this case, T cells would be given the primary activation signal and contacted with a selected cytokine, but would not be given the costimulatory signal. After washing and resting the T cells, the cells would be rechallenged with both a primary activation signal and a costimulatory signal. If the T cells do not respond (e.g., proliferate or produce IL-2) they have become tolerized and the cytokine has not prevented the induction of tolerance. However, if the T cells respond, induction of tolerance has been prevented by the cytokine. Those cytokines which are capable of preventing the induction of tolerance can be targeted for blockage *in vivo* in conjunction with reagents which block B lymphocyte antigens as a more efficient means to induce tolerance in transplant recipients or subjects with autoimmune diseases. For example, one could administer a B7-2 blocking reagent together with a cytokine blocking antibody to a subject.

I. Identification of Molecules which Inhibit Costimulation

Another application of the peptide having the activity of a novel B lymphocyte antigen of the invention (e.g., B7-2 and B7-3) is the use of one or more of these peptides in screening assays to discover as yet undefined molecules which are inhibitors of costimulatory ligand binding and/or of intracellular signaling through T cells following costimulation. For example, a solid-phase binding assay using a peptide having the activity of a B lymphocyte antigen, such as B7-2, could be used to identify molecules which inhibit binding of the antigen with the appropriate T cell ligand (e.g., CTLA4, CD28). In addition, an *in vitro* T cell costimulation assay as described above could be used to identify molecules which interfere with intracellular signaling through the T cells following costimulation as determined by the ability of these molecules to inhibit T cell proliferation and/or cytokine production (yet which do not prevent binding of B lymphocyte antigens to their receptors). For example, the compound cyclosporine A inhibits T cell activation through stimulation via the T cell receptor pathway but not via the CD28/CTLA4 pathway. Therefore, a different intracellular signaling pathway is involved in costimulation. Molecules which interfere with intracellular signaling via the CD28/CTLA4 pathway may be effective as immunosuppressive agents *in vivo* (similar to the effects of cyclosporine A).

J. Identification of Molecules which Modulate B Lymphocyte Antigen Expression

The monoclonal antibodies produced using the proteins and peptides of the current invention can be used in a screening assay for molecules which modulate the expression of B lymphocyte antigens on cells. For example, molecules which effect intracellular signaling which leads to induction of B lymphocyte antigens, e.g. B7-2 or B7-3, can be identified by assaying expression of one or more B lymphocyte antigens on the cell surface. Reduced immunofluorescent staining by an anti-B7-2 antibody in the presence of the molecule would indicate that the molecule inhibits intracellular signals. Molecules which upregulate B lymphocyte antigen expression result in an increased immunofluorescent staining. Alternatively, the effect of a molecule on expression of a B lymphocyte antigen, such as B7-2, can be determined by detecting cellular B7-2 mRNA levels using a B7-2 cDNA as a probe. For example, a cell which expresses a peptide having B7-2 activity can be contacted with a molecule to be tested, and an increase or decrease in B7-2 mRNA levels in the cell detected by standard technique, such as Northern hybridization analysis or conventional dot blot of mRNA or total poly(A⁺)RNAs using a B7-2 cDNA probe labeled with a detectable marker. Molecules which modulate B lymphocyte antigen expression may be useful therapeutically for either upregulating or downregulating immune responses alone or in conjunction with soluble blocking or stimulating reagents. For instance, a molecule which inhibits expression of B7-2 could be administered together with a B7-2 blocking reagent for immunosuppressive purposes. Molecules which can be tested in the above-described assays include cytokines such as IL-4, γ INF, IL-10, IL-12, GM-CSF and prostagladins.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references and published patent applications cited throughout this application are hereby incorporated by reference.

The following methodology was used in Examples 1, 2 and 3.

METHODS AND MATERIALS

A. Cells

Mononuclear cells were isolated by Ficoll-Hypaque density gradient centrifugation from single cell suspensions of normal human spleens and were separated into E- and E+ fractions by rosetting with sheep red blood cells (Boyd, A.W., et al. (1985) *J. Immunol.* 134, 1516). B cells were purified from the E- fraction by adherence of monocytes on plastic and depletion of residual T, natural killer cells (NK) and residual monocytes by two treatments with anti-MsIgG and anti-MsIgM coated magnetic beads (Advanced Magnetix, Cambridge, MA), using monoclonal antibodies: anti-CD4, -CD8, -CD11b, -

CD14 and -CD16. CD4⁺ T cells were isolated from the E⁺ fraction of the same spleens after adherence on plastic and depletion of NK, B cells and residual monocytes with magnetic beads and monoclonal antibodies: anti-CD20, -CD11b, -CD8 and -CD16. CD28⁺ T cells were identically isolated from the E⁺ fraction using anti-CD20, -CD11b, -CD14 and -CD16 monoclonal antibodies. The efficiency of the purification was analyzed by indirect immunofluorescence and flow cytometry using an EPICS flow cytometer (Coulter). B cell preparations were >95% CD20⁺, <2% CD3⁺, <1% CD14⁺. CD4⁺ T cell preparations were >98% CD3⁺, >98% CD4⁺, <1% CD8⁺, <1% CD20⁺, <1% CD14⁺. CD28⁺ T cell preparations were >98% CD3⁺, >98% CD28⁺, <1% CD20⁺, <1% CD14⁺.

B. Monoclonal Antibodies and Fusion Proteins

Monoclonal antibodies were used as purified Ig unless indicated otherwise: anti-B7:133, IgM is a blocking antibody and has been previously described (Freedman, A.S. et al. (1987) *Immunol.* 137, 3260-3267); anti-B7:B1.1, IgG1 (RepliGen Corp., Cambridge, MA) (Nickoloff, B., et al (1993) *Am. J. Pathol.* 142, 1029-1040) is a non-blocking monoclonal antibody; BB-1: IgM is a blocking antibody (Dr. E. Clark, University of Washington, Seattle, WA) (Yokochi, T., et al. (1982) *J. Immunol.* 128, 823-827); anti-CD20: B1, IgG2a (Stashenko, P., et al. (1980) *J. Immunol.* 125, 1678-1685); anti-B5: IgM (Freedman, A., et al. (1985) *J. Immunol.* 134, 2228-2235); anti-CD8: 7PT 3F9, IgG2a; anti-CD4: 19Thy5D7, IgG2a; anti-CD11b: Mo1, IgM and anti-CD14: Mo2, IgM (Todd, R, et al. (1981) *J. Immunol.* 126, 1435-1442); anti-MHC class II: 9-49, IgG2a (Dr R. Todd, University of Michigan, Ann Arbor) (Todd, R.I., et al. (1984) *Hum Immunol.* 10, 23-40; anti-CD28: 9.3, IgG2a (Dr. C. June, Naval Research Institute, Bethesda) (Hansen, J.A., et al. (1980) *Immunogenetics.* 10, 247-260); anti-CD16: 3G8, IgG1 (used as ascites) (Dr. J. Ritz, Dana-Farber Cancer Institute, Boston); anti-CD3: OKT3, IgG2a hybridoma was obtained from the American Type Culture Collection and the purified monoclonal antibody was adhered on plastic plates at a concentration of 1 µg/ml; anti-CD28 Fab fragments were generated from the 9.3 monoclonal antibody, by papain digestion and purification on a protein A column, according to the manufacturer's instructions (Pierce, Rockford, IL). Human CTLA4 fusion protein (CTLA4Ig) and control fusion protein (control-Ig) were prepared as previously described (Gimmi, C.D., et al. (1993) *Proc. Natl. Acad. Sci USA* 90:6586-6590); Boussiotis, V., et al *J. Exp. Med.* (accepted for publication)).

C. CHO Cell Transfection

B7-1 transfectants (CHO-B7) were prepared from the B7-1 negative chinese hamster ovary (CHO) cell line, fixed with paraformaldehyde and used as previously described (Gimmi, C.D., et al. *Proc. Natl. Acad. Sci USA* 88, 6575-6579).

D. In Vitro B Cell Activation and Selection of B7+ and B7- Cells

Splenic B cells were cultured at 2×10^6 cells/ml in complete culture media, {RPMI 1640 with 10% heat inactivated fetal calf serum (FCS), 2mM glutamine, 1 mM sodium pyruvate, penicillin (100 units/ml), streptomycin sulfate (100µg/ml) and gentamycin sulfate (5µg/ml)}, in tissue culture flasks and were activated by crosslinking of sIg with affinity purified rabbit anti-human IgM coupled to Affi-Gel 702 beads (Bio-Rad), Richmond, CA) (Boyd, A.W., et al., (1985) *J. Immunol.* **134**,1516) or by crosslinking of MHC class II with 9-49 antibody coupled to Affi-Gel 702 beads. B cells activated for 72 hours, were used as total activated B cell populations or were indirectly stained with anti-B7 (B1.1) monoclonal antibody and fluorescein isothiocyanate (FITC) labeled goat anti-mouse immunoglobulin (Fisher, Pittsburgh, PA), and fractionated into B7-1+ and B7-1- populations by flow cytometric cell sorting (EPICS Elite flow cytometer, Coulter).

E. Immunofluorescence and Flow Cytometry

For surface phenotype analysis populations of B cells activated by either sIg or MHC class II crosslinking for 6, 12, 24, 48, 72 and 96 hours were stained with either anti-B7 (133), BB-1 monoclonal antibodies, control IgM antibody, CTLA4Ig or control-Ig. Cell suspensions were stained by two step indirect membrane staining with 10µg/ml of primary monoclonal antibody followed by the appropriate secondary reagents. Specifically, immunoreactivity with anti-B7 (133) and BB-1 monoclonal antibodies was studied by indirect staining using goat anti-mouse Ig or immunoglobulin FITC (Fisher) as secondary reagent and immunoreactivity with fusion proteins was studied using biotinylated CTLA4Ig or biotinylated control-Ig and streptavidin-phycoerythrin as secondary reagent. PBS containing 10% AB serum was used as diluent and wash media. Cells were fixed with 0.1% paraformaldehyde and analyzed on a flow cytometer (EPICS Elite Coulter).

F. Proliferation Assay

T cells were cultured at a concentration of 1×10^5 cells per well in 96-well flat bottom microtiter plate at 37°C for 3 days in 5% CO₂. Syngeneic activated B cells (total B cell population or B7+ and B7- fractions) were irradiated (2500 rad) and added into the cultures at a concentration of 1×10^5 cells per well. Factors under study were added to the required concentration for a total final volume of 200 µl per well. When indicated, T cells were incubated with anti-CD28 Fab (final concentration of 10µg/ml), for 30 minutes at 4°C, prior to addition in experimental plates. Similarly, CHO-B7 or B cells were incubated with CTLA4Ig or control-Ig (10µg/ml) for 30 minutes at 4°C. Thymidine incorporation as an index of mitogenic activity, was assessed after incubation with 1µCi (37kBq) of {methyl-³H}

thymidine (Du Pont, Boston, MA) for the last 15 hours of the culture. The cells were harvested onto filters and the radioactivity on the dried filters was measured in a Pharmacia beta plate liquid scintillation counter.

5 G. IL-2 and IL-4 Assay

IL-2 and IL-4 concentrations were assayed by ELISA (R&D Systems, Minneapolis, MN and BioSource, Camarillo, CA) in culture supernatants collected at 24 hours after initiation of the culture.

10 EXAMPLE 1

Expression of a Novel CTLA4 Ligand on Activated B Cells
Which Induces T Cell Proliferation

Since crosslinking surface Ig induces human resting B cells to express B7-1
15 maximally (50-80%) at 72 hours, the ability of activated human B lymphocytes to induce submitogenically activated T cells to proliferate and secrete IL-2 was determined. Figure 1 depicts the costimulatory response of human splenic CD28+ T cells, submitogenically activated with anti-CD3 monoclonal antibody, to either B7 (B7-1) transfected CHO cells (CHO-B7) or syngeneic splenic B cells activated with anti-Ig for 72 hours. ³H-Thymidine
20 incorporation was assessed for the last 15 hours of a 72 hours culture. IL-2 was assessed by ELISA in supernatants after 24 hours of culture (Detection limits of the assay: 31-2000 pg/ml). Figure 1 is representative of seventeen experiments.

Submitogenically activated CD28+ T cells proliferated and secreted high levels of IL-2 in response to B7-1 costimulation provided by CHO-B7 (Figure 1, panel a).
25 Both proliferation and IL-2 secretion were totally inhibited by blocking the B7-1 molecule on CHO cells with either anti-B7-1 monoclonal antibody or by a fusion protein for its high affinity receptor, CTLA4. Similarly, proliferation and IL-2 secretion were abrogated by blocking B7-1 signalling via CD28 with Fab anti-CD28 monoclonal antibody. Control monoclonal antibody or control fusion protein had no effect. Nearly identical costimulation
30 of proliferation and IL-2 secretion was provided by splenic B cells activated with anti-Ig for 72 hours (panel b). Though anti-B7-1 monoclonal antibody could completely abrogate both proliferation and IL-2 secretion delivered by CHO-B7, anti-B7-1 monoclonal antibody consistently inhibited proliferation induced by activated B cells by only 50% whereas IL-2 secretion was totally inhibited. In contrast to the partial blockage of proliferation induced by
35 anti-B7-1 monoclonal antibody, both CTLA4Ig and Fab anti-CD28 monoclonal antibody completely blocked proliferation and IL-2 secretion. These results are consistent with the

hypothesis that activated human B cells express one or more additional CTLA4/CD28 ligands which can induce T cell proliferation and IL-2 secretion.

EXAMPLE 2

5 Activated Human Splenic B Cells Express CTLA4 Ligand(s) Distinct from B7-1

In light of the above observations, whether other CTLA4 binding counter-receptors were expressed on activated B cells was determined. To this end, human splenic B cells were activated for 72 hours with anti-Ig and then stained with an anti-B7-1 monoclonal antibody (B1.1) which does not inhibit B7-1 mediated costimulation. Fluorescein isothiocyanate (FITC) and mAb B1.1 were used with flow cytometric cell sorting to isolate B7-1⁺ and B7-1⁻ fractions. The resulting post-sort positive population was 99% B7-1⁺ and the post-sort negative population was 98% B7-1⁻ (Figure 2).

To examine the costimulatory potential of each population, human splenic CD28⁺ T cells were submitogenically stimulated with anti-CD3 monoclonal antibody in the presence of irradiated B7-1⁺ or B7-1⁻ anti-Ig activated (72 hours) splenic B cells. ³H-Thymidine incorporation was assessed for the last 15 hours of a 72 hours culture. IL-2 was assessed by ELISA in supernatants after 24 hours of culture (Detection limits of the assay: 31-2000 pg/ml). The results of Figure 3 are representative of ten experiments. B7-1⁺ B cells induced anti-CD3 activated T cells to proliferate and secrete IL-2 (Figure 3a) but not IL-4. As was observed with the unfractionated activated B cell population, anti-B7-1 monoclonal antibody (133) inhibited proliferation only 50% but consistently abrogated IL-2 secretion. As above, CTLA4Ig binding or blockade of CD28 with Fab anti-CD28 monoclonal antibody completely inhibited both proliferation and IL-2 secretion. Control monoclonal antibody and control-Ig were not inhibitory. In an attempt to identify other potential CTLA4/CD28 binding costimulatory ligand(s) which might account for the residual, non-B7 mediated proliferation delivered by B7⁺ B cells, the effect of BB-1 monoclonal antibody on proliferation and IL-2 secretion was examined. As seen, BB-1 monoclonal antibody completely inhibited both proliferation and IL-2 secretion (Figure 3a). Figure 3b displays the costimulatory potential of B7-1- activated human splenic B cells. Irradiated B7-1- activated (72 hr) B cells could also deliver a significant costimulatory signal to submitogenically activated CD4⁺ lymphocytes. This costimulation was not accompanied by detectable IL-2 (Figure 3b) or IL-4 accumulation and anti-B7-1 monoclonal antibody did not inhibit proliferation. However, CTLA4Ig, Fab anti-CD28 monoclonal antibody, and BB-1 monoclonal antibody all completely inhibited proliferation.

Phenotypic analysis of the B7-1⁺ and B7-1⁻ activated splenic B cells confirmed the above functional results. Figure 4 shows the cell surface expression of B7-1,

B7-2 and B7-3 on fractionated B7-1⁺ and B7-1⁻ activated B cell. As seen in Figure 4, B7-1⁺ activated splenic B cells stained with anti-B7-1 (133) monoclonal antibody, BB-1 monoclonal antibody, and bound CTLA4-Ig. In contrast, B7-⁻ activated splenic B cells did not stain with anti-B7-1 (133) monoclonal antibody but did stain with BB-1 monoclonal antibody and CTLA4Ig. These phenotypic and functional results demonstrate that both B7-1⁺ and B7-1⁻ activated (72 hours) human B lymphocytes express CTLA4 binding counter-receptor(s) which: 1) can induce submitogenically activated T cells to proliferate without detectable IL-2 secretion; and 2) are identified by the BB-1 monoclonal antibody but not anti-B7-1 monoclonal antibody. Thus, these CTLA4/CD28 ligands can be distinguished on the basis of their temporal expression after B cell activation and their reactivity with CTLA4Ig and anti-B7 monoclonal antibodies. The results of Figure 4 are representative of five experiments.

EXAMPLE 3

15 Three Distinct CTLA4/CD28 Ligands Are Expressed Following Human B Cell Activation

To determine the sequential expression of CTLA4 binding counter-receptors following activation, human splenic B cells were activated by crosslinking of either surface Ig or MHC class II and the expression of B7-1, B7-3 and B7-2 binding proteins were examined by flow cytometric analysis. Ig or MHC class II crosslinking induced a similar pattern of CTLA4Ig binding (Figures 5 and 6). Figure 5 is representative of the results of 25 experiments for anti-B7-1 and BB-1 binding and 5 experiments for CTLA4Ig binding. Figure 6 is representative of 25 experiments for anti-B7-1 binding and 5 experiments for CTLA4Ig binding. The results of these experiments indicates that prior to 24 hours, none of these molecules are expressed. At 24 hours post-activation, the majority of cells express a protein that binds CTLA4Ig (B7-2), however, fewer than 20% express either B7-1 or B7-3. Crosslinking of MHC class II induces maximal expression and intensity of B7-1 and B7-3 at 48 hours whereas crosslinking of Ig induces maximal expression at 72 hours and expression declines thereafter. These results suggest that an additional CTLA4 binding counter-receptor is expressed by 24 hours and that the temporal expression of the distinct B7-1 and B7-3 proteins appears to coincide.

A series of experiments was conducted to determine whether the temporal expression of CTLA4 binding counter-receptors differentially correlated with their ability to costimulate T cell proliferation and/or IL-2 secretion. Human splenic CD28⁺ T cells submitogenically stimulated with anti-CD3 were cultured for 72 hours in the presence of irradiated human splenic B cells that had been previously activated *in vitro* by sIg

crosslinking for 24, 48, or 72 hours. IL-2 secretion was assessed by ELISA in supernatants after 24 hours and T cell proliferation as assessed by ³H-thymidine incorporation for the last 15 hours of a 72 hour culture. The results of Figure 7 are representative of 5 experiments. As seen in Figure 7a, 24 hour activated B cells provided a costimulatory signal which was accompanied by modest levels of IL-2 production, although the magnitude of proliferation was significantly less than observed with 48 and 72 hours activated human B cells (note differences in scale for ³H-Thymidine incorporation). Neither proliferation nor IL-2 accumulation was inhibited by anti-B7-1 (133) or BB-1. In contrast, with CTLA4Ig and anti-CD28 Fab monoclonal antibody totally abrogated proliferation and IL-2 accumulation. B cells activated for 48 hours, provided costimulation which resulted in nearly maximal proliferation and IL-2 secretion (Figure 7b). Here, anti-B7-1 (133) monoclonal antibody, inhibited proliferation approximately 50% but totally blocked IL-2 accumulation. BB-1 monoclonal antibody totally inhibited both proliferation and IL-2 secretion. As above, CTLA4Ig and Fab anti-CD28 also totally blocked proliferation and IL-2 production. Finally, 72 hour activated B cells induced T cell response identical to that induced by 48 hour activated B cells. Similar results are observed if the submitogenic signal is delivered by phorbol myristic acid (PMA) and if the human splenic B cells are activated by MHC class II rather than Ig crosslinking. These results indicate that there are three CTLA4 binding molecules that are temporarily expressed on activated B cells and each can induce submitogenically stimulated T cells to proliferate. Two of these molecules, the early CTLA4 binding counter-receptor (B7-2) and B7-1 (133) induce IL-2 production whereas B7-3 induces proliferation without detectable IL-2 production.

Previous studies provided conflicting evidence whether the anti-B7 monoclonal antibody, 133 and monoclonal antibody BB-1 identified the same molecule (Freedman, A.S. et al. (1987) *Immunol.* 137, 3260-3267; Yokochi, T., et al. (1982) *J. Immunol.* 128, 823-827; Freeman, G.J., et al. (1989) *J. Immunol.* 143, 2714-2722.). Although both monoclonal antibodies identified molecules expressed 48 hours following human B-cell activation, several reports suggested that B7 (B7-1) and the molecule identified by monoclonal antibody BB-1 were distinct since they were differentially expressed on cell lines and B cell neoplasms (Freedman, A.S. et al. (1987) *Immunol.* 137, 3260-3267; Yokochi, T., et al. (1982) *J. Immunol.* 128, 823-827; Freeman, G.J., et al. (1989) *J. Immunol.* 143, 2714-2722; Clark, E and Yokochi, T. (1984) *Leukocyte Typing, 1st International References Workshop*. 339-346; Clark, E., et al. (1984) *Leukocyte Typing, 1st International References Workshop*. 740). In addition, immunoprecipitation and Western Blotting with these IgM monoclonal antibodies suggested that they identified different molecules (Clark, E and Yokochi, T. (1984) *Leukocyte Typing, 1st International References Workshop*. 339-346; Clark, E., et al. (1984) *Leukocyte Typing, 1st International References Workshop*. 740). The

original anti-B7 monoclonal antibody, 133, was generated by immunization with anti-immunoglobulin activated human B lymphocytes whereas the BB-1 monoclonal antibody was generated by immunization with a baboon cell line. Thus, the BB-1 monoclonal antibody must identify an epitope on human cells that is conserved between baboons and humans.

Following the molecular cloning and expression of the human B7 gene (B7-1), B7 transfected COS cells were found to be identically stained with the anti-B7 (133) and BB-1 monoclonal antibodies and that they both precipitated the identical broad molecular band (44-54kD) strongly suggesting that they identified the same molecule (Freeman, G.J., et al. (1989) *J. Immunol.* 143, 2714-2722). This observation was unexpected since the gene encoding the molecule identified by the BB-1 monoclonal antibody had been previously mapped to chromosome 12 (Katz, F.E., et al. (1985) *Eur. J. Immunol.* 103-6), whereas the B7 gene was located by two groups on chromosome 3 (Freeman, G.J., et al. (1992) *Blood* 79, 489-494; Selvakumar, A., et al. (1992) *Immunogenetics* 36, 175-181.). Subsequently, additional discrepancies between the phenotypic expression of B7 (B7-1) and the molecule identified by the BB-1 monoclonal antibody were noted. BB-1 monoclonal antibody stained thymic epithelial cells (Turka, L.A., et al. (1991) *J. Immunol.* 146, 1428-36; Munro, J.M., et al. *Blood* submitted.) and keratinocytes (Nickoloff, B., et al (1993) *Am. J. Pathol.* 142, 1029-1040; Augustin, M., et al. (1993) *J. Invest. Dermatol.* 100, 275-281.) whereas anti-B7 did not. Recently, Nickoloff et al. (1993) *Am. J. Pathol.* 142, 1029-1040, reported discordant expression of the molecule identified by the BB-1 monoclonal antibody and B7 on keratinocytes using a BB-1 and anti-B7 (B1.1 and 133) monoclonal antibodies. Nickoloff et al. also demonstrated that these BB-1 positive cells did not express B7 mRNA yet bound CD28 transfected COS cells providing further support for the existence of a distinct protein which binds monoclonal antibody BB-1.

The present findings confirm that there is an additional CTLA4 counter-receptor identified by the BB-1 monoclonal antibody, B7-3, and that this protein appears to be functionally distinct from B7-1 (133). Although the expression of B7-1 and B7-3 following B cell activation appears to be concordant on B7 positive B cells, these studies demonstrate that the B7-3 molecule is also expressed on B7 negative activated B cells. More importantly, the B7-3 molecule appears to be capable of inducing T cell proliferation without detectable IL-2 or IL-4 production. This result is similar to the previous observation that ICAM-1 could costimulate T cell proliferation without detectable IL-2 or IL-4 production (Boussiotis, V., et al *J. Exp. Med.* (accepted for publication)). These data indicate that the BB-1 monoclonal antibody recognizes an epitope on the B7-1 protein and that this epitope is also found on a distinct B7-3 protein, which also has costimulatory function. Phenotypic and blocking studies demonstrate that the BB-1 monoclonal antibody could detect one (on B7

negative cells) or both (on B7 positive cells) of these proteins. In contrast, the anti-B7 monoclonal antibodies, 133 and B1.1 detect only the B7-1 protein. Taken together, these results suggest that by 48 hours post B-cell activation by crosslinking of surface immunoglobulin or MHC class II, B cells express at least two distinct CTLA4 binding counter-receptors, one identified by both anti-B7 and BB-1 monoclonal antibodies and the other identified only by BB-1 monoclonal antibody.

The B7-2 antigen is not detectable on activated B cells after 12 hours, but by 24 hours it is strongly expressed and functional. This molecule appears to signal via CD28 since proliferation and IL-2 production are completely blocked by Fab anti-CD28 monoclonal antibody. At 48 hours post activation, IL-2 secretion seems to be accounted for by B7-1 costimulation, since anti-B7 monoclonal antibody completely inhibits IL-2 production.

Previous studies and results presented here demonstrate that B7 (B7-1) is neither expressed (Freedman, A.S. et al. (1987) *Immunol.* 137, 3260-3267; Freedman, A.S., et al. (1991) *Cell. Immunol.* 137, 429-437) nor capable of costimulating T cell proliferation or IL-2 secretion until 48 hours post B-cell activation. Previous studies have shown that activation of T cells via the TCR in the absence of costimulation (Gimmi, C.D., et al. (1993) *Proc. Natl. Acad. Sci USA* 90:6586-6590; Schwartz, R.H., et al. (1989) *Cold Spring Harb. Symp. Quant. Biol* 54, 605-10; Beverly, B., et al. (1992) *Int. Immunol.* 4, 661-671.) and lack of IL-2 (Boussiotis, V., et al *J. Exp. Med.* (submitted); Beverly, B., et al. (1992) *Int. Immunol.* 4, 661-671; Wood, M., et al. (1993) *J. Exp. Med.* 177, 597-603) results in anergy. If B7-1 were the only costimulatory molecule capable of inducing IL-2 secretion, T cells would be anergized within the first 24 hours following activation since there is no B7-1 present to costimulate IL-2 production. Therefore, the existence of another, early inducible costimulatory molecule, which can costimulate IL-2 secretion during the first 24 hours would be necessary to induce an effective immune response rather than anergy. The appearance of the early CTLA4 binding counter-receptor, B7-2, between 12 and 24 hours post B cell activation, fulfills this function.

Two observations shed light on the biologic and potential clinical significance of these two additional CTLA4 binding counter-receptors. First, B7 (B7-1) deficient mouse has been developed and its antigen presenting cells were found to still bind CTLA4Ig (Freeman and Sharpe manuscript in preparation). This mouse is viable and isolated mononuclear cells induce detectable levels of IL-2 when cultured with T cells *in vitro*. Therefore, an alternative CD28 costimulatory counter-receptor or an alternative IL-2 producing pathway must be functional. Second, thus far the most effective reagents to induce antigen specific anergy in murine and human systems are CTLA4Ig and Fab anti-CD28, whereas anti-B7 monoclonal antibodies have been much less effective (Harding, F.A., et al.

(1992) *Nature*. 356, 607-609; Lenschow, D.J., et al. (1992) *Science*. 257, 789-792; Chen, L., et al. (1992) *Cell*. 71, 1093-1102; Tan, P., et al. (1993) *J. Exp. Med.* 177, 165-173.). These observations are also consistent with the hypothesis that alternative CTLA4/CD28 ligands capable of inducing IL-2 exist, and taken together with the results presented herein, suggest that all three CTLA4 binding counter-receptors may be critical for the induction of T cell immunity. Furthermore, their blockade will likely be required for the induction of T cell anergy. Identical results have been observed in the murine system with the identification of two CTLA4 binding ligands, corresponding to the human B7-1 and B7-2 molecules. APCs in the B7 deficient mouse bind to the CTLA4 and can induce IL-2 secretion. Taken together, these observations indicate that multiple CTLA-4 binding counter-receptors exist and sequentially costimulate T cell activation in the murine system.

EXAMPLE 4

Cloning, Sequencing and Expression of the B7-2 Antigen

A. Construction of cDNA Library

A cDNA library was constructed in the pCDM8 vector (Seed, *Nature*, 329:840 (1987)) using poly (A)⁺ RNA from the human anti-IgM activated B cells as described (Aruffo et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)). Splenic B cells were cultured at 2x10⁶ cells/ml in complete culture media, {RPMI 1640 with 10% heat inactivated fetal calf serum (FCS), 2mM glutamine, 1 mM sodium pyruvate, penicillin (100 units/ml), streptomycin sulfate (100µg/ml) and gentamycin sulfate (5µg/ml)}, in tissue culture flasks and were activated by crosslinking of sIg with affinity purified rabbit anti-human IgM coupled to Affi-Gel 702 beads (Bio-Rad), Richmond, CA) (Boyd, A.W., et al., (1985) *J. Immunol.* 134,1516). Activated B cells were harvested after 1/6, 1/2, 4, 8 12, 24, 48, 72 and 96 hours.

RNA was prepared by homogenizing activated B cells in a solution of 4M guanidine thiocyanate, 0.5% sarkosyl, 25mM EDTA, pH 7.5, 0.13% Sigma anti-foam A, and 0.7% mercaptoethanol. RNA was purified from the homogenate by centrifugation for 24 hour at 32,000 rpm through a solution of 5.7M CsCl, 10mM EDTA, 25mM Na acetate, pH 7. The pellet of RNA was dissolved in 5% sarkosyl, 1mM EDTA, 10mM Tris, pH 7.5 and extracted with two volumes of 50% phenol, 49% chloroform, 1% isoamyl alcohol. RNA was ethanol precipitated twice. Poly (A)⁺ RNA used in cDNA library construction was purified by two cycles of oligo (dT)-cellulose selection.

Complementary DNA was synthesized from 5.5µg of anti-IgM activated human B cell poly(A)⁺ RNA in a reaction containing 50mM Tris, pH 8.3, 75mM KCl, 3mM MgCl₂, 10mM dithiothreitol, 500µM dATP, dCTP, dGTP, dTTP, 50µg/ml oligo(dT)₁₂₋₁₈,

180 units/ml RNasin, and 10,000 units/ml Moloney-MLV reverse transcriptase in a total volume of 55 μ l at 37° for 1 hr. Following reverse transcription, the cDNA was converted to double-stranded DNA by adjusting the solution to 25mM Tris, pH 8.3, 100mM KCl, 5mM MgCl₂, 250 μ M each dATP, dCTP, dGTP, dTTP, 5mM dithiothreitol, 250 units/ml DNA polymerase I, 8.5 units/ml ribonuclease H and incubating at 16° for 2 hr. EDTA was added to 18mM and the solution was extracted with an equal volume of 50% phenol, 49% chloroform, 1% isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate and with 4 micrograms of linear polyacrylamide as carrier. In addition, cDNA was synthesized from 4 μ g of anti-IgM activated human B cell poly(A)⁺ RNA in a reaction containing 50mM Tris, pH 8.8, 50 μ g/ml oligo(dT)₁₂₋₁₈, 327 units/ml RNasin, and 952 units/ml AMV reverse transcriptase in a total volume of 100 μ l at 42° for 0.67 hr. Following reverse transcription, the reverse transcriptase was inactivated by heating at 70° for 10 min. The cDNA was converted to double-stranded DNA by adding 320 μ l H₂O and 80 μ l of a solution of 0.1M Tris, pH 7.5, 25mM MgCl₂, 0.5M KCl, 250 μ g/ml bovine serum albumin, and 50mM dithiothreitol, and adjusting the solution to 200 μ M each dATP, dCTP, dGTP, dTTP, 50 units/ml DNA polymerase I, 8 units/ml ribonuclease H and incubating at 16° C for 2 hours. EDTA was added to 18 mM and the solution was extracted with an equal volume of 50 % phenol, 49 % chloroform, 1 % isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate and with 4 micrograms of linear polyacrylamide as carrier.

The DNA from 4 μ g of AMV reverse transcription and 2 μ g of Moloney MLV reverse transcription was combined. Non-selfcomplementary BstXI adaptors were added to the DNA as follows: The double-stranded cDNA from 6 μ g of poly(A)⁺ RNA was incubated with 3.6 μ g of a kinased oligonucleotide of the sequence CTTTAGAGCACA (SEQ ID NO:15) and 2.4 μ g of a kinased oligonucleotide of the sequence CTCTAAAG (SEQ ID NO:16) in a solution containing 6mM Tris, pH 7.5, 6mM MgCl₂, 5mM NaCl, 350 μ g/ml bovine serum albumin, 7mM mercaptoethanol, 0.1mM ATP, 2mM dithiothreitol, 1mM spermidine, and 600 units T4 DNA ligase in a total volume of 0.45ml at 15° C for 16 hours. EDTA was added to 34mM and the solution was extracted with an equal volume of 50% phenol, 49% chloroform, 1% isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate.

DNA larger than 600bp was selected as follows: The adapted DNA was redissolved in 10mM Tris, pH 8, 1mM EDTA, 600mM NaCl, 0.1% sarkosyl and chromatographed on a Sepharose CL-4B column in the same buffer. DNA in the void volume of the column (containing DNA greater than 600bp) was pooled and ethanol precipitated.

The pCDM8 vector was prepared for cDNA cloning by digestion with BstXI and purification on an agarose gel. Adapted DNA from 6µg of poly(A)⁺RNA was ligated to 2.25µg of BstXI cut pCDM8 in a solution containing 6mM Tris, pH 7.5, 6mM MgCl₂, 5mM NaCl, 350µg/ml bovine serum albumin, 7mM mercaptoethanol, 0.1mM ATP, 2mM dithiothreitol, 1mM spermidine, and 600 units T4 DNA ligase in a total volume of 1.5ml at 15° for 24 hr. The ligation reaction mixture was transformed into competent E.coli MC1061/P3 and a total of 4,290,000 independent cDNA clones were obtained.

Plasmid DNA was prepared from a 500 ml culture of the original transformation of the cDNA library. Plasmid DNA was purified by the alkaline lysis procedure followed by twice banding in CsCl equilibrium gradients (Maniatis et al, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, NY (1987)).

B. Cloning Procedure

In the first round of screening, thirty 100 mm dishes of 50% confluent COS cells were transfected with 0.05µg/ml anti-IgM activated human B cells library DNA using the DEAE-Dextran method (Seed et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)). The cells were trypsinized and re-plated after 24 hours. After 47 hours, the cells were detached by incubation in PBS/0.5 mM EDTA, pH 7.4/0.02% Na azide at 37°C for 30 min. The detached cells were treated with 10 µg/ml/CTLA4Ig and CD28Ig for 45 minutes at 4°C. Cells were washed and distributed into panning dishes coated with affinity-purified Goat anti-human IgG antibody and allowed to attach at room temperature. After 3 hours, the plates were gently washed twice with PBS/0.5mM EDTA, pH 7.4/0.02% Na azide, 5% FCS and once with 0.15M NaCl, 0.01 M Hepes, pH 7.4, 5% FCS. Episomal DNA was recovered from the panned cells and transformed into *E. coli* DH10B/P3. The plasmid DNA was re-introduced into COS cells via spheroplast fusion as described (Seed et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)) and the cycle of expression and panning was repeated twice. In the second and third rounds of selection, after 47 hours, the detached COS cells were first incubated with α-B7-1 mAbs (133 and B1.1, 10 µg/ml), and COS cells expressing B7-1 were removed by α-mouse IgG and IgM coated magnetic beads. COS cells were then treated with 10 µg/ml of human CTLA4Ig (hCTLA4Ig) and human CD28Ig (hCD28Ig) and human B7-2 expressing COS cells were selected by panning on dishes with goat anti-human IgG antibody plates. After the third round, plasmid DNA was prepared from individual colonies and transfected into COS cells by the DEAE-Dextran method. Expression of B7-2 on transfected COS cells was analyzed by indirect immunofluorescence with CTLA4Ig.

After the final round of selection, plasmid DNA was prepared from individual colonies. A total of 4 of 48 candidate clones contained a cDNA insert of approximately 1.2 kb. Plasmid DNA from these four clones was transfected into COS cells. All four clones

were strongly positive for B7-2 expression by indirect immunofluorescence using CTLA4Ig and flow cytometric analysis.

C. Sequencing

5 The B7-2 cDNA insert in clone29 was sequenced in the pCDM8 expression vector employing the following strategy. Initial sequencing was performed using sequencing primers T7, CDM8R (Invitrogen) homologous to pCDM8 vector sequences adjacent to the cloned B7-2 cDNA (see Table I). Sequencing was performed using dye terminator chemistry and an ABI automated DNA sequencer. (ABI, Foster City, CA). DNA sequence obtained
10 using these primers was used to design additional sequencing primers (see Table I). This cycle of sequencing and selection of additional primers was continued until the B7-2 cDNA was completely sequenced on both strands.

TABLE I

15	T7(F) (SEQ ID NO:3)	5'd[TAATACGACTCACTATAGGG]3'
	CDM8(R) (SEQ ID NO:4)	5'd[TAAGGTTCCCTTCACAAAG]3'
	CDM8 RGV(2) (SEQ ID NO:5)	5'd[ACTGGTAGGTATGGAAGATCC]3'
	HBX29-5P (2R) (SEQ ID NO:6)	5'd[ATGCGAATCATTCCCTGTGGGC]3'
20	HBX29-5P (2F) (SEQ ID NO:7)	5'd[AAAGCCCACAGGAATGATTCTG]3'
	HBX29-5P (SEQ ID NO:8)	5'd[CTCTCAAAACCAAABCCTGAG]3'
	5PA (SEQ ID NO:9)	5'd[TTAGGTCACAGCAGAAGCAGC]3'
	5PA (3FA) (SEQ ID NO:10)	5'd[TCTGGAAACTGACAAGACGCG]3'
	HBX29-5P(1R) (SEQ ID NO:11)	5'd[CTCAGGCTTTGGTTTTGAGAG]3'
25	HBX29-3P(1R) (SEQ ID NO:12)	5'd[CACTCTCTCCCTCTCCATTG]3'
	HBX29-5P(3R) (SEQ ID NO:13)	5'd[GACAAGCTGATGGAAACGTCG]3'
	HBX29-3P(1P) (SEQ ID NO:14)	5'd[CAATGGAGAGGGAAGAGAGTG]3'

30 The human B7-2 clone 29 contained an insert of 1,120 base pairs with a single long open reading frame of 987 nucleotides and approximately 27 nucleotides of 3' noncoding sequences (Figure 8 (SEQ ID NO:1)). The predicted amino acid sequence encoded by the open reading frame of the protein is shown below the nucleotide sequence in Figure 8. The encoded protein, human B7-2, is predicted to be 329 amino acids in length
35 (SEQ ID NO:2). This protein sequence exhibits many features common to other type 1 Ig superfamily embrane proteins. Protein translation is predicted to begin at the ATG codon (nucleotide 107-109) based on DNA homology in this region with the consensus eukaryotic

translation initiation site (Kozak, M. (1987) *Nucl. Acids Res.* 15:8125-8148). The amino terminus of the human B7-2 protein (amino acids 1 to 23) has the characteristics of a secretory signal peptide with a predicted cleavage between the alanines at positions 23 and 24 (von Heijne (1986) *Nucl. Acids Res.* 14:4683). Processing at this site would result in a

5 human B7-2 membrane bound protein of 306 amino acid with an unmodified molecular weight of approximately 34 kDa. This protein would consist of an extracellular Ig superfamily V and C like domains, of from about amino acid residue 24-245, a hydrophobic transmembrane domain of from about amino acid residue 246-268 and a long cytoplasmic domain of from about amino acid residue 269-329. The homologies to the Ig superfamily are

10 due to the two contiguous Ig-like domains in the extracellular region bound by the cysteines at positions 40 to 110 and 157 to 218. The extracellular domain also contains eight potential N-linked glycosylation sites. *E. coli* transfected with a vector containing the cDNA insert of clone 29, encoding the human B7-2 protein, was deposited with the American Type Culture Collection (ATCC) on July 26, 1993 as Accession No. 69357.

15 Comparison of both the nucleotide and amino acid sequences of human B7-2 with the GenBank and EMBL databases showed that only the human and murine B7-1 proteins are related. Alignment of the three B7 protein sequences (see Figure 13) shows that human B7-2 has approximately 26% amino acid identity with human B7-1. Figure 13 represents the comparison of the amino acid sequences for human B7-2 (hB7-2) (SEQ ID

20 NO:2), human B7-1 (hB7-1) (SEQ ID NO: 28 and 29) and murine B7 (mB7) (SEQ ID NO: 30 and 31). The amino acid sequences for the human B7-1 and murine B7 (referred to herein as murine B7-1) can be found in Genbank at Accession #M27533 and X60958 respectively. Vertical lines in Figure 13 show identical amino acids between the hB7-2 and hB7-1 or mB7. Identical amino acids between hB7-1 and mB7 are not shown. The hB7-2 protein exhibits

25 the same general structure as hB7-1 as defined by the common cysteines (positions 40 and 110, IgV domains; positions 157 and 217, IgC domain) which the Ig superfamily domains and by many other common amino acids. Since both hB7-1 and mB7 have been shown to bind to both human CTLA4 and human CD28, the amino acids in common between these two related proteins will be those necessary to comprise a CTLA4 or CD28 binding

30 sequence. An example of such a sequence would be the KYMGRTSFD (position 81-89, hB7-2) (SEQ ID NO:17) or KSQDNVTELYDVS (position 188-200, hB7-2) (SEQ ID NO:18). Additional related sequences are evident from the sequence comparison and others can be inferred by considering homologous related amino acids such as aspartic acid and glutamic acid, alanine and glycine and other recognized functionally related amino acids.

35 The B7 sequences share a highly positive charged domain with the cytoplasmic portion WKWKKKKRPRNSYKC (position 269-282, hB7-2) (SEQ ID NO:19) which is probably involved in intracellular signaling.

EXAMPLE 5**Characterization of the Recombinant B7-2 Antigen****5. A. B7-2 Binds CTLA4Ig and Not Anti-B7-1 and Anti-B7-3 Monoclonal Antibodies**

COS cells transfected with either vector DNA (pCDNAI), or an expression plasmid containing B7-1 (B7-1) or B7-2 (B7-2) were prepared. After 72 hours, the transfected COS cells were detached by incubation in PBS containing 0.5 mM EDTA and 0.02% Na azide for 30 min. at 37°C. Cells were analyzed for cell surface expression by indirect immunofluorescence and flow cytometric analysis using fluorescein isothiocyanate conjugated (FITC) goat-anti-mouse Ig or goat-anti-human IgG FITC (Figure 9). Cell surface expression of B7-1 was detected with mAbs 133 (anti-B7-1) and BB-1 (anti-B7-1 and anti-B7-3) and with CTLA4Ig, whereas B7-2 reacted only with CTLA4Ig. Neither of the B7 transfectants showed any staining with the isotype controls (IgM or control Ig). The vector transfected COS cells showed no staining with any of the detection reagents. In addition, none of the cells showed any staining with the FITC labeled detection reagents and alone. This demonstrates that B7-2 encodes a protein that is a CTLA4 counter-receptor but is distinct from B7-1 and B7-3.

20 B. RNA Blot Analysis of B7-2 Expression in Unstimulated and Activated Human B Cells, Cell Lines, and Myelomas

Human splenic B cells were isolated by removing T cells and monocytes as previously described (Freedman, A.S., Freeman, G.J., Horowitz, J.C., Daley, J., Nadler, L.M., *J. Immunol.* (1987) 137:3260-3267). Splenic B cells were activated using anti-Ig beads and cells were harvested at the indicated times (Freedman et al., (1987), cited supra). Human myelomas from bone marrow specimens were enriched by removing T cells and monocytes using E rosettes and adherence as previously described (Freeman, G.J., et al., *J. Immunol.* (1989) 143:2714-2722). RNA was prepared by guanidine thiocyanate homogenization and cesium chloride centrifugation. Equal amounts of RNA (20µg) were electrophoresed on an agarose gel, blotted, and hybridized to ³²P-labelled B7-2 cDNA. Figure 10, panel a, shows RNA blot analysis of unstimulated and anti-Ig activated human splenic B cells and of cell lines including Raji (B cell Burkitt's lymphoma), Daudi (B cell Burkitt's lymphoma), RPMI 8226 (myeloma), K562 (erythroleukemia), and REX (T cell acute lymphoblastic leukemia). Figure 10, panel b shows RNA blot analysis of human myeloma specimens.

35 Three mRNA transcripts of 1.35, 1.65 and 3.0 kb were identified by hybridization to the B7-2 cDNA (Figure 10, panel b). RNA blot analysis demonstrated that B7-2 mRNA is expressed in unstimulated human splenic B cells and increases 4-fold

following activation (Figure 10, panel a). B7-2 mRNA was expressed in B cell neoplastic lines (Raji, Daudi) and a myeloma (RPMI 8226) but not in the erythroleukemia K562 and the T cell line REX. In contrast, we have previously shown that B7-1 mRNA is not expressed in resting B cells and is transiently expressed following activation (G.J. Freeman et al. (1989) supra). Examination of mRNA isolated from human myelomas demonstrates that B7-2 mRNA is expressed in 6 of 6 patients, whereas B7-1 was found in only 1 of these 6 (G.J. Freeman et al. (1989) supra). Thus, B7-1 and B7-2 expression appears to be independently regulated.

10 C. Costimulation

Human CD28⁺ T cells were isolated by immunomagnetic bead depletion using monoclonal antibodies directed against B cells, natural killer cells and macrophages as previously described (Gimmi, C.D., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90, 6586-6590). B7-1, B7-2 and vector transfected COS cells were harvested 72 hours after transfection, 15 incubated with 25µg/ml of mitomycin-C for 1 hour, and then extensively washed. 10⁵ CD28⁺ and T cells were incubated with 1 ng/ml of phorbol myristic acid (PMA) and the indicated number of COS transfectants (Figure 11). As shown in Figure 11, panel a, T cell proliferation was measured by 3H-thymidine (1 µCi) incorporated for the last 12 hours of a 72 hour incubation. Panel b of Figure 11 shows IL-2 production by T cells as measured by 20 ELISA (Biosource, CA) using supernatants harvested 24 hours after the initiation of culture.

D. B7-2 Costimulation is not Blocked by Anti-B7-1 and Anti-B7-3 mAbs but is Blocked by CTLA4-Ig and Anti-CD28 Fab

Human CD28⁺ T cells were isolated by immunomagnetic bead depletion 25 using mAbs directed against B cells, natural killer cells, and macrophages as previously described (Gimmi, C.D., Freeman, G.J., Gribben, J.G., Gray, G., Nadler, L.M. (1993) *Proc. Natl. Acad. Sci USA* 90, 6586-6590). B7-1, B7-2, and vector transfected COS cells were harvested 72 hours after transfection, incubated with 25µg/ml of mitomycin-C for 1 hour, and then extensively washed. 10⁵ CD28⁺ T cells were incubated with 1 ng/ml of phorbol 30 myristic acetate (PMA) and 2 x 10⁴ COS transfectants. Blocking agents (10µg/ml) are indicated on the left side of Figure 12 and include: 1) no monoclonal antibody (no blocking agents), 2) mAb 133 (anti-B7-1 mAb), 3) mAb BB1 (anti-B7-1 and anti-B7-3 mAb), 4) mAb B5 (control IgM mAb), 5) anti-CD28 Fab (mAb 9.3), 6) CTLA-Ig, and 7) control Ig. Panel a of Figure 12 shows proliferation measured by 3H-thymidine (1µCi) incorporation for the last 35 12 hours of a 72 hour incubation. Figure 12, panel b, shows IL-2 production as measured by ELISA (Biosource, CA) using supernatants harvested 24 hours after the initiation of culture.

B7-1 and B7-2 transfected COS cells costimulated equivalent levels of T cell proliferation when tested at various stimulator to responder ratios (Figure 11). Like B7-1, B7-2 transfected COS cell costimulation resulted in the production of IL-2 over a wide range of stimulator to responder cell ratios (Figure 11). In contrast, vector transfected COS cells
 5 did not costimulate T cell proliferation or IL-2 production.

E. B7-2 Costimulation is not Blocked by Anti-B7-1 and Anti-B7-3 mAbs but is Blocked by CTLA4-Ig and Anti-CD28 Fab

Human CD28⁺ T cells were isolated by immunomagnetic bead depletion
 10 using mAbs directed against B cells, natural killer cells, and macrophages as previously described (Gimmi, C.D., Freeman, G.J., Gribben, J.G., Gray, G., Nadler, L.M. (1993) *Proc. Natl. Acad. Sci USA* 90, 6586-6590). B7-1, B7-2, and vector transfected COS cells were harvested 72 hours after transfection, incubated with 25µg/ml of mitomycin-C for 1 hour, and then extensively washed. 10⁵ CD28⁺ T cells were incubated with 1 ng/ml of phorbol
 15 myristic acetate (PMA) and 2 x 10⁴ COS transfectants. Blocking agents (10µg/ml) are indicated on the left side of Figure 12 and include: 1) no monoclonal antibody (no blocking agents), 2) mAb 133 (anti-B7-1 mAb), 3) mAb BB1 (anti-B7-1 and anti-B7-3 mAb), 4) mAb B5 (control IgM mAb), 5) anti-CD28 Fab (mAb 9.3), 6) CTLA-Ig, and 7) control Ig. Panel a of Figure 12 shows proliferation measured by ³H-thymidine (1µCi) incorporation for the last
 20 12 hours of a 72 hour incubation. Figure 12, panel b, shows IL-2 production as measured by ELISA (Biosource, CA) using supernatants harvested 24 hours after the initiation of culture.

To distinguish B7-2 from B7-1 and B7-3, mAbs directed against B7-1 and B7-3 were used to inhibit proliferation and IL-2 production of submitogenically activated human CD28⁺ T cells. Both B7-1 and B7-2 COS transfectants costimulated T cell proliferation and
 25 IL-2 production (Figure 12). MAbs 133 (Freedman, A.S. et al. (1987) *supra*) (anti-B7-1) and BB1 (Boussiotis, V.A., et al., (in review) *Proc. Natl. Acad. Sci. USA*; Yokochi, T., Holly, R.D., Clark, E.A. (1982) *J. Immunol.* 128, 823-827) (anti-B7-1 and anti-B7-3) completely inhibited proliferation and IL-2 secretion induced by B7-1 but had no effect upon costimulation by B7-2 transfected COS cells. Isotype matched control B5 mAb had no
 30 effect. To determine whether B7-2 signals via the CD28/CTLA4 pathway, anti-CD28 Fab and CTLA4-Ig fusion protein were tested to determine whether they inhibited B7-2 costimulation. Both anti-CD28 Fab and CTLA4-Ig inhibited proliferation and IL-2 production induced by either B7-1 or B7-2 COS transfectants whereas control Ig fusion protein had no effect (Figure 12). While CTLA4-Ig inhibited B7-2 costimulation of
 35 proliferation by only 90%, in other experiments inhibition was more pronounced (98-100%). None of the blocking agents inhibited T cell proliferation or IL-2 production induced by the combination of PMA and phytohemagglutinin.

Like B7-1, B7-2 is a counter-receptor for the CD28 and CTLA4 T cell surface molecules. Both proteins are similar in that they are: 1) expressed on the surface of APCs; 2) structurally related to the Ig supergene family with an IgV and IgC domain which share 26% amino acid identity, and 3) capable of costimulating T cells to produce IL-2 and proliferate. However, B7-1 and B7-2 differ in several fundamental ways. First, B7-2 mRNA is constitutively expressed in unstimulated B cells, whereas B7-1 mRNA does not appear until 4 hours and cell surface protein is not detected until 24 hours (Freedman, A.S., et al. (1987) *supra*; Freeman, G.J., et al. (1989) *supra*). Unstimulated human B cells do not express CTLA4 counter-receptors on the cell surface and do not costimulate T cell proliferation (Boussiotis, V.A., et al. *supra*). Therefore, expression of B7-2 mRNA in unstimulated B cells would allow rapid expression of B7-2 protein on the cell surface following activation, presumably from stored mRNA or protein. Costimulation by B7-2 transfectants is partially sensitive to paraformaldehyde fixation, whereas B7-2 costimulation is resistant (Gimmi, C.D., et al. (1991) *Proc. Natl. Acad. Sci. USA* **88**, 6575-6579). Second, expression of B7-1 and B7-2 in cell lines and human B cell neoplasms substantially differs. Third, B7-2 protein contains a longer cytoplasmic domain than B7-1 and this could play a role in signaling B-cell differentiation. These phenotypic and functional differences suggest that these homologous molecules may have biologically distinct functions.

20 **EXAMPLE 6**

Cloning and Sequencing of the Murine B7-2 Antigen

A. Construction of cDNA Library

A cDNA library was constructed in the pCDM8 vector (Seed, *Nature*, 329:840 (1987)) using poly (A)⁺ RNA from dibutyl cyclic AMP (cAMP) activated M12 cells (a murine B cell tumor line) as described (Aruffo et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)).

M12 cells were cultured at 1×10^6 cells/ml in complete culture media, {RPMI 1640 with 10% heat inactivated fetal calf serum (FCS), 2mM glutamine, 1 mM sodium pyruvate, penicillin (100 units/ml), streptomycin sulfate (100 μ g/ml) and gentamycin sulfate (5 μ g/ml)}, in tissue culture flasks and were activated by 300 μ g/ml dibutyl cAMP (Nabavi, N., et al. (1992) *Nature* **360**, 266-268). Activated M12 cells were harvested after 0, 6, 12, 18, 24 and 30 hours.

RNA was prepared by homogenizing activated M12 cells in a solution of 4M guanidine thiocyanate, 0.5% sarkosyl, 25mM EDTA, pH 7.5, 0.13% Sigma anti-foam A, and 0.7% mercaptoethanol. RNA was purified from the homogenate by centrifugation for 24 hour at 32,000 rpm through a solution of 5.7M CsCl, 10mM EDTA, 25mM Na acetate, pH 7.

The pellet of RNA was dissolved in 5% sarkosyl, 1mM EDTA, 10mM Tris, pH 7.5 and extracted with two volumes of 50% phenol, 49% chloroform, 1% isoamyl alcohol. RNA was ethanol precipitated twice. Poly (A)⁺ RNA used in cDNA library construction was purified by two cycles of oligo (dT)-cellulose selection

5. Complementary DNA was synthesized from 5.5µg of dibutyl cAMP activated murine M12 cell poly(A)⁺ RNA in a reaction containing 50mM Tris, pH 8.3, 75mM KCl, 3mM MgCl₂, 10mM dithiothreitol, 500µM dATP, dCTP, dGTP, dTTP, 50µg/ml oligo(dT)₁₂₋₁₈, 180 units/ml RNasin, and 10,000 units/ml Moloney-MLV reverse transcriptase in a total volume of 55µl at 37°C for 1 hr. Following reverse transcription, the cDNA was converted to double-stranded DNA by adjusting the solution to 25mM Tris, pH 8.3, 100mM KCl, 5mM MgCl₂, 250µM each dATP, dCTP, dGTP, dTTP, 5mM dithiothreitol, 250 units/ml DNA polymerase I, 8.5 units/ml ribonuclease H and incubating at 16°C for 2 hr. EDTA was added to 18mM and the solution was extracted with an equal volume of 50% phenol, 49% chloroform, 1% isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate and with 4 micrograms of linear polyacrylamide as carrier. Following reverse transcription, the reverse transcriptase was inactivated by heating at 70°C for 10 min. The cDNA was converted to double-stranded DNA by adding 320µl H₂O and 80µl of a solution of 0.1M Tris, pH 7.5, 25mM MgCl₂, 0.5M KCl, 250µg/ml bovine serum albumin, and 50mM dithiothreitol, and adjusting the solution to 200µM each dATP, dCTP, dGTP, dTTP, 50 units/ml DNA polymerase I, 8 units/ml ribonuclease H and incubating at 16°C for 2 hours. EDTA was added to 18 mM and the solution was extracted with an equal volume of 50% phenol, 49% chloroform, 1% isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate and with 4 micrograms of linear polyacrylamide as carrier.

25 2µg of non-selfcomplementary BstXI adaptors were added to the DNA as follows: The double-stranded cDNA from 5.5µg of poly(A)⁺ RNA was incubated with 3.6µg of a kinased oligonucleotide of the sequence CTTTAGAGCACA (SEQ ID NO:15) and 2.4µg of a kinased oligonucleotide of the sequence CTCTAAAG (SEQ ID NO:16) in a solution containing 6mM Tris, pH 7.5, 6mM MgCl₂, 5mM NaCl, 350µg/ml bovine serum albumin, 7mM mercaptoethanol, 0.1mM ATP, 2mM dithiothreitol, 1mM spermidine, and 600 units T4 DNA ligase in a total volume of 0.45ml at 15°C for 16 hours. EDTA was added to 34mM and the solution was extracted with an equal volume of 50% phenol, 49% chloroform, 1% isoamyl alcohol. DNA was precipitated with two volumes of ethanol in the presence of 2.5M ammonium acetate.

35 DNA larger than 600bp was selected as follows: The adapted DNA was redissolved in 10mM Tris, pH 8, 1mM EDTA, 600mM NaCl, 0.1% sarkosyl and chromatographed on a Sepharose CL-4B column in the same buffer. DNA in the void

volume of the column (containing DNA greater than 600bp) was pooled and ethanol precipitated.

The pCDM8 vector was prepared for cDNA cloning by digestion with BstXI and purification on an agarose gel. Adapted DNA from 5.5µg of poly(A)⁺RNA was
5 ligated to 2.25µg of BstXI cut pCDM8 in a solution containing 6mM Tris, pH 7.5, 6mM MgCl₂, 5mM NaCl, 350µg/ml bovine serum albumin, 7mM mercaptoethanol, 0.1mM ATP, 2mM dithiothreitol, 1mM spermidine, and 600 units T4 DNA ligase in a total volume of 1.5ml at 15° for 24 hr. The ligation reaction mixture was transformed into competent *E. coli* MC1061/P3 and a total of 200 x 10⁶ independent cDNA clones were obtained.

10 Plasmid DNA was prepared from a 500 ml culture of the original transformation of the cDNA library. Plasmid DNA was purified by the alkaline lysis procedure followed by twice banding in CsCl equilibrium gradients (Maniatis et al, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, NY (1987)).

15 B. Cloning Procedure

In the first round of screening, thirty 100 mm dishes of 50% confluent COS cells were transfected with 0.05µg/ml activated M12 murine B cell library DNA using the DEAE-Dextran method (Seed et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)). The cells were trypsinized and re-plated after 24 hours. After 47 hours, the cells were detached by
20 incubation in PBS/0.5 mM EDTA, pH 7.4/0.02% Na azide at 37°C for 30 min. The detached cells were treated with 10 µg/ml/human CTLA4Ig and murine CD28Ig for 45 minutes at 4°C. Cells were washed and distributed into panning dishes coated with affinity-purified Goat anti-human IgG antibody and allowed to attach at room temperature. After 3 hours, the plates were gently washed twice with PBS/0.5mM EDTA, pH 7.4/0.02% Na azide, 5% FCS and
25 once with 0.15M NaCl, 0.01 M Hepes, pH 7.4, 5% FCS. Episomal DNA was recovered from the panned cells and transformed into *E. coli* DH10B/P3. The plasmid DNA was re-introduced into COS cells via spheroplast fusion as described (Seed et al, *Proc. Natl. Acad. Sci. USA*, 84:3365 (1987)) and the cycle of expression and panning was repeated twice. In the second and third rounds of selection, after 47 hours, the detached COS cells were first
30 incubated with α-murine B7-1 mAb (16-10A1, 10 µg/ml), and COS cells expressing B7-1 were removed by α-mouse IgG and IgM coated magnetic beads. COS cells were then treated with 10µg/ml of human CTLA4Ig and murine CD28Ig and murine B7-2 expressing COS cells were selected by panning on dishes coated with goat anti-human IgG antibody. After the third round, plasmid DNA was prepared from individual colonies and transfected into
35 COS cells by the DEAE-Dextran method. Expression of B7-2 on transfected COS cells was analyzed by indirect immunofluorescence with CTLA4Ig.

After the final round of selection, plasmid DNA was prepared from individual colonies. A total of 6 of 8 candidate clones contained a cDNA insert of approximately 1.2 kb. Plasmid DNA from these eight clones was transfected into COS cells. All six clones with the 1.2 Kb cDNA insert were strongly positive for B7-2 expression by indirect immunofluorescence using CTLA4Ig and flow cytometric analysis.

C. Sequencing

The B7-2 cDNA insert in clone4 was sequenced in the pCDM8 expression vector employing the following strategy. Initial sequencing was performed using sequencing primers T7, CDM8R (Invitrogen) homologous to pCDM8 vector sequences adjacent to the cloned B7-2 cDNA (see Table II). Sequencing was performed using dye terminator chemistry and an ABI automated DNA sequencer. (ABI, Foster City, CA). DNA sequence obtained using these primers was used to design additional sequencing primers (see Table II). This cycle of sequencing and selection of additional primers was continued until the murine B7-2 cDNA was completely sequenced on both strands.

TABLE II

T7(F) (SEQ ID NO:3)	5'd[TAATACGACTCACTATAGGG]3'
CDM8(R) (SEQ ID NO:4)	5'd[TAAGGTTCTTCACAAAG]3'
MBX4-1F (SEQ ID NO:24)	5'd[ACATAAGCCTGAGTGAGCTGG]3'
MBX4-2R (SEQ ID NO:25)	5'd[ATGATGAGCAGCATCACAAGG]3'
MBX4-14 (SEQ ID NO:26)	5'd[TGGTCGAGTGAGTCCGAATAC]3'
MBX4-2F (SEQ ID NO:27)	5'd[GACGAGTAGTAACATACAGTG]3'

A murine B7-2 clone (mB7-2, clone 4) was obtained containing an insert of 1,163 base pairs with a single long open reading frame of 927 nucleotides and approximately 126 nucleotides of 3' noncoding sequences (Figure 14, SEQ ID NO:22). The predicted amino acid sequence encoded by the open reading frame of the protein is shown below the nucleotide sequence in Figure 14. The encoded murine B7-2 protein, is predicted to be 309 amino acid residues in length (SEQ ID NO:23). This protein sequence exhibits many features common to other type I Ig superfamily membrane proteins. Protein translation is predicted to begin at the methionine codon (ATG, nucleotides 111 to 113) based on the DNA homology in this region with the consensus eucaryotic translation initiation site (see Kozak, M. (1987) *Nucl. Acids Res.* 15:8125-8148). The amino terminus of the murine B7-2 protein (amino acids 1 to 23) has the characteristics of a secretory signal peptide with a predicted cleavage between the alanine at position 23 and the valine at position 24 (von Heijne (1987)

Nucl. Acids Res. 14:4683). Processing at this site would result in a murine B7-2 membrane bound protein of 286 amino acids having an unmodified molecular weight of approximately 32 kDa. This protein would consist of an approximate extracellular Ig superfamily V and C like domains of from about amino acid residue 24 to 246, a hydrophobic transmembrane domain of from about amino acid residue 247 to 265, and a long cytoplasmic domain of from about amino acid residue 266 to 309. The homologies to the Ig superfamily are due to the two contiguous Ig-like domains in the extracellular region bound by the cysteines at positions 40 to 110 and 157 to 216. The extracellular domain also contains nine potential N-linked glycosylation sites and, like murine B7-1, is probably glycosylated. Glycosylation of the murine B7-2 protein may increase the molecular weight to about 50-70 kDa. The cytoplasmic domain of murine B7-2 contains a common region which has a cysteine followed by positively charged amino acids which presumably functions as signaling or regulatory domain within an APC. Comparison of both the nucleotide and amino acid sequences of murine B7-2 with the GenBank and EMBL databases yielded significant homology (about 26% amino acid sequence identity) with human and murine B7-1. Murine B7-2 exhibits about 50% identity and 67% similarity with its human homologue, hB7-2. *E. coli* (DH106/p3) transfected with a vector (plasmid pmBx4) containing a cDNA insert encoding murine B7-2 (clone 4) was deposited with the American Type Culture Collection (ATCC) on August 18, 1993 as Accession No. 69388.

D. Costimulation

CD4⁺ murine T cells were purified by first depleting red blood cells by treatment with Tris-NH₄Cl. T cells were enriched by passage over a nylon wool column. CD4⁺ T cells were purified by two-fold treatment with a mixture of anti-MHC class II and anti-CD28 mAbs and rabbit complement. Murine B7-1 (obtained from Dr. Gordon Freeman, Dana-Farber Cancer Institute, Boston, MA; see also, Freeman, G.J. et al (1991) *J. Exp. Med.* 174, 625-631) murine B7-2, and vector transfected COS cells were harvested 72 hours after transfection, incubated with 25µg/ml mitomycin-C for one hour, and then extensively washed. 10⁵ murine CD4⁺ T cells were incubated with 1ng/ml of phorbol myristic acid (PMA) and 2 x 10⁴ COS transfectants (Table III). T cell proliferation was measured by ³H-thymidine (1µCi) incorporated for the last 12 hours of a 72 hour incubation.

TABLE III

3H-Thymidine Incorporation (cpm)

	CD4 ⁺ T cells	175
5	CD4 ⁺ T cells + 1ng/ml PMA	49
	CD4 ⁺ T cells + COS-vector	1750
	CD4 ⁺ T cells + COS-B7-1	4400
	CD4 ⁺ T cells + COS-B7-2	2236
	CD4 ⁺ T cells + 1ng/ml PMA + COS-vector	2354
10	CD4 ⁺ T cells + 1ng/ml PMA + COS-B7-1	67935
	CD4 ⁺ T cells + 1ng/ml PMA + COS-B7-2	43847

EXAMPLE 7

15 Construction and Characterization of Human B7-2 Immunoglobulin Fusion ProteinsA. Preparation Of Human B7-2Ig Fusion Proteins

The extracellular portion of human B7-2 was prepared as a fusion protein coupled to an immunoglobulin constant region. The immunoglobulin constant region may
 20 contain genetic modifications including those which reduce or eliminate effector activity inherent in the immunoglobulin structure. Briefly, DNA encoding the extracellular portion of hB7-2 was joined to DNA encoding the hinge, CH2 and CH3 regions of human IgC γ 1 or IgC γ 4 modified by directed mutagenesis. This was accomplished as described in the following subsections.

25

B. Preparation of Gene Fusions

DNA fragments corresponding to the DNA sequences of interest were prepared by polymerase chain reaction (PCR) using primer pairs described below. In general, PCR reactions were prepared in 100 μ l final volume composed of *Taq*, polymerase buffer
 30 (Gene Amp PCR Kit, Perkin-Elmer/Cetus, Norwalk, CT) containing primers (1 μ M each), dNTPs (200 μ M each) 1 ng of template DNA, and *Taq*, polymerase (Saiki, R.K., et al. (1988) *Science* 239:487-491). PCR DNA amplifications were run on a thermocycler (Ericomp, San Diego, CA) for 25 to 30 cycles each composed of a denaturation step (1 minute at 94°C), a renaturation step (30 seconds at 54°C), and a chain elongation step (1 minute at 72°C). The
 35 structure of each hB7-2 Ig genetic fusion consisted of a signal sequence to facilitate secretion coupled to the extracellular domain of B7-2 and the hinge, CH2 and CH3 domains of human IgC γ 1 or IgC γ 4. The IgC gamma 1 and IgC gamma 4 sequences contained nucleotide

changes within the hinge region to replace cysteine residues available for disulfide bond formation with serine residues and may contain nucleotide changes to replace amino acids within the CH2 domain thought to be required for IgC binding to Fc receptors and complement activation.

- 5 Sequence analysis confirmed structures of both $\mu\gamma_4$ and γ_1 clones, and each construct was used to transfect 293 cells to test transient expression. hIgG ELISA measured/confirmed transient expression levels approximately equal to 100 ng protein/ml cell supernatant for both constructs. NSO cell lines were transfected for permanent expression the the fusion proteins.

10

C. Genetic Construction of hB7-2Ig Fusion Proteins

(1). Preparation of Signal Sequence

- PCR amplification was used to generate an immunoglobulin signal sequence suitable for secretion of the B7-2Ig fusion protein from mammalian cells. The Ig signal
15 sequence was prepared from a plasmid containing the murine IgG heavy chain gene (Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA.* 86:38333837) using the oligonucleotide 5'-GGCACTAGGTCTCCAGCTTGAGATCACAGTTCTCTCTAC-3' (#01) (SEQ ID NO: 32) as the forward primer and the oligonucleotide 5'-GCTTGAATCTTCAGAGGAGCGGAGTGGACACCTGTGG-3' (#02) (SEQ ID NO: 33) as
20 the reverse PCR primer. The forward PCR primer (SEQ ID NO: 32) contains recognition sequences for restriction enzymes BsaI and is homologous to sequences 5' to the initiating methionine of the Ig signal sequence. The reverse PCR primer (SEQ ID NO: 33) is composed of sequences derived from the 5' end of the extracellular domain of hB7-2 and the
25 3' end of the Ig signal sequence. PCR amplification of the murine Ig signal template DNA using these primers resulted in a 224 bp product which is composed of BsaI restriction sites followed by the sequence of the Ig signal region fused to the first 20 nucleotides of the coding sequence of the extracellular domain of hB7-2. The junction between the signal sequence and hB7-2 is such that protein translation beginning at the signal sequence will continue into and through hB7-2 in the correct reading frame.

30

(2). Preparation of the hB7-2 Gene Segment

- The extracellular domain of the hB7.2 gene was prepared by PCR amplification of plasmid containing the hB7-2 cDNA inserted into expression vector pCDNAI (Freeman et al., *Science* 262:909-11 (1994)):
35 The extracellular domain of hB7-2 was prepared by PCR amplification using oligonucleotide 5'-GCTCCTCTGAAGATTCAAGC-3' (#03) (SEQ ID NO: 34) as the forward primer and oligonucleotide 5'-GGCACTATGATCAGGGGGAGGCTGAGGTCC-3'

(#04) (SEQ ID NO: 35) as the reverse primer. The forward PCR primer contained sequences corresponding to the first 20 nucleotides of the B7-2 extracellular domain and the reverse PCR primer contained sequences corresponding to the last 22 nucleotides of the B7-2 extracellular domain followed by a Bcl I restriction site and 7 noncoding nucleotides. PCR amplification with primer #03 (SEQ ID NO: 34) and #04 (SEQ ID NO: 35) yields a 673 bp product corresponding to the extracellular IgV and IgC like domains of hB7-2 followed by a unique Bcl I restriction site.

The signal sequence was attached to the extracellular portion of hB7-2 by PCR as follows. DNA-PCR products obtained above corresponding to the signal sequence and the hB7-2 extracellular domain were mixed in equimolar amounts, denatured by heating to 100°C, held at 54°C for 30°C to allow the complementary ends to anneal and the strands were filled in using dNTPs and Taq polymerase. PCR primers #01 (SEQ ID NO: 32) and #04 (SEQ ID NO: 35) were added and the entire fragment produced by PCR amplification to yield a ~880 fragment composed of a BsaI restriction site followed by the signal sequence fused to the extracellular domain of hB7-2, followed by a Bcl I restriction site.

(3). Cloning and Modification of Immunoglobulin Fusion Domain

Plasmid pSP721gG1 was prepared by cloning the 2000 bp segment of human IgG1 heavy chain genomic DNA (Ellison, J.W., et al. (1982) *Nucl. Acids. Res.* 10:4071-4079) into the multiple cloning site of cloning vector pSP72 (Promega, Madison, WI). Plasmid pSP721gG1 contained genomic DNA encoding the CH1, hinge, CH2 and CH3 domains of the heavy chain human IgG1 gene. PCR primers designed to amplify the hinge-CH2-CH3 portion of the heavy chain along with the intervening DNA were prepared as follows. The forward PCR primer 5'-GCATTTTAAGCTTTTCCTGATCAGGAGCCCAAATCTTCT GACAAAACCTCACACATCTCCACCGTCTCCAGGTAAGCC-3' (SEQ ID NO: 36) contained HindIII and Bcl I restriction sites and was homologous to the hinge domain sequence except for five nucleotide substitutions which would change the three cysteine residues to serines. The reverse PCR primer 5'TAATACGACTCACTATAGGG-3' (SEQ ID NO: 37) was identical to the commercially available T7 primer (Promega, Madison, WI). Amplification with these primers yielded a 1050 bp fragment bounded on the 5' end by HindIII and BclI restriction sites and on the 3' end by BamHI, SmaI, KpnI, SacI, EcoRI, ClaI, EcoRV and BglII restriction sites. This fragment contained the IgC hinge domain in which the three cysteine codons had been replaced by serine codons followed by an intron, the CH2 domain, an intron, the CH3 domain and additional 3' sequences. After PCR amplification, the DNA fragment was digested with HindIII and EcoRI and cloned into expression vector pNRDSH digested with the same restriction enzymes. This created plasmid pNRDSH/IgG1.

A similar PCR based strategy was used to clone the hinge-CH2-CH3 domains of human IgCgamma4 constant regions. A plasmid, p428D (Medical Research Council, London, England) containing the complete IgCgamma4 heavy chain genomic sequence (Ellison, J. Buxbaum, J. and Hood, L.E. (1981) *DNA* 1: 11 -18) was used as a template for

5 PCR amplification using oligonucleotide 5'GAGCATTTCCTGATCAGGA GTCCAAATATGGTCCCCCATCCCATCATCCCCAGGTAAGCCAACCC-3' (SEQ ID NO: 38) as the forward PCR primer and oligonucleotide

10 5'GCAGAGGAATCGAGCTCGGTACCCGGGGATCCCCAGTGTGGGGACAGTGGGA CCGCTCTGCCTCCC-3' (SEQ ID NO: 39) as the reverse PCR primer. The forward PCR primer (SEQ ID NO: 38) contains a BclI restriction site followed by the coding sequence for the hinge domain of IgCgamma4. Nucleotide substitutions have been made in the hinge region to replace the cysteines residues with serines. The reverse PCR primer (SEQ ID NO: 39) contains a PspAI restriction site (5'CCCGGG-3'). PCR amplification with these primers results in a 1179 bp DNA fragment. The PCR product was digested with BclI and PspAI and

15 ligated to pNRDSH/IgG1 digested with the same restriction enzymes to yield plasmid pNRDSH/IgG4. In this reaction, the IgCγ 4 domain replaced the IgCγ1 domain present in pNRDSH/IgG1.

Modification of the CH2 domain in IgC to replace amino acids thought to be involved in binding to Fc receptor was accomplished as follows. Plasmid pNRDSH/IgG1

20 served as template for modifications of the IgCγ1 CH2 domain and plasmid pNRDSH/IgG4 served as template for modifications of the IgCγ 4 CH2 domain. Plasmid pNRDSH/IgG1 was PCR amplified using a forward PCR primer (SEQ ID NO: 36) and oligonucleotide 5'-GGGTTTT GGGGGGAAGAGGAAGACTGACGGTGCCCCC

25 TCGGCTTCAGGTGCTGAGGAAG-3' (SEQ ID NO: 40) as the reverse PCR primer. The forward PCR primer (SEQ ID NO: 36) has been previously described and the reverse PCR primer (SEQ ID NO: 40) was homologous to

the amino terminal portion of the CH2 domain of IgG1 except for five nucleotide substitutions designed to change amino acids 234, 235, and 237 (Canfield, S. M. and Morrison, S. L. (1991) *J. Exp. Med.* 173: 1483-1491.) from Leu to Ala, Leu to Glu, and Gly

30 to Ala, respectively. Amplification with these PCR primers will yield a 239 bp DNA fragment consisting of a modified hinge domain, an intron and modified portion of the CH2 domain. Plasmid pNRDSH/IgG1 was also PCR amplified with the oligonucleotide 5'-CATCTCTTCCTCAGCACCTGAAGCCGAGGGGGCACCCTCAGTCTTCCTCTTCCC

35 CC-3' (SEQ ID NO: 41) as the forward primer and oligonucleotide (SEQ ID NO: 37) as the reverse PCR primer. The forward PCR primer (SEQ ID NO: 41) is complementary to primer (SEQ ID NO: 40) and contains the five complementary nucleotide changes necessary for the CH2 amino acid replacements. The reverse PCR primer (SEQ ID NO: 37) has been

previously described. Amplification with these primes yields a 875 bp fragment consisting of the modified portion of the CH2 domain, an intron, the CH3 domain, and 3' additional sequences. The complete IgC γ 1 segment consisting of modified hinge domain, modified CH2 domain and CH3 domain was prepared by an additional PCR reaction. The purified products of the two PCR reactions above were mixed, denatured (95°C, 1 minute) and then renatured (54°C, 30 seconds) to allow complementary ends of the two fragments to anneal. The strands were filled in using dNTP and Taq polymerase and the entire fragment amplified using forward PCR primer (SEQ ID NO: 36) and reverse PCR primer (SEQ ID NO: 37). The resulting fragment of 1050 bp was purified, digested with HindIII and EcoRI and ligated to pNRDSH previously digested with the same restriction enzymes to yield plasmid pNRDSHlgG1 m.

Two amino acids at immunoglobulin positions 235 and 237 were changed from Leu to Glu and Gly to Ala, respectively, within the IgC γ 4 CH2 domain to eliminate Fc receptor binding. Plasmid pNRDSH/IgG4 was PCR amplified using the forward primer (SEQ ID NO: 38) and the oligonucleotide 5'-CGCACGTGACCTCAGGGGTCCGGGAGATCATGAGAGTGTCTTGGGTTTGGGGGGAACAGGAAGACTGATGGTGCCCCCTCGAACTCAGGTGCTGAGG-3' (SEQ ID NO: 42) as the reverse primer. The forward primer has been previously described and the reverse primer was homologous to the amino terminal portion of the CH2 domain, except for three nucleotide substitutions designed to replace the amino acids described above. This primer also contained a PmlI restriction site for subsequent cloning. Amplification with these primers yields a 265 bp fragment composed of the modified hinge region, and intron, and the modified 5' portion of the CH2 domain.

Plasmid pNRDSH/IgG4 was also PCR amplified with the oligonucleotide 5'-CCTCAGCACCTGAGTTCGAGGGGGCACCATCAGTCTCCTGTTCCCCAAAACCCAAGGACACTCTCATGATCTCCCGGACCCCTGAGGTACAGTGCG-3' (SEQ ID NO: 43) as the forward primer and oligonucleotide (SEQ ID NO: 39) as the reverse PCR primer. The forward PCR primer (SEQ ID NO: 43) is complementary to primer (SEQ ID NO: 42) and contains the three complementary nucleotide changes necessary for the CH2 amino acid replacements. The reverse PCR primer (SEQ ID NO: 39) has been previously described. Amplification with these primers yields a 1012 bp fragment consisting of the modified portion of the CH2 domain, an intron, the CH3 domain, and 3' additional sequences. The complete IgC γ 4 segment consisting of modified hinge domain, modified CH2 domain and CH3 domain was prepared by an additional PCR reaction. The purified products of the two PCR reactions above were mixed, denatured (95°C, 1 minute) and then renatured (54°C, 30 seconds) to allow complementary ends of the two fragments to anneal. The strands were filled in using dNTP and Taq polymerase and the entire fragment amplified using forward

PCR primer (SEQ ID NO: 38) and reverse PCR primer (SEQ ID NO: 39). The resulting fragment of 1179 bp was purified, digested with BclI and PspAI and ligated to pNRDSH previously digested with the same restriction enzymes to yield plasmid pNRDSH/IgG4m.

5 (4). Assembly of Final hB7-2Ig Genes

The PCR fragment corresponding to the Ig signal-hB7-2 gene fusion prepared above was digested with BsaI and BclI restriction enzymes and ligated to pNRDSH/IgG1, pNRDSH/IgG1m, pNRDSH/IgG4, and pNRDSH/IgG4m previously digested with Hind III and BclI. The ligated plasmids were transformed into *E. coli* JM109 using CaC12 competent
10 cells and transformants were selected on L-agar containing ampicillin (50 µg/ml; Molecular Cloning: A Laboratory Manual (1982) Eds. Maniatis, T., Fritsch, E. E., and Sambrook, J. Cold Spring Harbor Laboratory). Plasmids isolated from the transformed *E. coli* were analyzed by restriction enzyme digestion. Plasmids with the expected restriction plasmid were sequenced to verify all portions of the signal-hB7-2-IgG gene fusion segments.

15

D. Expression Cloning of hB7-2V-IgG1 and hB7-2C IgG1

The variable and constant domains of human B7-2 were separately cloned into pNRDSH/IgG1. These clonings were accomplished using PCR. The portions of hB7-2
20 corresponding to the variable and constant regions were determined from intron/exon mapping and previously published gene structure analysis.

Human B7-2 Variable Domain

5'GCTCCTCTGAAGATT.....GAACTGTCAGTGCTT3' (SEQ ID
NO: 44)
25 A P L K I E L S V L (SEQ ID
NO: 45)

Human B7-2 Constant Domain

5'GCTAACTTCAGTCAA.....CCTTTCTCTATAGAG3' (SEQ ID
30 NO: 46)
 A N F S Q P F S I E (SEQ ID
NO: 47)

(1). Assembly of hB7-2VIg

35 The hB7-2V domain Ig sequence was assembled using a PCR strategy similar to that shown above. The signal sequence was derived from the onco M gene by PCR amplification of a plasmid containing the onco M gene using oligonucleotide 5'-

GCAACCGGAAGCTTGCCACCATGGGGGTACTGCTCACACAGAGGACG-3' (#05) (SEQ ID NO: 48) as the forward PCR primer and 5'-AGTCTCATTGAAATAAGCTTGAATCTTCAGAGGAGCCATGCTGGCCATGCTTGGA AACAGGAG-3' (#06) (SEQ ID NO: 49) as the reverse primer. The forward PCR primer
 5 (#05) (SEQ ID NO: 48) contains a Hind III restriction site and the amino terminal portion of the onco M signal sequence. The reverse PCR (#06) (SEQ ID NO: 49) contains the sequence corresponding to the 3' portion of the onco M signal sequence fused to the 5' end of the hB7-2 IgV like domain.

The hB7-2 IgV like domain was obtained by PCR amplification of the hB7-2
 10 cDNA using oligonucleotide 5'-CTCCTGTTTCCAAGCATGGCCAGCATGGCTCCTCTGAA GATTCAGGCTTATTTCAATGAGAC-3' (#07) (SEQ ID NO: 50) as the forward and oligonucleotide 5'-TGTGTGTGGAATTCTCATTACTGATCAAGCACTGACAGTTCAGAATTCATC-3'
 15 (#08) (SEQ ID NO : 51) as the reverse PCR primer. PCR amplification with these primers yields the hB7-2 IgV domain with a portion of the 3' end of the onco M signal sequence on the 5' end and a Bcl I restriction site on the 3' end. The signal and IgV domain were linked together in a PCR reaction in which equimolar amounts of the onco M signal and IgV domain DNA fragments were mixed, denatured, annealed, and the strands filled in. Subsequent PCR
 20 amplification using forward primer #05 (SEQ ID NO: 48) and reverse primer #08 (SEQ ID NO: 51) yielded a DNA fragment containing a Hind III restriction site, followed by the onco M signal fused to the B7-2 IgV domain followed by a Bcl I restriction site. This PCR fragment was digested with Hind II and Bcl I and cloned into expression vector pNRDSH/IgG1 digested with the same restriction enzymes to yield pNRDSH/B7-2CIg.

25

(2). Assembly of hB7-2CIg

The expression plasmid for hB7-2IgC domain was prepared as described above for the IgV domain except for using PCR primers specific for the IgC domain. The onco M signal sequence was prepared using oligonucleotide #05 (SEQ ID NO: 48) as the
 30 forward PCR primer and oligonucleotide 5'-AGAAATTGGTACTATTTAGGTTGACTGAAGTTAGCCATGCTGGCCATGCTTGGA AACAGGAG-3' (#09) (SEQ ID NO: 52) as the reverse PCR primer. The hB7-2 IgC domain was prepared using oligonucleotide 5'-CTCCTGTTTCCAAGCATGGCCAGCATGGCTAACTTCAGTC
 35 AACCTGAAATAGTACCAATTTC-3' (#11) (SEQ ID NO: 53) as the reverse PCR primer. The two PCR products were mixed and amplified with primers #05 (SEQ ID NO: 48) and #11 (SEQ ID NO: 53) to assemble the onco M signal sequence with the hB7-2IgC domain.

The PCR product was subsequently digested with Hind III and BclI and ligated to pNRDSH/IgG1 digested with similar restriction enzymes to yield the final expression plasmid pNRDSH/hB7-2CIgG1.

5 E. Competitive Binding Assays with Human B7-2Ig Fusion Proteins

To determine the affinity of binding of different forms of soluble B7-1 and B7-2 proteins to CTLA4, competitive binding assays were performed with these proteins. The soluble B7-2VIg, B7-2CIg, B7-2Ig, and B7-1Ig fusion proteins used in these assays were expressed and purified as follows.

- 10 The preparation of expression vectors encoding human B7-2VIg, B7-2CIg, and B7-2Ig fusion proteins is described above. The expression vector encoding the B7-1Ig fusion protein, containing an OncoM leader sequence linked the extracellular domain of B7-1 was prepared similarly, using the PCR primers OncoMB71F (5'CTCAAGCTTGCCACCATGGGGGTACTGCTCACACAGAGGACG
15 CTGCTCAGTCTGGTCCTTGCACTCCTGTTTCCGAGCATGGCGAGCATGGGTCTTTC TCACTTC3'; SEQ ID NO: 54) and B71/BclI (5'TGTGTGTGGAATTCTCA TTACTGATCAGGAAAATGCTCTTGCTTG3'; SEQ ID NO: 55). The plasmid pKShB7-1, containing the OncoM leader sequence linked to the human B7-1 cDNA sequence (the nucleotide sequence of the human B7-1 cDNA is disclosed in Freeman, G.J. et al., (1989) *J. Immunol.* 143:2714-2722) was used as a template in this PCR reaction.

The B7Ig fusion proteins were prepared by transfection of COS cells or Chinese Hamster Ovary (CHO) cells and purification of the protein from the supernatant of the cultures.

- 25 Cell culture reagents were obtained from Gibco-BRL, Gaithersburg, MD. CHO cells were maintained in alpha MEM supplemented with 10% Fetal bovine serum (FBS) and glutamine. Penicillin, streptomycin, and fungizone were typically added. COS cells were maintained in DMEM with 10% FBS and supplemented as described for CHO cells. All cells were kept at 5% CO₂ at 37°C in a humidified incubator.

- 30 All fusion constructs except the hB7-1Ig construct were expressed transiently in COS cells. Typical transient transfections were done using 200 µg/ml of DEAE-dextran, 100 µM chloroquine and 5 µg of DNA per 10 cm dish in serum-free DMEM. The cells were treated until vacuoles were noted and the cells appeared distressed (about 3 hours). Cells were shocked with 10%DMSO/PBS for 2 minutes, then incubated with DMEM/10%FBS overnight. The following morning the media was changed to DMEM/serum free and left
35 until harvest at 72 hours post transfection. The hB7-1Ig construct was transfected into CHO cells by calcium phosphate transfection. The line was made stable using Geneticin (G418)

resistance selection, and expression was amplified using methotrexate and alpha MEM lacking nucleosides.

For all transiently transfected constructs (i.e., all constructs except hB7-1Ig) media enriched for Ig fusion proteins produced by the transiently transfected host cells was harvested 72 hours post-transfection. The amount of Ig fusion proteins produced by the host cells was measured by performing an anti-human IgG Elisa assay with the supernatant of the cultures. For this assay, Maxisorp plates (Nunc, Denmark) were coated overnight with 20 µg/ml of goat anti-human IgG (H+L) (Zymed, San Francisco, CA) in PBS. The plates were blocked for 1 hour with PBS/0.1%BSA and then incubated for an additional hour with cell culture supernatants from the transfected cells. After 5 washes with PBS/0.05%Tween, HRP-coupled goat anti-human IgG(H+L) (Zymed) was added as a 1:1000 dilution in PBS. After a 1 hour incubation the plates were washed again, then enzymatically developed using an ABTS kit (Zymed) as described above.

Expression levels were approximately 3 µg/ml for the constructs. The B7Ig fusion proteins were purified from the supernatant of transfected host cells by protein A purification as follows. Protein A Sepharose IPA300 (Repligen, Cambridge, MA) was washed in OBB (1.5 M glycine, 3 M NaCl, pH 8.9), resuspended in DMEM, and added to cell culture supernatant at 1 ml/liter of supernatant. The solution was left to mix gently overnight at 4°C. The Protein A sepharose was then pelleted, the majority of the supernatant was removed, and the remaining supernatant was used to resuspend the Protein-A Sepharose prior to loading onto a Poly-prep column (BioRad, Hercules, CA). The column was washed extensively with OBB and the immunoglobulin fusion protein was eluted in 0.1 M sodium citrate. Half volume column washes were collected into 1/10th volume 1 M Tris (pH9.0). Protein containing fractions were identified by a standard colorimetric reaction (BioRad), pooled, and dialyzed overnight against PBS in 6000-8000 KD dialysis tubing. The purified proteins were of the expected size and high purity, representing >90% of total protein stained with Commassie Blue on acrylamide gels.

The ability of various B7 family-Ig fusion proteins to competitively inhibit the binding of biotinylated-CTLA4Ig to immobilized B7-2Ig was determined. Competition binding assays were done as follows and analysed according to McPherson (McPherson, G.A. (1985) *J. Pharmacol. Methods* 14:213-228). Soluble hCTLA4Ig was labelled with ¹²⁵I to a specific activity of approximately 2 x 10⁶ cpm/pmol. hB7-2-Ig fusion protein was coated overnight onto microtiter plates at 10mg/ml in 10 mM Tris-HCl, pH8.0, 50 µl /well. The wells were blocked with binding buffer (DMEM containing 10% heat-inactivated FBS, 0.1% BSA, and 50 mM BES, pH 6.8) for 2 h at room temperature. The labeled CTLA4-Ig (4nM) was added to each well in the presence or absence of unlabeled competing Ig fusion

proteins, including full-length B7-2 (hB7-2Ig), full-length B7-1 (hB7-1Ig), the variable region-like domain of B7-2 (hB7-2VIg) and the constant region-like domain of B7-2 (hB7-2CIg) and allowed to bind for 2.5 h at room temperature. The wells were washed once with ice-cold binding buffer and then four times with ice-cold PBS. Bound radioactivity was recovered by treatment of the wells with 0.5 N NaOH for 5 min and the solubilized material removed and counted in a gamma counter.

The results of these assays are shown in Figure 15 in which both hB7-2Ig (10-20 nM) and hB7-2VIg (30-40 nM) competitively inhibit the binding of CTLA4Ig to immobilized B7-2 protein. hB7-2CIg is unable to compete with soluble CTLA4, indicating that the B7-2 binding region is in found in the variable-region like domain.

F. Competitive Binding Assays for B7-1 and B7-2 Fusion Proteins

The ability of recombinant CTLA4Ig to bind to hB7-1 or hB7-2 was assessed in a competitive binding ELISA assay as follows. Purified recombinant CTLA4Ig (20 µg/ml in PBS) was bound to a Costar EIA/RIA 96 well microtiter dish (Costar Corp, Cambridge MA, USA) in 50 µL overnight at room temperature. The wells were washed three times with 200 µL of PBS and the unbound sites blocked by the addition of 1 % BSA in PBS (200 µl/well) for 1 hour at room temperature. The wells were washed as above. Biotinylated B7-1Ig or B7-2Ig (1 µg/ml serially diluted in twofold steps to 15.6 ng/mL; 50 µL) was added to each well and incubated for 2.5 hours at room temperature. The wells were washed as above. The bound biotinylated B7-Ig was detected by the addition of 50 µl/well of a 1:2000 dilution of streptavidin-HRP (Pierce Chemical Co., Rockford, IL) for 30 minutes at room temperature. The wells were washed as above and 50 µL of ABTS (Zymed, California) added and the developing blue color monitored at 405 nm after 30 min. The ability of unlabelled B7-1Ig or B7-2Ig to compete with biotinylated B7-1Ig or B7-2Ig, respectively, was assessed by mixing varying amounts of the competing protein with a quantity of biotinylated B7-1Ig or B7-2Ig shown to be non-saturating (i.e., 70 ng/mL; 1.5nM) and performing the binding assays as described above. A reduction in the signal (Abs 405 nm) expected for biotinylated B7-1Ig or B7-2Ig indicated a competition for binding to immobilized CTLA4Ig.

Considering the previous evidence that CTLA4 was the high affinity receptor for B7-1, the avidity of binding of CTLA4 and CD28 to B7-1 and B7-2 was compared in assays as described above. In a first experiment, B7-1Ig or B7-2-Ig was labelled with biotin and bound to immobilized CTLA4-Ig in the presence or absence of increasing concentrations of unlabeled B7-1Ig or B7-2Ig. The experiment was repeated with ¹²⁵I-labeled B7-1Ig or B7-2Ig. Representative results are shown in Figure 16 (Panel A: B7-1Ig; Panel B: B7-2Ig). Using this solid phase binding assay, the avidity of B7-2 (2.7 nM) for CTLA4 was

determined to be approximately two-fold lower than that observed for B7-1 (4.6 nM). The experimentally determined IC₅₀ values are indicated in the upper right corner of the panels. The affinity of both B7-1 and B7-2 for CD28 was lower and was difficult to confidently determine.

5

G. Direct Binding Assays of modified forms of B7 family members to CTLA4Ig

Direct binding ELISA assays were performed to determine the level of binding of B7 family members as Ig fusion proteins to CTLA4. For these assays, the immunoglobulin fusion proteins were attached to plates and the amount of biotinylated CTLA4 binding to the plates was determined as described below.

Nunc Maxisorp plates were coated overnight at room temperature with 50 μ l per well of a 20 μ g/ml stock of the various B7Ig fusion proteins, or purified human IgG (Zymed) in PBS as described above. Human CTLA4Ig (Repligen) was biotinylated using NHS-LC-biotin (Pierce, Rockford, IL). Varying amounts of biotinylated CTLA4Ig were added to the plates and incubated for 2 hours at room temperature. The plates were washed five times with PBS and then a 1:1000 dilution of streptavidin-HRP (Zymed) was added and left for 30' minutes on the plates. After another series of washes with PBS, the HRP reactivity was measured using an ABTS kit (Zymed) as described above.

The results, presented in Figure 20, show that half-saturation occurred at 500pM for B7-1Ig, and at 5 nM and 8 nM for B7-2VIg and B7-2Ig, respectively. Thus, CTLA4Ig binds to a similar extent to the B7-2VIg and B7-2Ig fusion proteins. CTLA4 does not, however bind to B7-2CIg. Thus, the variable domain of B7-2 is sufficient for binding to CTLA4.

H. Binding of B7-2VIg, B7-2Ig, and B7-1Ig to CHO-CTLA4 cells

The examples presented in section F and G of Example 7 showed that B7-2VIg and B7-2Ig bind soluble CTLA4Ig. The present example shows that B7-2VIg and B7-2Ig also bind to CTLA4 expressed on a cell.

For this example, labeled B7-1Ig and B7-2Ig fusion proteins were incubated with CHO cells transfected to express CTLA4 and binding was measured by flow cytometry as follows.

B7-1 and B7-2 immunoglobulin fusion proteins, prepared as described above, were diluted to 20 μ g/ml in PBS/1%BSA and incubated with 10⁶ CHO cells transfected to express CTLA4 on their surface (CHO/CTLA4 cells) for 30 minutes on ice. The cells were washed twice with cold PBS/BSA and incubated with a 1:50 dilution of goat anti-human IgG-FITC (Zymed) for 30 minutes on ice. The cells were washed once with cold PBS/BSA, once with cold PBS and then resuspended in 250 μ l cold PBS. The cells were then fixed by

adding 250 μ l of a 2% paraformaldehyde solution in PBS and incubation for at least 1 hour and the fluorescence analyzed using a FACS (Becton Dickinson, San Jose, CA). Similarly treated CHO/CTLA4 cells which recieved the secondary antibody alone served to measure background staining.

- 5 The results of the flow cytometric analysis are presented in Figure 21. The results show that hB7-2Ig and hB7-2VIg fusion proteins bind to a similar extent to CTLA4 positive cells (Figure 21, panels C and D, repectively) and that the binding is stronger than binding of hB7-1Ig to CHO/CTLA4 cells (panel E).

10 I. B7-2VIg binds with stronger affinity to CD28 than B7-1Ig and B7-2Ig

 The Example shown in the previous section showed that B7-2VIg and B7-2Ig fusion proteins bind with similar affinity to cell membrane bound CTLA4. This example shows that the fusion proteins bind with different affinities to CD28 and in particular, that B7-2VIg binds with higher affinity to CD28 than B7-2Ig.

- 15 B7-1 and B7-2 immunoglobulin fusion proteins diluted at 20 μ g/ml in PBS/1%BSA were incubated with 10^6 CHO cells transfected to express CD28 on their surface (CHO/CD28 cells) for 30 minutes on ice. The cells were washed twice with cold PBS/BSA and incubated with a 1:50 dilution of goat anti-human IgG-FITC (Zymed) for 30 minutes on ice. The cells were washed once with cold PBS/BSA, once with cold PBS and
20 then resuspended in 250 μ l cold PBS. The cells were then fixed by adding 250 μ l of a 2% paraformaldehyde solution in PBS and incubation for at least 1 hour and the fluorescence analyzed using a FACS (Becton Dickinson, San Jose, CA).

- Representative results, as presented in Figure 22, indicate that B7-2Ig and B7-2VIg fusion proteins bind specifically to CHO-CD28 cells. The results further indicate that
25 B7-2VIg protein binds to CD28 with stronger affinity than does B7-2Ig and B7-1Ig.

 Thus, B7-2VIg fusion protein binds with higher affinity than B7-2Ig to CD28, whereas both fusion proteins bind with similar affinity to CTLA4.

30 J. B7-2VIg is more potent than B7-2Ig and B7-1Ig at costimulating proliferation of CD28+ T cells

 Since B7-2VIg binds with higher affinity to CD28 than B7-2Ig, it was next investigated whether B7-2VIg fusion protein is also more potent at stimulating T cell proliferation than B7-2Ig fusion protein.

- CD28+ T cells were isolated from peripheral blood leukocytes (PBLs) as
35 described above. For measuring T cell proliferation, 1.2×10^5 CD28+ T cells were incubated in 200 μ l of culture media in 96 well plates and stimulated with PMA at 1 ng/ml and either of the following costimulatory signals: 6×10^4 CHO/B7-1 or CHO/B7-2 cells (pretreated

overnight with mitomycin C and then extensively washed), 30 or 100 µg/ml of B7-1Ig, B7-2Ig, or B7-2VIg. Alternatively, the fusion proteins can first be incubated with a 3 fold excess (w/w) of affinity purified goat anti-human IgGFc (Cappel) for 30 minutes prior to use. After 60 hours of incubation, the T cells were pulsed overnight with ³H-thymidine (Dupont/NEN) and harvested for counting, as described above.

The results of the proliferation assay are represented graphically in Figure 23. The results indicate that CHO expressed B7-1 and B7-2 strongly induced T cell proliferation. Purified B7-1Ig and B7-2Ig also induced proliferation, although not as potently as the CHO/B7-1 and CHO/B7-2 cells. However, B7-2VIg induced proliferation to the same extent as the CHO/B7-1 and CHO/B7-2 cells. Thus, B7-2VIg is as potent as CHO/B7-1 and CHO/B7-2 cells in costimulating proliferation of T cells.

In another example, CD28+ T cells were activated with anti-CD3 coated plates prepared as described above and costimulated with various amounts of B7-1Ig, B7-2Ig, and B7-2VIg. Proliferation of the CD28+ T cells was measured after 60 hours and overnight pulsing of the cells with ³H-thymidine. The results, presented graphically in Figure 24 indicate that B7-2VIg fusion protein is also more potent than B7-2Ig and B7-1Ig at costimulating CD28+ T cells when anti-CD3 is used as the primary activating agent. Moreover, the results indicate that B7-2VIg is more potent than B7-2Ig and B7-1Ig at costimulating proliferation of CD28+ T cells when low concentrations of the proteins are used. This is apparent when comparing the amount of thymidine incorporated in cells incubated with 1 µg/ml of costimulatory fusion protein. In addition, B7-2VIg costimulates proliferation of the T cells at doses as low as 10 ng/ml (250 pM).

Thus, the higher binding affinity of B7-2VIg versus B7-2Ig fusion protein for CD28 (Example 7, section I) correlates with a higher costimulatory activity of B7-2VIg versus B7-2Ig for proliferation of the T cells.

K. B7-2VIg is more potent than B7-1Ig and B7-2Ig at costimulating production of IL-2 by CD28+ T cells

It was next investigated whether B7-2VIg was also more potent than B7-1Ig and B7-2Ig at costimulating activated T cells for the production of IL-2.

In the second example described in the previous section (Section J of Example 7), in which CD28+ T cells were activated with anti-CD3 and costimulated with various amounts of B7-1Ig, B7-2Ig, and B7-2VIg, and cell proliferation measured, the level of IL-2 in the supernatant was determined after 18 hours of stimulation using an ELISA kit (Endogen, Cambridge, MA). The results, presented in Figure 24 show that more IL-2 is produced by T cells costimulated with B7-2VIg than by T cells costimulated with B7-1Ig or

B7-2Ig. This was most apparent at low concentrations of costimulatory proteins (for example at 1 µg/ml of the fusion proteins).

Another example was performed to compare production of IL-2 from CD28+ T cells costimulated with CHO/B7-2 cells or costimulated with B7-2VIg protein. CD28+ T cells were incubated in anti-CD3 coated plates in the presence of CHO/B7-2 cells or B7-2VIg protein for 1, 2, or 3 days, and the amount of IL-2 was measured in the supernatant. The results, presented in Figure 25, show that B7-2VIg is more potent at costimulating T cells for the production of IL-2 than CHO/B7-2 cells.

In a further example, the amount of IL-2 produced by CD28+ T cells costimulated with anti-CD28, B7-1Ig, B7-2Ig, or B7-2VIg was compared after 1, 2, or 5 days of costimulation. CD28+ T cells were activated and costimulated with anti-CD28 antibody, or with B7-1Ig, B7-2Ig, or B7-2VIg fusion protein. The amount of IL-2 in the supernatant was measured after 1, 2, and 5 days of costimulation. Figure 26, representing graphically the amount of IL-2 produced by the T cells, indicate that B7-2VIg is more potent than B7-2Ig and B7-1Ig at stimulating production of IL-2 by CD28+ T cells and further, that after 5 days of culture, only T cells costimulated with B7-2Ig or B7-2VIg fusion proteins were producing IL-2. Moreover, T cells costimulated with B7-2VIg produce more IL-2 than T cells costimulated with B7-2Ig after 5 days of culture.

Thus, B7-2VIg costimulate T cells to produce high levels of IL-2, even after at least 5 days of culture.

L. B7-2VIg is a potent costimulator of activated CD4+ T cells for the production of cytokines.

In this example, the amount of IL-2, IL-4, IFN-γ, and GM-CSF produced by T cells costimulated with B7-2VIg, B7-2Ig, or B7-1Ig was compared.

CD4+CD28+ T cells were cultured at 2×10^6 cells/ml in T75 flasks coated with anti-CD3 antibody alone, or with anti-CD28 mAb 9.3, or one of the fusion proteins B7-1Ig, B7-2Ig, or B7-2VIg. After 18 hours of culture, the amount of IL-2, IL-4, interferon-γ (IFN-γ), and granulocyte macrophage-colony stimulating factor (GM-CSF) was measured by ELISA using commercially available kits (IL-2 (BioSource, Camarillo, CA), IL-4 (Endogen, Cambridge, MA), IFN-γ (Bio-Source, Camarillo, CA), and GM-CSF (R&D Systems, Minneapolis, MN)). The amount of IL-2 and IL-4 was also measured in cell costimulated for 120 hours.

The results are presented in Table VI. The results indicate that costimulation with B7-2VIg leads to production of high levels of IL-2, IL-4, IFN-γ, and GM-CSF. Moreover, the amount of all 4 cytokines produced from T cells costimulated with B7-2VIg was higher than the amount of cytokines produced from T cells costimulated with B7-1Ig.

Compared to B7-2Ig, B7-2VIg also costimulated the production of higher amounts of IL-2, IL-4 and GM-CSF and similar amounts of IFN- γ . Thus, B7-2VIg is a potent costimulator for production of cytokines by T cells.

Moreover, after 120 hours of culture, the T cells costimulated with B7-2VIg produced more than twice the amount of IL-4 produced by T cells costimulated with B7-2Ig and approximately 8 fold the amount of IL-4 produced by T cells costimulated with B7-1Ig. Thus, T cells costimulated with B7-2VIg fusion protein induces production of high levels of IL-4 and this production is longlasting. Costimulation of T cells with B7-2VIg could thus drive T cells to a T helper 2 (Th2) state in long term culture.

10

Table VI

Cytokine production by CD4+CD28+T cells costimulated with B7-2VIg, B7-2Ig, B7-1Ig or anti-CD28 antibody

costimulus	IL-2 18hr	IL-2 120hr	IL-4 18hr	IL-4 120hr	IFN-g 18hr	GM-CSF 18hr
none	19	0	6.5	0	27.6	0
anti-CD28	2412	N.D.	83.2	0	28	25
B7-1Ig	734	0	10.4	5.7	27.8	10
B7-2Ig	1557	104	10.8	18.6	42.6	12
B7-2vIg	3073	262	29.8	40.8	37.5	30

15

M. B7-2Ig and B7-2VIg, but not B7-1Ig promote sustained T cell growth.

In this example, the capability of B7Ig fusion proteins to promote growth of T cells for extended periods was analysed.

CD28+ T lymphocytes were incubated in the presence of anti-CD3 plus anti-CD28, or B7-1Ig, B7-2Ig, or B7-2VIg immobilized on beads and the total cell numbers monitored over a period of 12 days. The results are presented in Figure 27. Cells stimulated with anti-CD3 alone fail to proliferate, and die. Cells stimulated with anti-CD3 plus B7-2Ig go through one cycle of replication and then apoptose. Cells stimulated with anti-CD3 plus anti-CD28 or B7-2Ig or B7-2vIg continue to replicate. Thus, B7-2Ig and B7-2VIg fusion proteins, but not B7-1Ig, are capable of stimulating prolonged growth of CD28+ T cells.

20

EXAMPLE 8**Production and Characterization of Monoclonal Antibodies to Human B7-2****A. Immunizations and Cell Fusions**

- 5 Balb/c female mice (obtained from Taconic Labs, Germantown, NY) were immunized intraperitoneally with 50 µg human B7.2-Ig emulsified in complete Freund's adjuvant (Sigma Chemical Co., St. Louis, MO) or 10⁶ CHO-human B7.2 cells per mouse. The mice were given two booster immunizations with 10-25 µg human B7.2-Ig emulsified in incomplete Freund's adjuvant (Sigma Chemical Co., St. Louis, MO) or CHO-human B7.2
- 10 cells at fourteen day intervals following the initial immunization for the next two months. The mice were bled by retro-orbital bleed and the sera assayed for the presence of antibodies reactive to the immunogen by ELISA against human B7.2-Ig. ELISA against hCTLA4-Ig was also used to control for Ig tail directed antibody responses. Mice showing a strong serological response were boosted intravenously via the tail vein with 25 µg human hB7.2-Ig
- 15 diluted in phosphate-buffered saline (PBS), pH 7.2 (GIBCO, Grand Island, NY). Three to four days following this boost, the spleens from these mice were fused 5:1 with SP 2/0 myeloma cells (American Type Culture Collection, Rockville, MD, No. CRL8006), which are incapable of secreting both heavy and light immunoglobulin chains (Kearney et al. (1979) *J. Immunol.* 123:1548). Standard methods based upon those developed by Kohler and
- 20 Milstein (*Nature* (1975) 256:495) were used.

B. Antibody Screening

- After 10-21 days, supernatants from wells containing hybridoma colonies from the fusion were screened for the presence of antibodies reactive to human B7.2 as
- 25 follows: Each well of a 96 well flat bottomed plate (Costar Corp., Cat. #3590) was coated with 50 µl per well of a 1 µg/ml human B7.2-Ig solution or 5 x 10⁴ 3T3-hB7.2 cells on lysine coated plates in phosphate-buffered saline, pH 7.2, overnight at 4 °C. The human B7.2-Ig solution was aspirated off, or the cells were cross-linked to the plates with glutaraldehyde, and the wells were washed three times with PBS, then blocked with 1% BSA solution (in
- 30 PBS) (100µl/well) for one hour at room temperature. Following this blocking incubation, the wells were washed three times with PBS and 50 µl of hybridoma supernatant was added per well and incubated for 1.5 hours at room temperature. Following this incubation, the wells were washed three times with PBS and then incubated for 1.5 hours at room temperature with
- 35 50 µl per well of a 1:4000 dilution of horseradish peroxidase-conjugated, affinity purified, goat anti-mouse IgG or IgM heavy and light chain-specific antibodies (HRP; Zymed Laboratories, San Francisco, CA). The wells were then washed three times with PBS, followed by a 30 minute incubation in 50 µl per well of 1 mM 2,2-azino-bis-3-

ethylbenzthiazoline-6-sulfonic acid (ABTS) in 0.1 M Na-Citrate, pH 4.2 to which a 1:1000 dilution of 30 % hydrogen peroxide had been added as a substrate for HRP to detect bound antibody. The absorbance was then determined at OD₄₁₀ on a spectrophotometric autoreader (Dynatech, Virginia).

- 5 Three hybridomas, HA3.1F9, HA5.2B7 and HF2.3D1, were identified that produced antibodies to human B7.2-Ig. HA3.1F9 was determined to be of the IgG1 isotype, HA5.2B7 was determined to be of the IgG2b isotype and HF2.3D1 as determined to be of the IgG2a isotype. Each of these hybridomas were subcloned two additional times to insure that they were monoclonal. Hybridoma cells were deposited with the American Type Culture
10 Collection, which meets the requirements of the Budapest Treaty, on July 19, 1994 as ATCC Accession No. HB 11688 (hybridoma HA3.1F9), ATCC Accession No. HB 11687 (HA5.2B7) and ATCC Accession No. HB 11686 (HF2.3D1).

C. Competitive ELISA

- 15 Supernatants from the hybridomas HA3.1F9, HA5.2B7 and HF2.3D1 were further characterized by competitive ELISA, in which the ability of the monoclonal antibodies to inhibit the binding of biotinylated hCTLA4Ig to immobilized hB7-2 immunoglobulin fusion proteins was examined. Biotinylation of hCTLA4Ig was performed using Pierce Immunopure NHS-LC Biotin (Cat. No. 21335). B7-2 immunoglobulin fusion
20 proteins used were: hB7.2-Ig (full-length hB7-2), hB7.2-VIg (hB7-2 variable domain only) and hB7.2-CIg (B7-2 constant domain only). A hB7.1-Ig fusion protein was used as a control. For the ELISA, 96 well plates were coated with the Ig fusion protein (50 µl/well of a 20 µg/ml solution) overnight at room temperature. The wells were washed three times with PBS, blocked with 10 % fetal bovine serum (FBS), 0.1 % bovine serum albumin (BSA) in
25 PBS for 1 hour at room temperature, and washed again three times with PBS. To each well was added 50 µl of Bio-hCTLA4-Ig (70 ng/ml) and 50 µl of competitor monoclonal antibody supernatant. Control antibodies were an anti-B7.1 mAb (EW3.5D12) and the anti-hB7-2 mAb B70 (IgG2bk, obtained from Pharmingen). The wells were washed again and streptavidin-conjugated horse radish peroxidase (from Pierce, Cat. No. 21126; 1:2000
30 dilution, 50 µl/well) was added and incubated for 30 minutes at room temperature. The wells were washed again, followed by a 30 minute incubation in 50 µl per well of ABTS in 0.1 M Na-Citrate, pH 4.2 to which a 1:1000 dilution of 30 % hydrogen peroxide had been added as a substrate for HRP to detect bound antibody. The absorbance was then determined at OD₄₁₀ on a spectrophotometric autoreader (Dynatech, Virginia). The results, shown in
35 Table IV below, demonstrate that each of the mAbs produced by the hybridomas HA3.1F9, HA5.2B7 and HF2.3D1 are able to competitively inhibit the binding of hCTLA4Ig to full-length hB7.2-Ig or hB7.2-VIg (hCTLA4Ig does not bind to hB7.2CIg).

TABLE IV

	<u>Blocking of Binding</u>			
	<u>hB7.1-Ig</u>	<u>hB7.2-Ig</u>	<u>hB7.2-VIg</u>	<u>hB7.2-CIg</u>
EW3.5D12 (anti-hB7.1 mAb)	Yes	No	No	No
B70 (anti-hB7-2)	No	Yes	Yes	No
HA3.1F9 (anti-hB7-2)	No	Yes	Yes	No
HA5.2B7 (anti-hB7-2)	No	Yes	Yes	No
HF2.3D1 (anti-hB7-2)	No	Yes	Yes	No

5

D. Flow Cytometry

Supernatants from the hybridomas HA3.1F9, HA5.2B7 and HF2.3D1 were also characterized by flow cytometry. Supernatants collected from the clones were screened by flow cytometry on CHO and 3T3 cells transfected to express hB7.2 (CHO-hB7.2 and 3T3-hB7.2, respectively) or control transfected 3T3 cells (3T3-Neo). Flow cytometry was performed as follows: 1×10^6 cells were washed three times in 1 % BSA in PBS, then the cells were incubated in 50 μ l hybridoma supernatant or culture media per 1×10^6 cells for 30 minutes at 4 °C. Following the incubation, the cells were washed three times with 1 % BSA in PBS, then incubated in 50 μ l fluorescein-conjugated goat anti-mouse IgG or IgM antibodies (Zymed Laboratories, San Francisco, CA) at 1:50 dilution per 1×10^6 cells for 30 minutes at 4 °C. The cells were then washed three times in 1 % BSA in PBS and fixed with 1 % paraformaldehyde solution. The cell samples were then analyzed on a FACScan flow cytometer (Becton Dickinson, San Jose CA). The results, shown in Figures 17, 18 and 19, demonstrate the monoclonal antibodies produced by the hybridomas HA3.1F9, HA5.2B7 and HF2.3D1 each bind to hB7-2 on the surface of cells.

E. Inhibition of Proliferation of Human T Cells by Anti-hB7-2 mAbs

Hybridoma supernatants containing anti-human B7-2 mAbs were tested for their ability to inhibit hB7-2 costimulation of human T cells. In this assay, purified CD28⁺ human T cells were treated with submitogenic amounts of PMA (1ng/ml) to deliver the primary signal and with CHO cells expressing hB7-2 on their surface to deliver the costimulatory signal. Proliferation of the T cells was measured after three days in culture by the addition of ³H-thymidine for the remaining 18 hours. As shown in Table V, resting T cells show little proliferation as measured by ³H-thymidine incorporation (510 pm). Delivery of signal 1 by PMA results in some proliferation (3800 pm) and T cells receiving both the primary (PMA) and costimulatory (CHO/hB7-2) signals proliferate maximally (9020 cpm).

All three anti-hB7-2 mAbs tested reduce the costimulatory signal induced proliferation to that found for PMA treated cells alone showing that these mAbs can inhibit T cell proliferation by blocking the B7/CD28 costimulatory pathway.

5 TABLE V

Addition to CD28 ⁺ T Cells	hB7-2 mAb	CPM
---	---	510
+PMA	---	3800
+PMA + CHO/hB7-2	---	9020
+PMA + CHO/hB7-2	HF2.301	3030
---	HA5.2B7	1460
---	HA3.1F9	2980

F. Antibodies to the B7-2 Variable domain block B7-2 function

10 In this example, the ability of a series of monoclonal antibodies to B7-2 to bind to the Ig-variable or Ig-constant domains of B7-2, and to inhibit T cell proliferation was analyzed.

Monoclonal antibodies to human B7-1 and B7-2 were prepared from Balb/c mice using SP2/0 cells and standard protocols. Briefly, Balb/c female mice (Taconic Labs, Germantown, NY) were immunized intraperitoneally with either 50 µgs B7-2Ig emulsified in CFA (Sigma, St. Louis, MO) or 10⁶ CHO/B7-2 cells. The mice were boosted twice at 14 day intervals following the initial immunization and once with B7-2Ig protein in PBS. Hybridoma colonies were established in 96 well tissue culture plates and the culture supernatants were assayed for direct binding to B7-2Ig. All mAbs were purified from ascites fluid on Protein-A sepharose as described above. MAb B70 was purchased from PharMingen (San Diego, CA).

20 Purified mAbs were tested for their ability to bind to the various B7-Ig forms as follows. Maxisorp plates (Nunc) were coated overnight at room temperature with 20 µg/ml of purified B7-2Ig protein in PBS. The plates were then blocked with PBS/0.1%BSA for 1 h. Purified antibody (5 µgs/ml in PBS) was added to the test wells, the plates incubated for 1 hour, and then washed 5 times with PBS/0.05% Tween20. Goat anti-mouse IgG-HRP (Zymed) was added and allowed to react for 1 hour, followed by 5 washes. The plates were developed as described above.

The binding characteristics of the antibodies is indicated in Table VII under the heading "Recognition". As indicated in Table VII, all antibodies recognized B7-2Ig. Binding of the antibodies to B7-2Vlg and B7-2CIg fusion proteins indicated that the antibodies recognize either the variable region construct or the constant region construct, but not both constructs.

The antibodies were further analyzed for their ability to inhibit binding of B7-2Ig to CTLA4 and to CD28 and to inhibit T cell proliferation. Using a competition ELISA format, varying amounts of mAbs were added to wells coated B7-2Ig and containing 35 ngs/ml biotinylated CTLA4Ig or CD28Ig. The ability of mAbs to disrupt the binding was measured as a decrease in the specific signal of captured biotinylated CTLA4Ig or CD28Ig. The capability of the antibodies to inhibit proliferation of T cells was determined by performing proliferation assays, as described above in which one of the antibodies was added.

The results are presented in Table VII under the heading "Inhibition". In general, antibodies to the V-domain of B7-2 inhibit binding of the B7-2Ig to CD28 and CTLA4 and also inhibit CHO/B7-2 driven T cell proliferation. Antibodies to the C-domain are not inhibitory suggesting that the functionality of B7-2 resides in the V-domain.

Table VII**Characterization of anti-B7-2 antibodies**

Anti- B7-2 mAbs		Recognition			Inhibition		
		B7	B7v domain	B7c domain	CTLA4 binding	CD28 binding	T cell prolif.
HA5.1F9	IgG1	+	-	-	+	+	+
HA5.2B7	IgG2b	+	+	-	+	+	+
HF2.3D11	IgG2a	+	+	-	+	+	+
HF4.3C11	IgG1	+	+	-	+	+	+
HF4.3E8	IgG1	+	+	-	+	+/-	+/-
HF4.5B4	IgG1	+	-	+	nd	-	-
HF4.5H12	IgM	+	-	+	-	-	-
HF4.6B1	IgG2a	+	-	+	-	-	-
HF4.6H4	IgG1	+	+	-	+	+	+
B70	IgG2b	+	+	-	+	+	+

20

Moreover, all of the anti-B7-1 and anti-B7-2 mAbs tested also recognized their respective ligand when expressed on the surface of CHO cells and on the surface of

activated human B cells where tested. None of the anti-B7-1 or anti-B7-2 mAbs showed any crossreactivity with the other B7 protein.

EQUIVALENTS

- 5 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

- (A) NAME: DANA-FARBER CANCER INSTITUTE
(B) STREET: 44 BINNEY STREET
(C) CITY: BOSTON
10 (D) STATE: MASSACHUSETTS
(E) COUNTRY: USA
(F) POSTAL CODE (ZIP): 02115
(G) TELEPHONE: (617) 632-4016
(H) TELEFAX: (617) 632-4012
15
(A) NAME: REPLIGEN CORPORATION
(B) STREET: ONE KENDALL SQUARE, BLDG 700
(C) CITY: CAMBRIDGE
20 (D) STATE: MASSACHUSETTS
(E) COUNTRY: USA
(F) POSTAL CODE (ZIP): 02139
(G) TELEPHONE: (617) 225-6000
(H) TELEFAX: (617) 494-1975

25

(ii) TITLE OF INVENTION: Novel CTLA4/CD28 Ligands and
Uses Therefor

(iii) NUMBER OF SEQUENCES: 55

30

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: LAHIVE & COCKFIELD
(B) STREET: 60 State Street, Suite 510
(C) CITY: Boston
(D) STATE: Massachusetts
35 (E) COUNTRY: USA
(F) ZIP: 02109

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
40 (B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
45 (B) FILING DATE:

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 08/479,744
50 (B) FILING DATE: 7-JUN-1995

(viii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 08/280,757
55 (B) FILING DATE: 26-JUL-1994

(ix) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/101,624
(B) FILING DATE: 26-JUL-1993

(x) PRIOR APPLICATION DATA:

5 (A) APPLICATION NUMBER: 08/109,393
(B) FILING DATE: 19-AUG-1993

(xi) PRIOR APPLICATION DATA:

10 (A) APPLICATION NUMBER: 08/147,773
(B) FILING DATE: 03-NOV-1993

(xii) ATTORNEY/AGENT INFORMATION:

15 (A) NAME: Mandragouras, Amy E.
(B) REGISTRATION NUMBER: 36,207
(C) REFERENCE/DOCKET NUMBER: RPI-004C3PC

(xiii) TELECOMMUNICATION INFORMATION:

20 (A) TELEPHONE: (617) 227-7400
(B) TELEFAX: (617) 227-5941

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 1120 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

35 (A) NAME/KEY: CDS
(B) LOCATION: 107..1093

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

40	CACAGGGTGA AAGCTTTGCT TCTCTGCTGC TGTAACAGGG ACTAGCACAG ACACACGGAT	60
	GAGTGGGGTC ATTTCCAGAT ATTAGGTCAC AGCAGAAGCA GCCAAA ATG GAT CCC	115
		Met Asp Pro
		1
45	CAG TGC ACT ATG GGA CTG AGT AAC ATT CTC TTT GTG ATG GCC TTC CTG	163
	Gln Cys Thr Met Gly Leu Ser Asn Ile Leu Phe Val Met Ala Phe Leu	
	5 10 15	
50	CTC TCT GGT GCT GCT CCT CTG AAG ATT CAA GCT TAT TTC AAT GAG ACT	211
	Leu Ser Gly Ala Ala Pro Leu Lys Ile Gln Ala Tyr Phe Asn Glu Thr	
	20 25 30 35	
	GCA GAC CTG CCA TGC CAA TTT GCA AAC TCT CAA AAC CAA AGC CTG AGT	259
55	Ala Asp Leu Pro Cys Gln Phe Ala Asn Ser Gln Asn Gln Ser Leu Ser	
	40 45 50	

	GAG CTA GTA GTA TTT TGG CAG GAC CAG GAA AAC TTG GTT CTG AAT GAG	307
	Glu Leu Val Val Phe Trp Gln Asp Gln Glu Asn Leu Val Leu Asn Glu	
	55 60 65	
5	GTA TAC TTA GGC AAA GAG AAA TTT GAC AGT GTT CAT TCC AAG TAT ATG	355
	Val Tyr Leu Gly Lys Glu Lys Phe Asp Ser Val His Ser Lys Tyr Met	
	70 75 80	
10	GGC CGC ACA AGT TTT GAT TCG GAC AGT TGG ACC CTG AGA CTT CAC AAT	403
	Gly Arg Thr Ser Phe Asp Ser Asp Ser Trp Thr Leu Arg Leu His Asn	
	85 90 95	
15	CTT CAG ATC AAG GAC AAG GGC TTG TAT CAA TGT ATC ATC CAT CAC AAA	451
	Leu Gln Ile Lys Asp Lys Gly Leu Tyr Gln Cys Ile Ile His His Lys	
	100 105 110 115	
20	AAG CCC ACA GGA ATG ATT CGC ATC CAC CAG ATG AAT TCT GAA CTG TCA	499
	Lys Pro Thr Gly Met Ile Arg Ile His Gln Met Asn Ser Glu Leu Ser	
	120 125 130	
25	GTG CTT GCT AAC TTC AGT CAA CCT GAA ATA GTA CCA ATT TCT AAT ATA	547
	Val Leu Ala Asn Phe Ser Gln Pro Glu Ile Val Pro Ile Ser Asn Ile	
	135 140 145	
30	ACA GAA AAT GTG TAC ATA AAT TTG ACC TGC TCA TCT ATA CAC GGT TAC	595
	Thr Glu Asn Val Tyr Ile Asn Leu Thr Cys Ser Ser Ile His Gly Tyr	
	150 155 160	
35	CCA GAA CCT AAG AAG ATG AGT GTT TTG CTA AGA ACC AAG AAT TCA ACT	643
	Pro Glu Pro Lys Lys Met Ser Val Leu Leu Arg Thr Lys Asn Ser Thr	
	165 170 175	
40	ATC GAG TAT GAT GGT ATT ATG CAG AAA TCT CAA GAT AAT GTC ACA GAA	691
	Ile Glu Tyr Asp Gly Ile Met Gln Lys Ser Gln Asp Asn Val Thr Glu	
	180 185 190 195	
45	CTG TAC GAC GTT TCC ATC AGC TTG TCT GTT TCA TTC CCT GAT GTT ACG	739
	Leu Tyr Asp Val Ser Ile Ser Leu Ser Val Ser Phe Pro Asp Val Thr	
	200 205 210	
50	AGC AAT ATG ACC ATC TTC TGT ATT CTG GAA ACT GAC AAG ACG CGG CTT	787
	Ser Asn Met Thr Ile Phe Cys Ile Leu Glu Thr Asp Lys Thr Arg Leu	
	215 220 225	
55	TTA TCT TCA CCT TTC TCT ATA GAG CTT GAG GAC CCT CAG CCT CCC CCA	835
	Leu Ser Ser Pro Phe Ser Ile Glu Leu Glu Asp Pro Gln Pro Pro Pro	
	230 235 240	
60	GAC CAC ATT CCT TGG ATT ACA GCT GTA CTT CCA ACA GTT ATT ATA TGT	883
	Asp His Ile Pro Trp Ile Thr Ala Val Leu Pro Thr Val Ile Ile Cys	
	245 250 255	
65	GTG ATG GTT TTC TGT CTA ATT CTA TGG AAA TGG AAG AAG AAG AAG CGG	931
	Val Met Val Phe Cys Leu Ile Leu Trp Lys Trp Lys Lys Lys Lys Arg	

	260		265		270		275	
	CCT CGC AAC TCT TAT AAA TGT GGA ACC AAC ACA ATG GAG AGG GAA GAG							979
	Pro Arg Asn Ser Tyr Lys Cys Gly Thr Asn Thr Met Glu Arg Glu Glu							
5		280		285		290		
	AGT GAA CAG ACC AAG AAA AGA GAA AAA ATC CAT ATA CCT GAA AGA TCT							1027
	Ser Glu Gln Thr Lys Lys Arg Glu Lys Ile His Ile Pro Glu Arg Ser							
		295		300		305		
10								
	GAT GAA GCC CAG CGT GTT TTT AAA AGT TCG AAG ACA TCT TCA TGC GAC							1075
	Asp Glu Ala Gln Arg Val Phe Lys Ser Ser Lys Thr Ser Ser Cys Asp							
		310		315		320		
15								
	AAA AGT GAT ACA TGT TTT TAATTAAAGA GTAAAGCCCA AAAAAAA							1120
	Lys Ser Asp Thr Cys Phe							
		325						

20 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 329 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30	Met Asp Pro Gln Cys Thr Met Gly Leu Ser Asn Ile Leu Phe Val Met	
	1 5 10 15	
35	Ala Phe Leu Leu Ser Gly Ala Ala Pro Leu Lys Ile Gln Ala Tyr Phe	
	20 25 30	
	Asn Glu Thr Ala Asp Leu Pro Cys Gln Phe Ala Asn Ser Gln Asn Gln	
	35 40 45	
40	Ser Leu Ser Glu Leu Val Val Phe Trp Gln Asp Gln Glu Asn Leu Val	
	50 55 60	
	Leu Asn Glu Val Tyr Leu Gly Lys Glu Lys Phe Asp Ser Val His Ser	
	65 70 75 80	
45	Lys Tyr Met Gly Arg Thr Ser Phe Asp Ser Asp Ser Trp Thr Leu Arg	
	85 90 95	
	Leu His Asn Leu Gln Ile Lys Asp Lys Gly Leu Tyr Gln Cys Ile Ile	
	100 105 110	
50	His His Lys Lys Pro Thr Gly Met Ile Arg Ile His Gln Met Asn Ser	
	115 120 125	
55	Glu Leu Ser Val Leu Ala Asn Phe Ser Gln Pro Glu Ile Val Pro Ile	
	130 135 140	

Ser Asn Ile Thr Glu Asn Val Tyr Ile Asn Leu Thr Cys Ser Ser Ile
 145 150 155 160
 5 His Gly Tyr Pro Glu Pro Lys Lys Met Ser Val Leu Leu Arg Thr Lys
 165 170 175
 Asn Ser Thr Ile Glu Tyr Asp Gly Ile Met Gln Lys Ser Gln Asp Asn
 180 185 190
 10 Val Thr Glu Leu Tyr Asp Val Ser Ile Ser Leu Ser Val Ser Phe Pro
 195 200 205
 Asp Val Thr Ser Asn Met Thr Ile Phe Cys Ile Leu Glu Thr Asp Lys
 15 210 215 220
 Thr Arg Leu Leu Ser Ser Pro Phe Ser Ile Glu Leu Glu Asp Pro Gln
 225 230 235 240
 20 Pro Pro Pro Asp His Ile Pro Trp Ile Thr Ala Val Leu Pro Thr Val
 245 250 255
 Ile Ile Cys Val Met Val Phe Cys Leu Ile Leu Trp Lys Trp Lys Lys
 260 265 270
 25 Lys Lys Arg Pro Arg Asn Ser Tyr Lys Cys Gly Thr Asn Thr Met Glu
 275 280 285
 Arg Glu Glu Ser Glu Gln Thr Lys Lys Arg Glu Lys Ile His Ile Pro
 30 290 295 300
 Glu Arg Ser Asp Glu Ala Gln Arg Val Phe Lys Ser Ser Lys Thr Ser
 305 310 315 320
 35 Ser Cys Asp Lys Ser Asp Thr Cys Phe
 325

(2) INFORMATION FOR SEQ ID NO:3:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 45 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

50 TAATACGACT CACTATAGGG

20

(2) INFORMATION FOR SEQ ID NO:4:

- 55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

10 TAAGGTTTCCT TCACAAAG

18

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

25

ACTGGTAGGT ATGGAAGATC C

21

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

40

ATGCGAATCA TTCCTGTGGG C

21

(2) INFORMATION FOR SEQ ID NO:7:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AAAGCCCACA GGAATGATTC G

21

(2) INFORMATION FOR SEQ ID NO:8:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

15

CTCTCAAAAC CAAAGCCTGA G

21

(2) INFORMATION FOR SEQ ID NO:9:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

30

TTAGGTCACA GCAGAAGCAG C

21

(2) INFORMATION FOR SEQ ID NO:10:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

45

TCTGGAAACT GACAAGACGC G

21

(2) INFORMATION FOR SEQ ID NO:11:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

5 CTCAGGCTTT GGTTTGTAGA G 21

(2) INFORMATION FOR SEQ ID NO:12:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

20 CACTCTCTTC CCTCTCCATT G 21

(2) INFORMATION FOR SEQ ID NO:13:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

35 GACAAGCTGA TGGAAACGTC G 21

(2) INFORMATION FOR SEQ ID NO:14:

40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

50 CAATGGAGAG GGAAGAGAGT G 21

(2) INFORMATION FOR SEQ ID NO:15:

55 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 5 (ii) MOLECULE TYPE: oligonucleotide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
- 10 CTTTAGAGCA CA 12
- (2) INFORMATION FOR SEQ ID NO:16:
- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 20 (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
- 25 CTCTAAAG 8
- (2) INFORMATION FOR SEQ ID NO:17:
- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
- 35 (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
- 40 Lys Tyr Met Gly Arg Thr Ser Phe Asp
5
- (2) INFORMATION FOR SEQ ID NO:18:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 13 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
- 50 (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
- 55

Lys Ser Gln Asp Asn Val Thr Glu Lys Tyr Asp Val Ser
5 10

5 (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Trp Lys Trp Lys Lys Lys Lys Arg Pro Arg Asn Ser Tyr Lys Cys
5 10 15

20

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
25 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

35 TGGCCCATGG CTTCAGA

17

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
40 (B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

50

GCCAAAATGG ATCCCA

17

(2) INFORMATION FOR SEQ ID NO:22:

55

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1163 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

10

(A) NAME/KEY: CDS
 (B) LOCATION: 111..1040

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

15

CCCACGCGTC CGGGAGCAAG CAGACGCGTA AGAGTGGCTC CTGTAGGCAG CACGGACTTG 60

AACAACCAGA CTCCTGTAGA CGTGTTCAG AACTTACGGA AGCACCCACG ATG GAC 116
 Met Asp

20

1

CCC AGA TGC ACC ATG GGC TTG GCA ATC CTT ATC TTT GTG ACA GTC TTG 164
 Pro Arg Cys Thr Met Gly Leu Ala Ile Leu Ile Phe Val Thr Val Leu
 5 10 15

25

CTG ATC TCA GAT GCT GTT TCC GTG GAG ACG CAA GCT TAT TTC AAT GGG 212
 Leu Ile Ser Asp Ala Val Ser Val Glu Thr Gln Ala Tyr Phe Asn Gly
 20 25 30

30

ACT GCA TAT CTG CCG TGC CCA TTT ACA AAG GCT CAA AAC ATA AGC CTG 260
 Thr Ala Tyr Leu Pro Cys Pro Phe Thr Lys Ala Gln Asn Ile Ser Leu
 35 40 45 50

35

AGT GAG CTG GTA GTA TTT TGG CAG GAC CAG CAA AAG TTG GTT CTG TAC 308
 Ser Glu Leu Val Val Phe Trp Gln Asp Gln Gln Lys Leu Val Leu Tyr
 55 60 65

40

GAG CAC TAT TTG GGC ACA GAG AAA CTT GAT AGT GTG AAT GCC AAG TAC 356
 Glu His Tyr Leu Gly Thr Glu Lys Leu Asp Ser Val Asn Ala Lys Tyr
 70 75 80

45

CTG GGC CGC ACG AGC TTT GAC AGG AAC AAC TGG ACT CTA CGA CTT CAC 404
 Leu Gly Arg Thr Ser Phe Asp Arg Asn Asn Trp Thr Leu Arg Leu His
 85 90 95

50

AAT GTT CAG ATC AAG GAC ATG GGC TCG TAT GAT TGT TTT ATA CAA AAA 452
 Asn Val Gln Ile Lys Asp Met Gly Ser Tyr Asp Cys Phe Ile Gln Lys
 100 105 110

AAG CCA CCC ACA GGA TCA ATT ATC CTC CAA CAG ACA TTA ACA GAA CTG 500
 Lys Pro Pro Thr Gly Ser Ile Ile Leu Gln Gln Thr Leu Thr Glu Leu
 115 120 125 130

55

	TCA GTG ATC GCC AAC TTC AGT GAA CCT GAA ATA AAA CTG GCT CAG AAT	548
	Ser Val Ile Ala Asn Phe Ser Glu Pro Glu Ile Lys Leu Ala Gln Asn	
	135 140 145	
5	GTA ACA GGA AAT TCT GGC ATA AAT TTG ACC TGC ACG TCT AAG CAA GGT	596
	Val Thr Gly Asn Ser Gly Ile Asn Leu Thr Cys Thr Ser Lys Gln Gly	
	150 155 160	
10	CAC CCG AAA CCT AAG AAG ATG TAT TTT CTG ATA ACT AAT TCA ACT AAT	644
	His Pro Lys Pro Lys Lys Met Tyr Phe Leu Ile Thr Asn Ser Thr Asn	
	165 170 175	
15	GAG TAT GGT GAT AAC ATG CAG ATA TCA CAA GAT AAT GTC ACA GAA CTG	692
	Glu Tyr Gly Asp Asn Met Gln Ile Ser Gln Asp Asn Val Thr Glu Leu	
	180 185 190	
20	TTC AGT ATC TCC AAC AGC CTC TCT CTT TCA TTC CCG GAT GGT GTG TGG	740
	Phe Ser Ile Ser Asn Ser Leu Ser Leu Ser Phe Pro Asp Gly Val Trp	
	195 200 205 210	
25	CAT ATG ACC GTT GTG TGT GTT CTG GAA ACG GAG TCA ATG AAG ATT TCC	788
	His Met Thr Val Val Cys Val Leu Glu Thr Glu Ser Met Lys Ile Ser	
	215 220 225	
30	TCC AAA CCT CTC AAT TTC ACT CAA GAG TTT CCA TCT CCT CAA ACG TAT	836
	Ser Lys Pro Leu Asn Phe Thr Gln Glu Phe Pro Ser Pro Gln Thr Tyr	
	230 235 240	
35	TGG AAG GAG ATT ACA GCT TCA GTT ACT GTG GCC CTC CTC CTT GTG ATG	884
	Trp Lys Glu Ile Thr Ala Ser Val Thr Val Ala Leu Leu Leu Val Met	
	245 250 255	
40	CTG CTC ATC ATT GTA TGT CAC AAG AAG CCG AAT CAG CCT AGC AGG CCC	932
	Leu Leu Ile Ile Val Cys His Lys Lys Pro Asn Gln Pro Ser Arg Pro	
	260 265 270	
45	AGC AAC ACA GCC TCT AAG TTA GAG CGG GAT AGT AAC GCT GAC AGA GAG	980
	Ser Asn Thr Ala Ser Lys Leu Glu Arg Asp Ser Asn Ala Asp Arg Glu	
	275 280 285 290	
50	ACT ATC AAC CTG AAG GAA CTT GAA CCC CAA ATT GCT TCA GCA AAA CCA	1028
	Thr Ile Asn Leu Lys Glu Leu Glu Pro Gln Ile Ala Ser Ala Lys Pro	
	295 300 305	
55	AAT GCA GAG TGAAGGCAGT GAGAGCCTGA GGAAAGAGTT AAAAATTGCT	1077
	Asn Ala Glu	
50	TTGCCTGAAA TAAGAAGTGC AGAGTTTCTC AGAATTCAAA AATGTTCTCA GCTGATTGGA	1137
	ATTCTACAGT TGAATAATTA AAGAAC	1163

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 309 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

```

10 Met Asp Pro Arg Cys Thr Met Gly Leu Ala Ile Leu Ile Phe Val Thr
    1           5           10           15

    Val Leu Leu Ile Ser Asp Ala Val Ser Val Glu Thr Gln Ala Tyr Phe
        20           25           30

15 Asn Gly Thr Ala Tyr Leu Pro Cys Pro Phe Thr Lys Ala Gln Asn Ile
    35           40           45

    Ser Leu Ser Glu Leu Val Val Phe Trp Gln Asp Gln Gln Lys Leu Val
    50           55           60

20 Leu Tyr Glu His Tyr Leu Gly Thr Glu Lys Leu Asp Ser Val Asn Ala
    65           70           75           80

    Lys Tyr Leu Gly Arg Thr Ser Phe Asp Arg Asn Asn Trp Thr Leu Arg
    85           90           95

    Leu His Asn Val Gln Ile Lys Asp Met Gly Ser Tyr Asp Cys Phe Ile
        100           105           110

30 Gln Lys Lys Pro Pro Thr Gly Ser Ile Ile Leu Gln Gln Thr Leu Thr
    115           120           125

    Glu Leu Ser Val Ile Ala Asn Phe Ser Glu Pro Glu Ile Lys Leu Ala
    130           135           140

35 Gln Asn Val Thr Gly Asn Ser Gly Ile Asn Leu Thr Cys Thr Ser Lys
    145           150           155           160

    Gln Gly His Pro Lys Pro Lys Lys Met Tyr Phe Leu Ile Thr Asn Ser
    165           170           175

    Thr Asn Glu Tyr Gly Asp Asn Met Gln Ile Ser Gln Asp Asn Val Thr
        180           185           190

45 Glu Leu Phe Ser Ile Ser Asn Ser Leu Ser Leu Ser Phe Pro Asp Gly
    195           200           205

    Val Trp His Met Thr Val Val Cys Val Leu Glu Thr Glu Ser Met Lys
    210           215           220

50 Ile Ser Ser Lys Pro Leu Asn Phe Thr Gln Glu Phe Pro Ser Pro Gln
    225           230           235           240

    Thr Tyr Trp Lys Glu Ile Thr Ala Ser Val Thr Val Ala Leu Leu Leu
    245           250           255

```

Val Met Leu Leu Ile Ile Val Cys His Lys Lys Pro Asn Gln Pro Ser
 260 265 270

Arg Pro Ser Asn Thr Ala Ser Lys Leu Glu Arg Asp Ser Asn Ala Asp
 5 275 280 285

Arg Glu Thr Ile Asn Leu Lys Glu Leu Glu Pro Gln Ile Ala Ser Ala
 290 295 300

10 Lys Pro Asn Ala Glu
 305

(2) INFORMATION FOR SEQ ID NO:24:

15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

ACATAAGCCT GAGTGAGCTG G

21

30

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- 35 (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

45 ATGATGAGCA GCATCACAAG G

21

(2) INFORMATION FOR SEQ ID NO:26:

50

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:
TGGTCGAGTG AGTCCGAATA C 21

10 (2) INFORMATION FOR SEQ ID NO:27:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: oligonucleotide

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:
GACGAGTAGT AACATACAGT G 21

25 (2) INFORMATION FOR SEQ ID NO:28:
(i) SEQUENCE CHARACTERISTICS:
30 (A) LENGTH: 1491 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear
35 (ii) MOLECULE TYPE: cDNA to mRNA
(iii) HYPOTHETICAL: no
40 (iv) ANTI-SENSE: no
(vi) ORIGINAL SOURCE:
45 (A) ORGANISM: Homo sapien
(F) TISSUE TYPE: lymphoid
(G) CELL TYPE: B cell
(H) CELL LINE: Raji
(vii) IMMEDIATE SOURCE:
50 (A) LIBRARY: cDNA in pCDM8 vector
(B) CLONE: B7, Raji clone #13
(viii) POSITION IN GENOME:
55 (A) CHROMOSOME/SEGMENT: 3

(ix) FEATURE:

- 5 (A) NAME/KEY: Open reading frame (translated region)
(B) LOCATION: 318 to 1181 bp
(C) IDENTIFICATION METHOD: similarity to other pattern

(ix) FEATURE:

- 10 (A) NAME/KEY: Alternate polyadenylation signal
(B) LOCATION: 1474 to 1479 bp
(C) IDENTIFICATION METHOD: similarity to other pattern

(x) PUBLICATION INFORMATION:

- 15 (A) AUTHORS: FREEMAN, GORDON J.
FREEDMAN, ARNOLD S.
SEGIL, JEFFREY M.
LEE, GRACE
20 WHITMAN, JAMES F.
NADLER, LEE M.

(B) TITLE: B7, A New Member Of The Ig Superfamily With
Unique Expression On Activated And Neoplastic B Cells
25
(C) JOURNAL: The Journal of Immunology
(D) VOLUME: 143
(E) ISSUE: 8
(F) PAGES: 2714-2722
30 (G) DATE: 15-OCT-1989
(H) RELEVANT RESIDUES In SEQ ID NO:28: FROM 1 TO 1491

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

35 CCAAAGAAAA AGTGATTTGT CATTGCTTTA TAGACTGTAA GAAGAGAACA TCTCAGAAGT 60
GGAGTCTTAC CCTGAAATCA AAGGATTTAA AGAAAAAGTG GAATTTTCTC TCAGCAAGCT 120
40 GTGAAACTAA ATCCACAACC TTTGGAGACC CAGGAACACC CTCCAATCTC TGTGTGTTTT 180
GTAAACATCA CTGGAGGGTC TTCTACGTGA GCAATTGGAT TGTCATCAGC CCTGCCTGTT 240
TTGCACCTGG GAAGTGCCCT GGTCTTACTT GGGTCCAAAT TGTGGCTTT CACTTTTGAC 300
45 CCTAAGCATC TGAAGCC ATG GGC CAC ACA CGG AGG CAG GGA ACA TCA CCA TCC 353
Met Gly His Thr Arg Arg Gln Gly Thr Ser Pro Ser
-30 -25
50
AAG TGT CCA TAC CTG AAT TTC TTT CAG CTC TTG GTG CTG GCT GGT CTT 401
Lys Cys Pro Tyr Leu Asn Phe Gln Leu Leu Val Leu Ala Gly Leu
55 -20 -15 -10

5	TCT CAC TTC TGT TCA GGT GTT ATC CAC GTG ACC AAG GAA GTG AAA GAA Ser His Phe Cys Ser Gly Val Ile His Val Thr Lys Glu Val Lys Glu -5 1 5 10	449
10	GTG GCA ACG CTG TCC TGT GGT CAC AAT GTT TCT GTT GAA GAG CTG GCA Val Ala Thr Leu Ser Cys Gly His Asn Val Ser Val Glu Glu Leu Ala 15 20 25	497
15	CAA ACT CGC ATC TAC TGG CAA AAG GAG AAG AAA ATG GTG CTG ACT ATG Gln Thr Arg Ile Tyr Trp Gln Lys Glu Lys Lys Met Val Leu Thr Met 30 35 40	545
20	ATG TCT GGG GAC ATG AAT ATA TGG CCC GAG TAC AAG AAC CGG ACC ATC Met Ser Gly Asp Met Asn Ile Trp Pro Glu Tyr Lys Asn Arg Thr Ile 45 50 55	593
25	TTT GAT ATC ACT AAT AAC CTC TCC ATT GTG ATC CTG GCT CTG CGC CCA Phe Asp Ile Thr Asn Asn Leu Ser Ile Val Ile Leu Ala Leu Arg Pro 60 65 70	641
30	TCT GAC GAG GGC ACA TAC GAG TGT GTT GTT CTG AAG TAT GAA AAA GAC Ser Asp Glu Gly Thr Tyr Glu Cys Val Val Leu Lys Tyr Glu Lys Asp 75 80 85 90	689
35	GCT TTC AAG CGG GAA CAC CTG GCT GAA GTG ACG TTA TCA GTC AAA GCT Ala Phe Lys Arg Glu His Leu Ala Glu Val Thr Leu Ser Val Lys Ala 95 100 105	737
40	GAC TTC CCT ACA CCT AGT ATA TCT GAC TTT GAA ATT CCA ACT TCT AAT Asp Phe Pro Thr Pro Ser Ile Ser Asp Phe Glu Ile Pro Thr Ser Asn 110 115 120	785
45	ATT AGA AGG ATA ATT TGC TCA ACC TCT GGA GGT TTT CCA GAG CCT CAC Ile Arg Arg Ile Ile Cys Ser Thr Ser Gly Gly Phe Pro Glu Pro His 125 130 135	833
50	CTC TCC TGG TTG GAA AAT GGA GAA GAA TTA AAT GCC ATC AAC ACA ACA Leu Ser Trp Leu Glu Asn Gly Glu Glu Leu Asn Ala Ile Asn Thr Thr 140 145 150	881
55	GTT TCC CAA GAT CCT GAA ACT GAG CTC TAT GCT GTT AGC AGC AAA CTG Val Ser Gln Asp Pro Glu Thr Glu Leu Tyr Ala Val Ser Ser Lys Leu	929

	155	160	165	170	
5	GAT TTC AAT ATG ACA ACC AAC CAC AGC TTC ATG TGT CTC ATC AAG TAT				977
	Asp Phe Asn Met Thr Thr Asn His Ser Phe Met Cys Leu Ile Lys Tyr				
		175	180	185	
10	GGA CAT TTA AGA GTG AAT CAG ACC TTC AAC TGG AAT ACA ACC AAG CAA				1025
	Gly His Leu Arg Val Asn Gln Thr Phe Asn Trp Asn Thr Thr Lys Gln				
		190	195	200	
15	GAG CAT TTT CCT GAT AAC CTG CTC CCA TCC TGG GCC ATT ACC TTA ATC				1073
	Glu His Phe Pro Asp Asn Leu Leu Pro Ser Trp Ala Ile Thr Leu Ile				
		205	210	215	
20	TCA GTA AAT GGA ATT TTT GTG ATA TGC TGC CTG ACC TAC TGC TTT GCC				1121
	Ser Val Asn Gly Ile Phe Val Ile Cys Cys Leu Thr Tyr Cys Phe Ala				
		220	225	230	
25	CCA AGA TGC AGA GAG AGA AGG AGG AAT GAG AGA TTG AGA AGG GAA AGT				1169
	Pro Arg Cys Arg Glu Arg Arg Arg Asn Glu Arg Leu Arg Arg Glu Ser				
		235	240	245	250
30	GTA CGC CCT GTA TAACAGTGTG CGCAGAAGCA AGGGGCTGAA AAGATCTGAA				1221
	Val Arg Pro Val				
	GGTAGCCTCC GTCATCTCTT CTGGGATACA TGGATCGTGG GGATCATGAG GCATTCTTCC				1281
35	CTTAACAAAT TTAAGCTGTT TTACCCACTA CCTCACCTTC TTAAAAACCT CTTTCAGATT				1341
	AAGCTGAACA GTTACAAGAT GGCTGGCATC CCTCTCCTTT CTCCCCATAT GCAATTGCT				1401
40	TAATGTAACC TCTTCTTTTG CCATGTTTCC ATTCTGCCAT CTTGAATTGT CTTGTCAGCC				1461
45	AATTCATTAT CTATTAAACA CTAATTTGAG				
	1491				
50	(3) INFORMATION FOR SEQ ID NO:29:				
	(i) SEQUENCE CHARACTERISTICS:				
	(A) LENGTH: 288 amino acids				
	(B) TYPE: amino acid				
55	(C) TOPOLOGY: linear				

(ii) MOLECULE TYPE: protein

(A) DESCRIPTION: B cell activation antigen; natural ligand
for CD28 T cell surface antigen; transmembrane protein

5

(ix) FEATURE:

(A) NAME/KEY: signal sequence

(B) LOCATION: -34 to -1

10 (C) IDENTIFICATION METHOD: amino terminal sequencing of
soluble protein

(D) OTHER INFORMATION: hydrophobic

(ix) FEATURE:

15

(A) NAME/KEY: extracellular domain

(B) LOCATION: 1 to 208

(C) IDENTIFICATION METHOD: similarity with known
sequence

20

(ix) FEATURE:

(A) NAME/KEY: transmembrane domain

25

(B) LOCATION: 209 to 235

(C) IDENTIFICATION METHOD: similarity with known
sequence

30

(ix) FEATURE:

(A) NAME/KEY: intracellular domain

(B) LOCATION: 236 to 254

35

(C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

40

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 19 to 21

(C) IDENTIFICATION METHOD: similarity with known
sequence

45

(ix) FEATURE:

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 55 to 57

50

(C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

55

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 64 to 66

(C) IDENTIFICATION METHOD: similarity with known
sequence

5

(ix) FEATURE:

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 152 to 154

10 (C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

15

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 173 to 175

(C) IDENTIFICATION METHOD: similarity with known
sequence

20

(ix) FEATURE:

(A) NAME/KEY: N-linked glycosylation

25

(B) LOCATION: 177 to 179

(C) IDENTIFICATION METHOD: similarity with known
sequence

30

(ix) FEATURE:

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 192 to 194

35 (C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

40

(A) NAME/KEY: N-linked glycosylation

(B) LOCATION: 198 to 200

(C) IDENTIFICATION METHOD: similarity with known
sequence

45

(ix) FEATURE:

(A) NAME/KEY: Ig V-set domain

(B) LOCATION: 1 to 104

50 (C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

55

(A) NAME/KEY: Ig C-set domain

(B) LOCATION: 105 to 202

(C) IDENTIFICATION METHOD: similarity with known sequence

5 (x) PUBLICATION INFORMATION:

(A) AUTHORS: FREEMAN, GORDON J.
 FREEDMAN, ARNOLD S.
 SEGIL, JEFFREY M.
 10 LEE, GRACE
 WHITMAN, JAMES F.
 NADLER, LEE M.

15 (B) TITLE: B7, A New Member Of The Ig Superfamily With
 Unique Expression On Activated And Neoplastic B Cells

(C) JOURNAL: The Journal of Immunology

(D) VOLUME: 143

(E) ISSUE: 8

(F) PAGES: 2714-2722

20 (G) DATE: 15-OCT-1989

(H) RELEVANT RESIDUES IN SEQUENCE ID NO:29: From -26 to 262

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

25

Met Gly His Thr Arg Arg Gln Gly Thr Ser Pro Ser Lys Cys Pro Tyr
 -30 -25 -20

30 Leu Asn Phe Phe Gln Leu Leu Val Leu Ala Gly Leu Ser His Phe Cys
 -15 -10 -5

Ser Gly Val Ile His Val Thr Lys Glu Val Lys Glu Val Ala Thr Leu
 -1 1 5 10

35 Ser Cys Gly His Asn Val Ser Val Glu Glu Leu Ala Gln Thr Arg Ile
 15 20 25 30

40 Tyr Trp Gln Lys Glu Lys Lys Met Val Leu Thr Met Met Ser Gly Asp
 35 40 45

Met Asn Ile Trp Pro Glu Tyr Lys Asn Arg Thr Ile Phe Asp Ile Thr
 50 55 60

45 Asn Asn Leu Ser Ile Val Ile Leu Ala Leu Arg Pro Ser Asp Glu Gly
 65 70 75

Thr Tyr Glu Cys Val Val Leu Lys Tyr Glu Lys Asp Ala Phe Lys Arg
 80 85 90

50 Glu His Leu Ala Glu Val Thr Leu Ser Val Lys Ala Asp Phe Pro Thr
 95 100 105 110

55 Pro Ser Ile Ser Asp Phe Glu Ile Pro Thr Ser Asn Ile Arg Arg Ile
 115 120 125

-113-

Ile Cys Ser Thr Ser Gly Gly Phe Pro Glu Pro His Leu Ser Trp Leu
 130 135 140
 5 Glu Asn Gly Glu Glu Leu Asn Ala Ile Asn Thr Thr Val Ser Gln Asp
 145 150 155

 Pro Glu Thr Glu Leu Tyr Ala Val Ser Ser Lys Leu Asp Phe Asn Met
 160 165 170
 10 Thr Thr Asn His Ser Phe Met Cys Leu Ile Lys Tyr Gly His Leu Arg
 175 180 185 190

 15 Val Asn Gln Thr Phe Asn Trp Asn Thr Thr Lys Gln Glu His Phe Pro
 195 200 205

 Asp Asn Leu Leu Pro Ser Trp Ala Ile Thr Leu Ile Ser Val Asn Gly
 210 215 220
 20 Ile Phe Val Ile Cys Cys Leu Thr Tyr Cys Phe Ala Pro Arg Cys Arg
 225 230 235

 25 Glu Arg Arg Arg Asn Glu Arg Leu Arg Arg Glu Ser Val Arg Pro Val
 240 245 250

30 (4) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 1716 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: cDNA to mRNA

40 (iii) HYPOTHETICAL: no

(vi) ORIGINAL SOURCE:

45 (A) ORGANISM: Mus musculus
 (D) DEVELOPMENTAL STAGE: germ line
 (F) TISSUE TYPE: lymphoid
 (G) CELL TYPE: B lymphocyte
 (H) CELL LINE: 70Z and A20

50 (vii) IMMEDIATE SOURCE:

(A) LIBRARY: cDNA in pCDM8 vector
 (B) CLONE: B7 #'s 1 and 29

55 (ix) FEATURE:

- (A) NAME/KEY: translated region
 (B) LOCATION: 249 to 1166 bp
 (C) IDENTIFICATION METHOD: similarity to other pattern

5 (ix) FEATURE:

- (A) NAME/KEY: Alternate ATG initiation codons
 (B) LOCATION: 225 to 227 and 270 to 272
 (C) IDENTIFICATION METHOD: similarity to other pattern

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

```

15  GAGTTTATA CCTCAATAGA CTCTTACTAG TTTCTCTTTT TCAGGTTGTG AAACCTCAACC   60
    TTCAAAGACA CTCTGTTCCA TTTCTGTGGA CTAATAGGAT CATCTTTAGC ATCTGCCGGG  120
    TGGATGCCAT CCAGGCTTCT TTTTCTACAT CTCTGTTTCT CGATTTTGTG GAGCCTAGGA  180
20  GGTGCCTAAG CTCCATTGGC TCTAGATTCC TGGCTTTCCC CATCATGTTC TCCAAAGCAT   240
    CTGAAGCT ATG GCT TGC AAT TGT CAG TTG ATG CAG GAT ACA CCA CTC CTC   290
        Met Ala Cys Asn Cys Gln Leu Met Gln Asp Thr Pro Leu Leu
        -35                -30                -25

    AAG TTT CCA TGT CCA AGG CTC AAT CTT CTC TTT GTG CTG CTG ATT CGT   338
    Lys Phe Pro Cys Pro Arg Leu Ile Leu Leu Phe Val Leu Leu Ile Arg
        -20                -15                -10

30  CTT TCA CAA GTG TCT TCA GAT GTT GAT GAA CAA CTG TCC AAG TCA GTG   386
    Leu Ser Gln Val Ser Ser Asp Val Asp Glu Gln Leu Ser Lys Ser Val
        -5                -1  1                5

35  AAA GAT AAG GTA TTG CTG CCT TGC CGT TAC AAC TCT CCT CAT GAA GAT   434
    Lys Asp Lys Val Leu Leu Pro Cys Arg Tyr Asn Ser Pro His Glu Asp
        10                15                20                25

40  GAG TCT GAA GAC CGA ATC TAC TGG CAA AAA CAT GAC AAA GTG GTG CTG   482
    Glu Ser Glu Asp Arg Ile Tyr Trp Gln Lys His Asp Lys Val Val Leu
        30                35                40

    TCT GTC ATT GCT GGG AAA CTA AAA GTG TGG CCC GAG TAT AAG AAC CGG   530
    Ser Val Ile Ala Gly Lys Leu Lys Val Trp Pro Glu Tyr Lys Asn Arg
        45                50                55

    ACT TTA TAT GAC AAC ACT ACC TAC TCT CTT ATC ATC CTG GGC CTG GTC   578
    Thr Leu Tyr Asp Asn Thr Thr Tyr Ser Leu Ile Ile Leu Gly Leu Val
        60                65                70

50  CTT TCA GAC CGG GGC ACA TAC AGC TGT GTC GTT CAA AAG AAG GAA AGA   626
    Leu Ser Asp Arg Gly Thr Tyr Ser Cys Val Val Gln Lys Lys Glu Arg
        75                80                85

55  GGA ACG TAT GAA GTT AAA CAC TTG GCT TTA GTA AAG TTG TCC ATC AAA   674
    Gly Thr Tyr Glu Val Lys His Leu Ala Leu Val Lys Leu Ser Ile Lys

```


	90		95		100		105	
	GCT GAC TTC TCT ACC CCC AAC ATA ACT GAG TCT GGA AAC CCA TCT GCA							722
	Ala Asp Phe Ser Thr Pro Asn Ile Thr Glu Ser Gly Asn Pro Ser Ala							
5		110			115		120	
	GAC ACT AAA AGG ATT ACC TGC TTT GCT TCC GGG GGT TTC CCA AAG CCT							770
	Asp Thr Lys Arg Ile Thr Cys Phe Ala Ser Gly Gly Phe Pro Lys Pro							
10		125			130		135	
	CGC TTC TCT TGG TTG GAA AAT GGA AGA GAA TTA CCT GGC ATC AAT ACG							818
	Arg Phe Ser Trp Leu Glu Asn Gly Arg Glu Leu Pro Gly Ile Asn Thr							
15		140			145		150	
	ACA ATT TCC CAG GAT CCT GAA TCT GAA TTG TAC ACC ATT AGT AGC CAA							866
20	Thr Ile Ser Gln Asp Pro Glu Ser Glu Leu Tyr Thr Ile Ser Ser Gln							
		155			160		165	
	CTA GAT TTC AAT ACG ACT CGC AAC CAC ACC ATT AAG TGT CTC ATT AAA							914
	Leu Asp Phe Asn Thr Thr Arg Asn His Thr Ile Lys Cys Leu Ile Lys							
25		170			175		180	
	TAT GGA GAT GCT CAC GTG TCA GAG GAC TTC ACC TGG GAA AAA CCC CCA							962
	Tyr Gly Asp Ala His Val Ser Glu Asp Phe Thr Trp Glu Lys Pro Pro							
		190			195		200	
30	GAA GAC CCT CCT GAT AGC AAG AAC ACA CTT GTG CTC TTT GGG GCA GGA							1010
	Glu Asp Pro Pro Asp Ser Lys Asn Thr Leu Val Leu Phe Gly Ala Gly							
		205			210		215	
35	TTC GGC GCA GTA ATA ACA GTC GTC GTC ATC GTT GTC ATC ATC AAA TGC							1058
	Phe Gly Ala Val Ile Thr Val Val Val Ile Val Val Ile Ile Lys Cys							
		220			225		230	
40	TTC TGT AAG CAC AGA AGC TGT TTC AGA AGA AAT GAG GCA AGC AGA GAA							1106
	Phe Cys Lys His Arg Ser Cys Phe Arg Arg Asn Glu Ala Ser Arg Glu							
		235			240		245	
45	ACA AAC AAC AGC CTT ACC TTC GGG CCT GAA GAA GCA TTA GCT GAA CAG							1154
	Thr Asn Asn Ser Leu Thr Phe Gly Pro Glu Glu Ala Leu Ala Glu Gln							
		250			255		260	
	ACC GTC TTC CTT TAGTTCTTCT CTGTCCATGT GGGATACATG GTATTATGTG							1206
	Thr Val Phe Leu							
50	GCTCATGAGG TACAATCTTT CTTTCAGCAC CGTGCTAGCT GATCTTTCGG ACAACTTGAC							1266
	ACAAGATAGA GTTAACTGGG AAGAGAAAGC CTTGAATGAG GATTTCTTTC CATCAGGAAG							1326
	CTACGGGCAA GTTTGCTGGG CCTTTGATTG CTTGATGACT GAAGTGGAAA GGCTGAGCCC							1386
55	ACTGTGGGTG GTGCTAGCCC TGGGCAGGGG CAGGTGACCC TGGGTGGTAT AAGAAAAAGA							1446

GCTGTCACTA AAAGGAGAGG TGCCTAGTCT TACTGCAACT TGATATGTCA TGTTTGGTTG 1506
GTGTCTGTGG GAGGCCTGCC CTTTTCTGAA GAGAAGTGGT GGGAGAGTGG ATGGGGTGGG 1566
5 GGCAGAGGAA AAGTGGGGGA GAGGCCTGG GAGGAGAGGA GGGAGGGGGA CGGGGTGGGG 1626
GTGGGGAAAA CTATGTTGG GATGTAAAAA CGGATAATAA TATAATATT AAATAAAAAG 1686
AGAGTATTGA GCAAAAAAAA AAAAAAAAAA
10 1716

15 (5) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 306 amino acids
(B) TYPE: amino acid
(C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

25 (A) DESCRIPTION: B lymphocyte activation antigen; Ig
superfamily member; T cell costimulatory signal
via activation of CD28 pathways, binds to CD28⁺
T cells, transmembrane protein

30 (ix) FEATURE:

(A) NAME/KEY: signal sequence
(B) LOCATION: -37 to -1
35 (C) IDENTIFICATION METHOD: similarity with known
sequence
(D) OTHER INFORMATION: hydrophobic

(ix) FEATURE:

40 (A) NAME/KEY: extracellular domain
(B) LOCATION: 1 to 210
(C) IDENTIFICATION METHOD: similarity with known
sequence

45

(ix) FEATURE:

50 (A) NAME/KEY: transmembrane domain
(B) LOCATION: 211 to 235
(C) IDENTIFICATION METHOD: similarity with known
sequence

(ix) FEATURE:

55

(A) NAME/KEY: intracellular (cytoplasmic) domain

(B) LOCATION: 236 to 269

(C) IDENTIFICATION METHOD: similarity with known sequence

5

(ix) FEATURE:

(A) NAME/KEY: Ig V-set domain

(B) LOCATION: 1 to 105

10 (C) IDENTIFICATION METHOD: similarity with known sequence

(ix) FEATURE:

15

(A) NAME/KEY: Ig C-set domain

(B) LOCATION: 106 to 199

(C) IDENTIFICATION METHOD: similarity with known sequence

20

(x) PUBLICATION INFORMATION:

(A) AUTHORS: FREEMAN, GORDON J.

25

GRAY, GARY S.

GIMMI, CLAUDE D.

LOMBARD, DAVID B.

ZHOU, LIANG-JI

WHITE, MICHAEL

30

FINGEROTH, JOYCE D.

GRIBBEN, JOHN G.

NADLER, LEE M.

(B) TITLE: Structure, Expression, and T Cell Costimulatory Activity Of The Murine Homologue Of The Human B Lymphocyte Activation Antigen B7

35

(C) JOURNAL: Journal of Experimental Medicine

(D) VOLUME:

(E) ISSUE:

40

(F) PAGES:

(G) DATE: IN PRESS

(H) RELEVANT RESIDUES IN SEQUENCE ID NO:31: From -37 to 269

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Met Ala Cys Asn Cys Gln Leu Met Gln Asp Thr Pro Leu Leu Lys Phe
-35 -30 -25

50

Pro Cys Pro Arg Leu Ile Leu Leu Phe Val Leu Leu Ile Arg Leu Ser
-20 -15 -10

55

Gln Val Ser Ser Asp Val Asp Glu Gln Leu Ser Lys Ser Val Lys Asp
-5 -1 1 5 10

-118-

Lys Val Leu Leu Pro Cys Arg Tyr Asn Ser Pro His Glu Asp Glu Ser
 15 20 25
 5 Glu Asp Arg Ile Tyr Trp Gln Lys His Asp Lys Val Val Leu Ser Val
 30 35 40
 Ile Ala Gly Lys Leu Lys Val Trp Pro Glu Tyr Lys Asn Arg Thr Leu
 45 50 55
 10 Tyr Asp Asn Thr Thr Tyr Ser Leu Ile Ile Leu Gly Leu Val Leu Ser
 60 65 70 75
 Asp Arg Gly Thr Tyr Ser Cys Val Val Gln Lys Lys Glu Arg Gly Thr
 80 85 90
 15 Tyr Gly Val Lys His Leu Ala Leu Val Lys Leu Ser Ile Lys Ala Asp
 95 100 105
 Phe Ser Thr Pro Asn Ile Thr Glu Ser Gly Asn Pro Ser Ala Asp Thr
 110 115 120
 Lys Arg Ile Thr Cys Phe Ala Ser Gly Gly Phe Pro Lys Pro Arg Phe
 125 130 135
 25 Ser Trp Leu Glu Asn Gly Arg Glu Leu Pro Gly Ile Asn Thr Thr Ile
 140 145 150 155
 Ser Gln Asp Pro Glu Ser Glu Leu Tyr Thr Ile Ser Ser Gln Leu Asp
 160 165 170
 30 Phe Asn Thr Thr Arg Asn His Thr Ile Lys Cys Leu Ile Lys Tyr Gly
 175 180 185
 Asp Ala His Val Ser Glu Asp Phe Thr Trp Glu Lys Pro Pro Glu Asp
 190 195 200
 Pro Pro Asp Ser Lys Asn Thr Leu Val Leu Phe Gly Ala Gly Phe Gly
 205 210 215
 40 Ala Val Ile Thr Val Val Val Ile Val Val Ile Ile Lys Cys Phe Cys
 220 225 230 235
 Lys His Arg Ser Cys Phe Arg Arg Asn Glu Ala Ser Arg Glu Thr Asn
 240 245 250
 45 Asn Ser Leu Thr Phe Gly Pro Glu Glu Ala Leu Ala Glu Gln Thr Val
 255 260 265
 Phe Leu
 50

(2) INFORMATION FOR SEQ ID NO:32:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: oligonucleotide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GGCACTAGGT CTCCAGCTTG AGATCACAGT TCTCTCTAC

39

(2) INFORMATION FOR SEQ ID NO:33:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: oligonucleotide

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GCTTGAATCT TCAGAGGAGC GGAGTGGACA CCTGTGG

37

(2) INFORMATION FOR SEQ ID NO:34:

30

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: oligonucleotide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GCTCCTCTGA AGATTCAAGC

20

45 (2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: cDNA

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GGCACTATGA TCAGGGGGAG GCTGAGGTCC 30

5 (2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 78 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

20 GCATTTTAAG CTTTTTCCTG ATCAGGAGCC CAAATCTTCT GACAAAAC TC ACACATCTCC 60

ACCGTCTCCA GGTAAGCC 78

(2) INFORMATION FOR SEQ ID NO:37:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

TAATACGACT CACTATAGGG 20

(2) INFORMATION FOR SEQ ID NO:38:

40

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GAGCATTTTC CTGATCAGGA GTCCAAATAT GGTCCCCCAT CCCATCATCC CCAGGTAAGC 60

55 CAACCC 66

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

15

GCAGAGGAAT CGAGCTCGGT ACCCGGGGAT CCCAGTGTG GGGACAGTGG GACCGCTCTG 60
CCTCCC 66

(2) INFORMATION FOR SEQ ID NO:40:

20

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 59 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GGGTTTTGGG GGGAAGAGGA AGACTGACGG TGCCCCCTCG GCTTCAGGTG CTGAGGAAG 59

35

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

40

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

CATCTCTTCC TCAGCACCTG AAGCCGAGGG GGCACCGTCA GTCTTCCTCT TCCCCC 56

50

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

55

- (A) LENGTH: 99 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

CGCACGTGAC CTCAGGGGTC CGGGAGATCA TGAGAGTGTC CTTGGGTTTT GGGGGGAACA 60
 10 GGAAGACTGA TGGTGCCCCC TCGAACTCAG GTGCTGAGG 99

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 98 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

25 CCTCAGCACC TGAGTTCGAG GGGGCACCAT CAGTCTCCTG TTCCCCCAA AACCCAAGGA 60
 CACTCTCATG ATCTCCCGGA CCCCTGAGGT CACGTGCG 98

30 (2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 330 base pairs
 35 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..330

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GCT CCT CTG AAG ATT CAA GCT TAT TTC AAT GAG ACT GCA GAC CTG CCA 48
 Ala Pro Leu Lys Ile Gln Ala Tyr Phe Asn Glu Thr Ala Asp Leu Pro
 50 1 5 10 15
 TGC CAA TTT GCA AAC TCT CAA AAC CAA AGC CTG AGT GAG CTA GTA GTA 96
 Cys Gln Phe Ala Asn Ser Gln Asn Gln Ser Leu Ser Glu Leu Val Val
 20 25 30
 55 TTT TGG CAG GAC CAG GAA AAC TTG GTT CTG AAT GAG GTA TAC TTA GGC 144

Phe Trp Gln Asp Gln Glu Asn Leu Val Leu Asn Glu Val Tyr Leu Gly
 35 40 45

5 AAA GAG AAA TTT GAC AGT GTT CAT TCC AAG TAT ATG GGC CGC ACA AGT 192
 Lys Glu Lys Phe Asp Ser Val His Ser Lys Tyr Met Gly Arg Thr Ser
 50 55 60

10 TTT GAT TCG GAC AGT TGG ACC CTG AGA CTT CAC AAT CTT CAG ATC AAG 240
 Phe Asp Ser Asp Ser Trp Thr Leu Arg Leu His Asn Leu Gln Ile Lys
 65 70 75 80

15 GAC AAG GGC TTG TAT CAA TGT ATC ATC CAT CAC AAA AAG CCC ACA GGA 288
 Asp Lys Gly Leu Tyr Gln Cys Ile Ile His His Lys Lys Pro Thr Gly
 85 90 95

ATG ATT CGC ATC CAC CAG ATG AAT TCT AGG CTG TCA GTG CTT 330
 Met Ile Arg Ile His Gln Met Asn Ser Arg Leu Ser Val Leu
 100 105 110

20

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 110 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: protein

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Ala Pro Leu Lys Ile Gln Ala Tyr Phe Asn Glu Thr Ala Asp Leu Pro
 1 5 10 15

35 Cys Gln Phe Ala Asn Ser Gln Asn Gln Ser Leu Ser Glu Leu Val Val
 20 25 30

Phe Trp Gln Asp Gln Glu Asn Leu Val Leu Asn Glu Val Tyr Leu Gly
 35 40 45

40 Lys Glu Lys Phe Asp Ser Val His Ser Lys Tyr Met Gly Arg Thr Ser
 50 55 60

45 Phe Asp Ser Asp Ser Trp Thr Leu Arg Leu His Asn Leu Gln Ile Lys
 65 70 75 80

Asp Lys Gly Leu Tyr Gln Cys Ile Ile His His Lys Lys Pro Thr Gly
 85 90 95

50 Met Ile Arg Ile His Gln Met Asn Ser Arg Leu Ser Val Leu
 100 105 110

(2) INFORMATION FOR SEQ ID NO:46:

55

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 306 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- 10 (B) LOCATION: 1..310

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

15	GCT AAC TTC AGT CAA CCT GAA ATA GTA CCA ATT TCT AAT ATA ACA GAA	48
	Ala Asn Phe Ser Gln Pro Glu Ile Val Pro Ile Ser Asn Ile Thr Glu	
	1 5 10 15	
20	AAT GTG TAC ATA AAT TTG ACC TGC TCA TCT ATA CAC GGT TAC CCA GAA	96
	Asn Val Tyr Ile Asn Leu Thr Cys Ser Ser Ile His Gly Tyr Pro Glu	
	20 25 30	
25	CCT AAG AAG ATG AGT GTT TTG CTA AGA ACC AAG AAT TCA ACT ATC GAG	144
	Pro Lys Lys Met Ser Val Leu Leu Arg Thr Lys Asn Ser Thr Ile Glu	
	35 40 45	
30	TAT GAT GGT ATT ATG CAG AAA TCT CAA GAT AAT GTC ACA GAA CTG TAC	192
	Tyr Asp Gly Ile Met Gln Lys Ser Gln Asp Asn Val Thr Glu Leu Tyr	
	50 55 60	
35	GAC GTT TCC ATC AGC TTG TCT GTT TCA TTC CCT GAT GTT ACG AGC AAT	240
	Asp Val Ser Ile Ser Leu Ser Val Ser Phe Pro Asp Val Thr Ser Asn	
	65 70 75 80	
40	ATG ACC ATC TTC TGT ATT CTG GAA ACT GAC AAG ACG CGG CTT TTA TCT	288
	Met Thr Ile Phe Cys Ile Leu Glu Thr Asp Lys Thr Arg Leu Leu Ser	
	85 90 95	
45	TCA CCT TTC TCT ATA GAG	306
	Ser Pro Phe Ser Ile Glu	
	100	

(2) INFORMATION FOR SEQ ID NO:47:

- 45 (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 103 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

55	Ala Asn Phe Ser Gln Pro Glu Ile Val Pro Ile Ser Asn Ile Thr Glu
	1 5 10 15

Asn Val Tyr Ile Asn Leu Thr Cys Ser Ser Ile His Gly Tyr Pro Glu
 20 25 30
 5 Pro Lys Lys Met Ser Val Leu Leu Arg Thr Lys Asn Ser Thr Ile Glu
 35 40 45
 Tyr Asp Gly Ile Met Gln Lys Ser Gln Asp Asn Val Thr Glu Leu Tyr
 50 55 60
 10 Asp Val Ser Ile Ser Leu Ser Val Ser Phe Pro Asp Val Thr Ser Asn
 65 70 75 80
 15 Met Thr Ile Phe Cys Ile Leu Glu Thr Asp Lys Thr Arg Leu Leu Ser
 85 90 95
 Ser Pro Phe Ser Ile Glu
 100

20 (2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 base pairs
 (B) TYPE: nucleic acid
 25 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

35 GCAACCGGAA GCTTGCCACC ATGGGGGTAC TGCTCACACA GAGGACG

47

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 63 base pairs
 40 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

50 AGTCTCAT TG AAATAAGCTT GAATCTTCAG AGGAGCCATG CTGGCCATGC TTGGAAACAG
 GAG

60

63

(2) INFORMATION FOR SEQ ID NO:50:

55

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 62 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: oligonucleotide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

CTCCTGTTTC CAAGCATGGC CAGCATGGCT CCTCTGAAGA TTCAGGCTTA TTTCAATGAG 60
AC 62

15 (2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 51 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: oligonucleotide

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

30 TGTGTGTGGA ATTCTCATTA CTGATCAAGC ACTGACAGTT CAGAATTCAT C 51

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 63 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: oligonucleotide

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

45 AGAAATTGGT ACTATTTTCAG GTTGA CTGACTGAA GTTAGCCATG CTGGCCATGC TTGGAACAG 60
GAG 63

50 (2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 62 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: oligonucleotide

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CTCCTGTTTC CAAGCATGGC CAGCATGGCT AACTTCAGTC AACCTGAAAT AGTACCAATT 60

10 TC 62

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 105 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

CTCAAGCTTG CCACCATGGG GGTACTGCTC ACACAGAGGA CGCTGCTCAG TCTGGTCCTT 60

25 GCACTCCTGT TTCCGAGCAT GGCGAGCATG GGTCTTTCTC ACTTC 105

30 (2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: oligonucleotide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

TGTGTGTGGA ATTCTCATTA CTGATCAGGA AAATGCTCTT GCTTG 45

CLAIMS

1. An isolated nucleic acid encoding a B7-2 fusion protein comprising a
5 nucleotide sequence encoding a first peptide having a B7-2 activity and a nucleotide sequence
encoding a second peptide corresponding to a moiety that alters the solubility, binding
affinity or valency of the first peptide.
2. The isolated nucleic acid of claim 1 which is a DNA.
- 10 3. The isolated nucleic acid of claim 2, wherein the first peptide
comprises an extracellular domain of a human B7-2 protein.
4. The isolated nucleic acid of claim 3, wherein the first peptide
15 comprises amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2).
5. The isolated nucleic acid of claim 3, wherein the first peptide
comprises a variable region-like domain of human B7-2.
- 20 6. The isolated nucleic acid of claim 3, wherein the first peptide
comprises a constant region-like domain of human B7-2.
7. The isolated nucleic acid of claim 2, wherein the second peptide
comprises an immunoglobulin constant region.
- 25 8. The isolated nucleic acid of claim 7, wherein the immunoglobulin
constant region is a C γ 1 domain, including the hinge, CH2 and CH3 region.
9. The isolated nucleic acid of claim 7, wherein the immunoglobulin
30 constant region is modified to reduce constant region-mediated biological effector functions.
10. The isolated nucleic acid of claim 9, wherein the biological effector
function is selected from the group consisting of complement activation, Fc receptor
interaction, and complement activation and Fc receptor interaction.

-129-

11. The isolated nucleic acid of claim 10, wherein the immunoglobulin constant region is a C γ 4 domain, including the hinge, CH2 and CH3 region.
- 5 12. The isolated nucleic acid of claim 11, wherein at least one amino acid residue of the CH2 domain is modified by substitution, addition or deletion.
- 10 13. An isolated B7-2 fusion protein comprising a first peptide having a B7-2 activity and a second peptide corresponding to a moiety that alters the solubility, binding affinity or valency of the first peptide.
14. The isolated B7-2 fusion protein of claim 13, wherein the first peptide comprises an extracellular domain of human B7-2 protein.
- 15 15. The isolated B7-2 fusion protein of claim 14, wherein the first peptide comprises amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2).
16. The isolated B7-2 fusion protein of claim 14, wherein the first peptide comprises a variable region-like domain of human B7-2.
- 20 17. The isolated B7-2 fusion protein of claim 14, wherein the first peptide comprises a constant region-like domain of human B7-2.
18. The isolated B7-2 fusion protein of claim 13, wherein the second
- 25 peptide comprises an immunoglobulin constant region.
19. The isolated B7-2 fusion protein of claim 18, wherein the immunoglobulin constant region is a C γ 1 domain, including the hinge, CH2 and CH3 region.

20. The isolated B7-2 fusion protein of claim 18, wherein the immunoglobulin constant region is modified to reduce constant region-mediated biological effector functions.
- 5 21. The isolated B7-2 fusion protein of claim 20, wherein the biological effector function is selected from the group consisting of complement activation, Fc receptor interaction, and complement activation and Fc receptor interaction.
- 10 22. The isolated B7-2 fusion protein of claim 21, wherein the immunoglobulin constant region is a C γ 4 domain, including the hinge, CH2 and CH3 region.
23. The isolated B7-2 fusion protein of claim 22, wherein at least one amino acid residue of the CH2 domain is modified by substitution, addition or deletion.
- 15 24. A composition suitable for pharmaceutical administration comprising a fusion protein of claim 13 and a pharmaceutically acceptable carrier.
25. A composition suitable for pharmaceutical administration comprising a fusion protein of claim 14 and a pharmaceutically acceptable carrier.
- 20 26. A composition suitable for pharmaceutical administration comprising a fusion protein of claim 16 and a pharmaceutically acceptable carrier.
- 25 27. A composition suitable for pharmaceutical administration comprising a fusion protein of claim 18 and a pharmaceutically acceptable carrier.

-131-

28. A method for inhibiting an interaction of a B lymphocyte antigen, B7-2, with its natural ligand(s) on the surface of immune cells, comprising contacting an immune cell with a B7-2 fusion protein which inhibits B7-2 binding with its natural ligand(s), to
5 thereby inhibit costimulation of the immune cell through the B7-2-ligand interaction.

29. The method of claim 28, wherein the B7-2 fusion protein comprises a first peptide having B7-2 activity and a second peptide comprising a moiety that alters the solubility, binding affinity or valency of the first peptide.

10

30. The method of claim 29, wherein the first peptide comprises an extracellular domain of the human B7-2 protein.

31. The method of claim 30, wherein the first peptide comprises amino
15 acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2).

32. The method of claim 29, wherein the second peptide comprises an immunoglobulin constant region.

20

33. The method of claim 32, wherein the immunoglobulin constant region is a C γ 1 domain, including the hinge, CH2 and CH3 region.

34. A method for treating an autoimmune disease in a subject mediated by interaction of a B lymphocyte antigen, B7-2, with its natural ligand(s) on the surface of
25 immune cells, comprising administering to the subject an inhibitory form of a B7-2 fusion protein, to thereby inhibit costimulation of the immune cells through the B7-2-ligand interaction.

35. The method of claim 34, wherein the inhibitory form of a B7-2 fusion
30 protein is a B7-2 immunoglobulin fusion protein (B7-2Ig) comprising a first peptide comprising an extracellular domain of the B7-2 protein and a second peptide comprising an immunoglobulin constant domain.

36. The method of claim 35, wherein the extracellular domain of the B7-2 protein comprises amino acid residues 24-245 of the sequence shown in Figure 8 (SEQ ID NO:2).

5

37. A method for treating allergy in a subject mediated by interaction of a B lymphocyte antigen, B7-2, with its natural ligand(s) on the surface of immune cells, comprising administering to the subject an inhibitory form of a B7-2 fusion protein, to thereby inhibit costimulation of the immune cells through the B7-2 -ligand interaction.

10

38. An isolated variable region form of the B cell activation antigen B7-2 which comprises a B7-2 immunoglobulin-like variable region domain but does not comprise a B7-2 immunoglobulin-like constant region domain.

15

39. The B7-2 variable region form of claim 38, which is human.

40. The B7-2 variable region form of claim 38, which is a fusion protein comprising a B7-2 variable region polypeptide operatively linked to a heterologous polypeptide.

20

41. The B7-2 variable region form of claim 40, wherein the B7-2 variable region polypeptide is a human B7-2 variable region polypeptide.

42. The B7-2 variable region form of claim 41, wherein the human B7-2 variable region polypeptide comprises an amino acid sequence of about positions 24 to 133 of SEQ ID NO: 2.

25

43. The B7-2 variable region form of claim 40, wherein the heterologous polypeptide comprises an immunoglobulin constant region.

30

44. The B7-2 variable region form of claim 43, wherein the immunoglobulin constant region comprises the hinge, CH2 and CH3 domains of IgG1.

45. The B7-2 variable region form of claim 38, comprising a B7-2 immunoglobulin-like variable region domain operatively linked to a transmembrane domain, the B7-2 variable region form being expressed on the surface of a cell.

35

-133-

46. The B7-2 variable region form of claim 45, further comprising a non-B7-2 linker polypeptide located between the B7-2 immunoglobulin-like variable region domain and the transmembrane domain.
- 5 47. The B7-2 variable region form of claim 45, further comprising a cytoplasmic domain.
48. The B7-2 variable region form of claim 38, comprising a B7-2
10 immunoglobulin-like variable region domain bound to a solid support.
49. The B7-2 variable region form of claim 48, wherein the solid support is a bead or plate.
- 15 50. The B7-2 variable region form of claim 48, further comprising a non-B7-2 linker polypeptide located between the B7-2 immunoglobulin-like variable region domain and the solid support.
- 20 51. An isolated B7-2 fusion protein comprising a human B7-2 immunoglobulin-like variable region domain operatively linked to a heterologous polypeptide, wherein the B7-2 fusion protein does not comprise a B7-2 immunoglobulin-like constant region domain.
- 25 52. The B7-2 fusion protein of claim 51, wherein the human B7-2 immunoglobulin-like variable region domain comprises an amino acid sequence from about position 24 to position 133 of SEQ ID NO: 2.
- 30 53. The B7-2 fusion protein of claim 51, wherein the heterologous polypeptide comprises an immunoglobulin constant region polypeptide.
54. An isolated nucleic acid molecule encoding a variable region form of a B7-2 fusion protein, the B7-2 fusion protein comprising a human B7-2 immunoglobulin-like variable region domain operatively linked to a heterologous polypeptide, wherein the B7-2 fusion protein does not comprise a B7-2 immunoglobulin-like constant region domain.

55. The nucleic acid of claim 58, wherein the heterologous polypeptide is an immunoglobulin constant region polypeptide.
- 5 56. A recombinant expression vector comprising the nucleic acid molecule of claim 54.
57. A host cell containing the recombinant expression vector of claim 56.
- 10 58. An isolated nucleic acid molecule encoding a variable region form of B7-2, the nucleic acid comprising a contiguous nucleotide sequence encoding a signal peptide, a human B7-2 immunoglobulin-like variable region domain, a transmembrane domain and a cytoplasmic domain.
- 15 59. The nucleic acid molecule of claim 58, wherein the human B7-2 immunoglobulin-like variable region domain comprises an amino acid sequence from about position 24 to position 133 of SEQ ID NO: 2.
- 20 60. The nucleic acid molecule of claim 58, further comprising a nucleotide sequence encoding a non-B7-2 linker polypeptide located between the nucleotide sequence encoding the B7-2 immunoglobulin-like variable region domain and the transmembrane domain.
- 25 61. A recombinant expression vector comprising the nucleic acid molecule of claim 58.
62. A host cell containing the recombinant expression vector of claim 61, wherein the variable region form of B7-2 is expressed on the surface of the cell.

-135-

63. A method for stimulating a response by an activated T cell, comprising contacting the activated T cell with a variable region form of the B cell activation antigen B7-2, the variable region form of B7-2 comprising a B7-2 immunoglobulin-like variable region domain but not comprising a B7-2 immunoglobulin-like constant region domain such that a
5 response by the activated T cell is stimulated.

64. The method of claim 63, wherein a T_{helper} -Type 2 (TH_2) response is preferentially stimulated.

1/34

FIG. 1A





FIG. 1B



SUBSTITUTE SHEET (RULE 26)

2/34

FIG. 2C




FIG. 2B





FIG. 2A



3/34

FIG. 3A





FIG. 3B



4/34




FIG. 4

B Cells Activated by sIg Crosslinking





FIG. 5

6/34

Cell Number

FIG. 6

B Cells Activated by Class II Crosslinking



7/34

FIG. 7A





FIG. 7B



SUBSTITUTE SHEET (RULE 26)

9/34


FIG. 8B

421 GGGCTTGATCAATGTATCATCCATCACAAAAGCCACAGGAATGATTCGCATCCACCAGATGAATTCT 490
 106 G L Y Q C I I H H K K P T G M I R I H Q M N S 128
 491 GAACTGTCAGTCTTGCTAACTTCAGTCAACCTGAAATAGTACCAATTTCTAATAACAGAAAATGTGT 560
 129 E L S V L A N F S Q P E I V P I S N I T E N V 151
 561 ACATAAATTGACCTGCTCATCTATACACGGTTACCCAGAACCTAAGAAGATGAGTGTTCGCTAAGAAC 630
 152 Y I N L T C S S I H G Y P E P K K M S V L L R T 175
 631 CAAGAATTCAACTATCGAGTATGATGGTATTATGCAGAAATCTCAAGATAATGTCACAGAACTGTACGAC 700
 176 K N S T I E Y D G I M Q K S Q D N V T E L Y D 198
 701 GTTCCATCAGCTGTCTGTTTCATTCCCTGAATGTTACGAGCAATATGACCATCTTCGTATTCCTGGAAA 770
 199 V S I S L S V S F P D V T S N M T I F C I L E 221
 771 CTGACAAGACGGGCTTTTATCTTCACCTTCTCTATAGAGCTTGAGGACCCCTCAGCCTCCCCCAGACCA 840
 222 T D K T R L L S S P F S I E L E D P Q P P D H 245

FIG. 8C

841	CATTCC	TTGGATTACAGCTGTA	CTTCCAAACAGTTATATATG	TGCTGATGGTTTCTG	TCTAATTC	TATGG	910
246	<u>I P W I T A V L P T V I I C V M V F C L I L W</u>						268
911	AAATGGAAGAAGAGCGGCCTCGCAACTCTTTATAAATGTGGAACCAACACAATCGAGAGGGAAGAGA						980
269	K W K K K R P R N S Y K C G T N T M E R E E						291
981	GTGAACAGACCAAGAAAGAGAAAAATCCATATACCTGAAAGATCTGATGAAGCCAGCGTGTTTTAA						1050
292	S E Q T K K R E K I H I P E R S D E A Q R V F K						315
11051	AAGTTCGAAGACATCTTCATCGGACAAAAGTGATACATGTTTTTAATTAAGAGTAAAGCCCAAAAAA						1120
316	S S K T S S C D K S D T C F *						329

11/34






FIG. 10A




FIG. 10B

13/34

FIG. IIA






FIG. IIB




SUBSTITUTE SHEET (RULE 26)

14/34



SUBSTITUTE SHEET (RULE 26)

15/34



hB7-1	1	M..GHTRRQGTSPKCPYLNFFQLLV.LAGLSHFCSGV.IHVTKEVKEVA	46
hB7-2	1	M....DPQCTMGLSN.....ILFVMAFLLSGA...APLKIQAIFYNETA	36
mB7	1	MACNCQLMQDTPLLKFPCLILLFVLLIRLSQVSSDVDEQLSKSVKDKV	50
hB7-1	47	TLSCGHNVSVEE.LAQTRIYWQKEKKMVL.T.MMSGDMNI...WPEYKNRT	91
hB7-2	37	DLPCQFANSQNSLSELVFWQDQENLVNEVYLGKEKFDVHSHKYMGR	86
mB7	51	LLPCRY.NSPHEDESEDRIYWQKHKDVLS.VIAGKLV...WPEYKNRT	95
hB7-1	92	IFDITNNLSIVILALRPSDEGTYESCVVLKYEKDAFKREHLAEVTLSVKAD	141
hB7-2	87	SFD.SDSWTLRLHNLQIKDKGLYQCI.IHHKKPTGMIRIHOMNSELSVLN	135
mB7	96	LYDNNT.YSLIILGLVLSDRGTYSVCVQKKERGTYEVKHLALVKLSIKAD	144
hB7-1	142	FPTPSISDFEIPTSNI.RRIICSTSGGFPEPH....LSWLENGEELNAIN	186
hB7-2	136	FSQPEIVPISNITENVYNLTCSSIHGYPEPKMSVLLRTKNSTIEYDGI	185
mB7	145	FSTPNITESGNPSADT.KRITCFASGGFPKPR....FSWLENGRELPGIN	189

17/34

FIG. 13B

hB7-1	187	TTVSQDPETELYAVSSKLDEN...MTNHSFMCLIKYGHLRVNQTFNWNT	233
hB7-2	186	MQKSQDNVTELYDVSLVSFPDVTSNMTIFCILETDKTRLLSSPFSIE	235
mB7	190	TTISQDPESELYTISSQLDEN...TRNHTIKCLIKYGDHVSSEDFTWK	236
		
hB7-1	234	TKQEHF.PDNLPSWAITLISVNGIFVICCLTYCFAPRCRERRRNERLRR	282
hB7-2	236	.LEDPPPPDHIPWITAVLP....TVIICVMVFCILILWKWKKKRPRNSY	280
mB7	237	PPEDPPDSKNTLVLFAGAGFVITVVIVVVIKCFCKHRSCFRNEA.SR	285
		
hB7-1	283	ESVRPV*	288
hB7-2	281	KCG...TNTMERESEQTKKREKIHIPERSDEAQRVFKSSKTSSCDKSDT	327
mB7	286	ETNNSLTFGPEEALAEQTVFL*	306
		
hB7-2	328	CG*	329

18/34

FIG. 14A

```

1  CCCACGGTCCGGAGCAAGCAGACGCGTAAGAGTGGCTCCTGTAGGCAGCACGGACTTG  60
   -----+-----+-----+-----+-----+-----+
   GGGTGGCGCAGGCCCTCGTTGCTGCGCATTCTCACCAGGACATCCGTCGTGCCTGAAC

61  AACAAACCAGACTCCTGTAGACGTGTTCCAGAACTTACGGAAGCACCCACGATGGACCCCA  120
   -----+-----+-----+-----+-----+-----+
   TTGTTGGTCTGAGGACATCTGCAAGGTCTTGAATGCCCTTCGTGGGTGCTACCTGGGGT

   M D P R -

121  GATGCACCATGGGCTTGGCAATCCTTATCTTTGTGACAGTCTTGCTGATCTCAGATGCTG  180
   -----+-----+-----+-----+-----+-----+
   CTACGTGGTACCCGAACCGTTAGGAATAGAAACACTGTTCAGAACGACTAGAGTCTACGAC

   C T M G L A I L I F V T V L L I S D A V -

181  TTTCGGTGAGACGCAAGCTTATTCAATGGGACTGCATATCTGCCGTGCCATTACAA  240
   -----+-----+-----+-----+-----+-----+
   AAAGGCACCTCTGCGTTCGAATAAAGTTACCCCTGACGTATAGACGGCACGGGTAAATGTT

   S V E T Q A Y F N G T A Y L P C P F T K -

241  AGGCTCAAAACATAAGCCTGAGTGAGCTGGTAGTATTTTGGCAGGACCAGCAAAAGTTGG  300
   -----+-----+-----+-----+-----+-----+
   TCCGAGTTTGTATTTCGGACTCACTCGACCATCATAAACCGTCTGCTCGTTTCAACC

   A Q N I S L S E L V V F W Q D Q Q K L V -

```

FIG. 14B

```

301 TTCTGTACGAGCACTATTGGGCACAGAGAACTTGATAGTGTGAATGCCAAGTACCTGG
-----+-----+-----+-----+-----+
AAGACATGCTCGTGATAAACCCGTGTCTCTTTGAACTATCACACTTACGGTTCATGGACC
360
L Y E H Y L G T E K L D S V N A K Y L G -
GCCGCACGAGCTTTGACAGGAACAACCTGGACTCTACGACTTCACAATGTTCAAGATCAAGG
361 -----+-----+-----+-----+-----+
CGCGTGCTCGAAACTGTCTCTTGTGACCTGAGATGCTGAAGTGTACAAGTCTAGTTCC
420
R T S F D R N N W T L R L H N V Q I K D -
ACATGGGCTCGTATGATTGTTTATACAAAAAAGCCACCACAGGATCAATTATCCTCC
421 -----+-----+-----+-----+-----+
TGTAACCGAGCATACTAACAAAAATATGTTTTTTTCGGTGGTGTCTAGTTAATAGGAGG
480
M G S Y D C F I Q K K P P T G S I I L Q -
AACAGACATTAAACAGAACTGTGAGTATCGCCAACTTCAGTGAACTGAAATAAAACTGG
481 -----+-----+-----+-----+-----+
TTGTCTGTAAATTGTCTTGACAGTCACTAGCGGTTGAAGTCACCTTGGACTTTATTTTGACC
540
Q T L T E L S V I A N F S E P E I K L A -
CTCAGAATGTAACAGGAAATTCTGGCATAAAATTGACCTGCACGCTTAAGCAAGGTCACC
541 -----+-----+-----+-----+-----+
GAGTCTTACATTGTCCCTTAAGACCGTATTTAAACTGGACGTGCAGATTTCGTTCCAGTGG
600
Q N V T G N S G I N L T C T S K Q G H P -

```


20/34

FIG. 14C

```

CGAAACCTAAGAAGATGATTTTCTGATAAATAAATCAACTAATGAGTATGGTGATAACA
601 -----+-----+-----+-----+-----+-----+-----+
GCTTTGGATTCTTCTACATAAAGACTATTGATTAGTTGATTACTCATACCACCTATTGT
660 -----+-----+-----+-----+-----+-----+-----+

K P K K M Y F L I T N S T N E Y G D N M -

TGCAGATATCACAGAATAATGTCACAGAACTGTTTCAGTATCTCCAACAGCCTCTCTCTTT
661 -----+-----+-----+-----+-----+-----+-----+
ACGTCATAGTGTCTATTACAGTGTCTTGACAAGTCATAGAGGTTGTCGGAGAGAGAAA
720 -----+-----+-----+-----+-----+-----+-----+

Q I S Q D N V T E L F S I S N S L S L S -

CATTCCCGGATGGTGTGGCATATGACCGTTGTGTGTGTCTGGAAACGGAGTCAATGA
721 -----+-----+-----+-----+-----+-----+-----+
GTAAGGGCCTACACACACCGTATACTGGCAACACACACACAGACCTTTGCCCTCAGTTACT
780 -----+-----+-----+-----+-----+-----+-----+

F P D G V W H M T V V C V L E T E S M K -

AGATTTCCTCCAAACCTCTCAATTTCACTCAAGAGTTTCCCATCTCCTCAAACGATTGGA
781 -----+-----+-----+-----+-----+-----+-----+
TCTAAGGAGGTTTGGAGAGTTAAAGTGAGTTCTCAAAAGGTAGAGGAGTTTGCATAAACCT
840 -----+-----+-----+-----+-----+-----+-----+

I S S K P L N F T Q E F P S P Q T Y W K -

AGGAGATTACAGCTTCAGTTACTGTGGCCCTCCTCTTGTGTGATGCTGCTCATATTGTAT
841 -----+-----+-----+-----+-----+-----+-----+
TCCTCTAATGTCGAAGTCAATGACACCGGGAGGAGGAACACTACGACGAGTAGTAACATA
900 -----+-----+-----+-----+-----+-----+-----+

E I T A S V T V A L L L V M L L I I V C -

```

21/34

FIG. 14D

```

901  GTCAAGAAGCCGAATCAGCCTAGCAGGCCCCAGCAACACAGCCTCTAAGTTAGAGCGGG
      -----+-----+-----+-----+-----+
      CAGTGTCTTCGGCTTAGTCGGATCGTCCGGGTCGTTGTGTCGGAGATTCAATCTCGCCC
      -----+-----+-----+-----+-----+
      H K K P N Q P S R P S N T A S K L E R D -
      -----+-----+-----+-----+-----+
961  ATAGTAACGCTGACAGAGAGACTATCAACCTGAAGGAACCTGAACCCCAAAATGCTTCAG
      -----+-----+-----+-----+-----+
      TATCATGCGACTGTCTCTGTATAGTTGGACTTCCTTGAACCTTGGGGTTTAAACGAAGTC
      -----+-----+-----+-----+-----+
      S N A D R E T I N L K E L E P Q I A S A -
      -----+-----+-----+-----+-----+
1021 CAAACCAAAATGCAGAGTGAAGGCAGTGAGAGCCTGAGGAAAGAGTTAAAAATGCTTTG
      -----+-----+-----+-----+-----+
      GTTTTGGTTTACGTCCTCACTCCGTCACCTCTCGGACTCCTTCTCAATTTTAAACGAAAC
      -----+-----+-----+-----+-----+
      K P N A E +
      -----+-----+-----+-----+-----+
1081 CCTGAAATAAGAAGTGCAGAGTTTCTCAGAATTCAAAAATGTTCTCAGCTGATGGAATT
      -----+-----+-----+-----+-----+
      GGACTTTATTCTTCACGTCCTCAAAGAGTCTTAAGTTTTTACAAGAGTCGACTAACCTTAA
      -----+-----+-----+-----+-----+
      CTACAGTTGAATAATTAAGAAC
      -----+-----+-----+-----+-----+
1141 GATGCAACTTATTAATTCTTG
      -----+-----+-----+-----+-----+
      1163
```

22/34




FIG. 15

SUBSTITUTE SHEET (RULE 26)

23/34

FIG. 16A





FIG. 16B



SUBSTITUTE SHEET (RULE 26)

FIG. 17B




FIG. 17C




FIG. 17A




FIG. 18B




FIG. 18C




FIG. 18A




FIG. 19A






FIG. 19C



27/34




Figure 20

28/34








Figure 21



SUBSTITUTE SHEET (RULE 26)

29/34




FIG. 22

30/34




Figure 23




Figure 24

32/34




Figure 25

33/34




Figure 26

34/34




Figure 27

SUBSTITUTE SHEET (RULE 26)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/705, C12N 15/62, C07K 19/00, A61K 38/17, 47/48, C12N 5/10, C07K 16/28, 17/14, G01N 33/545	A3	(11) International Publication Number: WO 96/40915 (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09052 (22) International Filing Date: 6 June 1996 (06.06.96) (30) Priority Data: 08/479,744 7 June 1995 (07.06.95) US (71) Applicants: DANA-FARBER CANCER INSTITUTE [US/US]; 44 Binney Street, Boston, MA 02115 (US). REPLIGEN CORPORATION [US/US]; One Kendall Square, Cambridge, MA 02139 (US). (72) Inventors: FREEMAN, Gordon, J.; 305 Walnut Street, Brookline, MA 02146 (US). NADLER, Lee, M.; 36 Cross Hill Road, Newton, MA 02159 (US). GRAY, Gary, S.; 32 Milton Road, Brookline, MA 02146 (US). (74) Agents: MANDRAGOURAS, Amy, E. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 23 January 1997 (23.01.97)	
(54) Title: NOVEL CTLA4/CD28 LIGANDS AND USES THEREFOR (57) Abstract Nucleic acids encoding novel CTLA4/CD28 ligands which costimulate T cell activation are disclosed. In one embodiment, the nucleic acid has a sequence which encodes a fusion protein consisting of at least part of the B lymphocyte antigen, B7-2, fused to one or more constant domains of immunoglobulin heavy chain. Preferably, the nucleic acid is a DNA molecule comprising at least a portion of a nucleotide sequence shown in Figure 8, SEQ ID NO:1 or Figure 14, SEQ ID NO:22. The nucleic acid sequences of the invention can be integrated into various expression vectors, which in turn direct the synthesis of the corresponding proteins or peptides in a variety of hosts, particularly eukaryotic cells, such as mammalian and insect cell culture. Also disclosed are host cells transformed to produce proteins or peptides encoded by the nucleic acid sequences of the invention and isolated proteins and peptides which comprise at least a portion of a novel B lymphocyte antigen. Proteins and peptides described herein can be administered to subjects to enhance or suppress T cell-mediated immune responses.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/09052

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 C12N15/62 C07K19/00 A61K38/17
A61K47/48 C12N5/10 C07K16/28 C07K17/14 G01N33/545

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,95 03408 (DANA FARBER CANCER INST INC ;REPLIGEN CORP (US)) 2 February 1995	1-44,48, 49, 51-59, 61-64 38-50,60
Y	see page 4, line 27 - page 5, line 29 see page 14, line 29 - page 17, line 37 see page 36, line 15 - page 43, line 3 see page 83 - page 90; table VI see page 94, line 10 - page 95, line 10 see claims; figures 8,15 --- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

27 November 1996

Date of mailing of the international search report

06.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Macchia, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 96/09052

C.(Continuation) D. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 181, no. 4, April 1995, pages 1345-1355, XP002018930 YONG GUO ET AL.: "Mutational analysis and an alternatively spliced product of B7 defines its CD28/CTLA4-binding site on Immunoglobulin C-like domain" see page 1346 see page 1348; figure 1B see page 1351</p> <p>---</p>	38-42
Y	<p>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 200, no. 1, 15 April 1994, pages 443-449, XP002018931 INOBE ET AL.: "Identification of an alternatively spliced form of the murine homologue of B7" see figures 1C,D</p> <p>---</p>	38-50,60
A	<p>THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 173, no. 3, March 1991, pages 721-730, XP000611265 LINSLEY ET AL.: "Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and Interleukin 2 mRNA accumulation" cited in the application see the whole document</p> <p>---</p>	1-8, 13-19, 28-33, 63,64
A	<p>THE JOURNAL OF IMMUNOLOGY, vol. 154, 1 January 1995, pages 97-105, XP002018932 LANIER ET AL.: "CD80 (B7) and CD86 (B70) provide similar costimulatory signals for T cell proliferation, cytokine production and generation of CTL" see abstract</p> <p>---</p>	63,64
A	<p>WO,A,94 12520 (ENZON INC) 9 June 1994 see page 11, line 7 - page 12, line 5; claims 1,7,8</p> <p>---</p>	46,50,60
A	<p>INTERNATIONAL IMMUNOLOGY, vol. 4, no. 2, 1992, pages 215-225, XP002019178 CLASSON ET AL.: "The hinge region of the CD8 alpha chain: structure, antigenicity and utility in expression of immunoglobulin superfamily domains" see abstract; figure 1</p> <p>-----</p>	46,50,60

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/09052

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 28-37 and 63,64
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although these claims (as far as in vivo methods are concerned) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 96/09052

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9503408	02-02-95	AU-A- 7405294 EP-A- 0711345	20-02-95 15-05-96
WO-A-9412520	09-06-94	AU-A- 5670194	22-06-94